

CISC-102
Fall 2016
Week 1

David Rappaport daver@cs.queensu.ca

Goodwin G-532

Office Hours:

Monday 1:00-3:00 (or by appointment)

Homework

- Homework every week. Keep up to date or you risk falling behind.
- Homework will be solved in class on due date.
- Homework is not handed in, and not graded.
- Quizzes and Final exam are based on homework questions.

Assessment

- 4 quizzes, 17% each
- 1 final 32%
- You may miss up to two quizzes with my explicit permission.
If you miss quizzes then the marking scheme will be revised for you as follows:
1 missed - 3 quizzes 20% each and 40% final.
2 missed - 2 quizzes 25%, and 50% final
- You must pass the final (obtain at least 50%) to pass the course.
- Attending class is not compulsory, and I will not take attendance.

However...


2. Do you have any specific suggestions for improvements to this course?

Somehow get the students that don't come to class to come: because they are seriously MISSING out.

Less proofs? lol.

Motivation

- According to a *fairly* recent poll (33,000 CDN students interviewed in 2014) the preferred employer is Google.
- I have been given a communication from an applicant to Google with tips on how to conduct an interview.
- *#1 tip Algorithm Complexity: You need to know Big-O. If you struggle with basic big-O complexity analysis, then you are almost guaranteed not to get hired.*
- Mastering discrete mathematics is the direct pre-requisite to mastering algorithms and complexity.
- You should view this course as a language course. You will be learning the language of mathematics and computing!
- Math can be fun.
- Math is beautiful!



Beauty

- The picture on the previous page is a work of art titled “Beauty”.
(Prints can be purchased on-line.)
- The equation

$$e^{i\pi} + 1 = 0$$

consists of the most important concepts in mathematics:

- numbers
 - 0, 1 (integers)
 - π , e (irrational real numbers)
 - i (a complex number)
- operations
 - $+$ \times and exponentiation (exp. function)
- and the relation =

Motivation

- Math is a human invention just like music, painting, sculpture, poetry, hockey, basketball, soccer, fishing ...

And how do you become proficient at music, hockey, fishing ... ?

Practice, practice, practice.

**The Perfect Introductory Problem:
Counting hand shakes**

Alice is having a birthday party at her house, and has invited Bob, Carl, Diane, Eve, Frank, and George.

They all shake hands with each other.

Q: How many handshakes are there?

George says, “ I know the answer and I can prove it to you. There are 7 of us, so I shake hands with 6 other people. That’s also true for everyone else. So the total number of hand shakes is $6 \times 7 = 42$.”

Frank says, “ I have another way of working this out. Suppose there’s only two of us, just George and I. That’s 1 handshake. For 3 of us Eve, Frank and George, we have

E and F shake hands

E and G shake hands

F and G shake hands. 3 hand shakes.

And for 4 of us, Diane, Eve, Frank, and George we have

D and E shake hands

D and F shake hands

D and G shake hands

E and F shake hands

E and G shake hands

F and G shake hands. 6 hand shakes.

I see the pattern $(3 \times 2) / 2 = 3$,

$$(4 \times 3) / 2 = 6.$$

So with 7 of us the correct answer is

$$(6 \times 7) / 2 = 21.$$

Who’s right?

Sets

- We convert the hand shake problem into an “official” math problem using proper notation.
- The basic building block will be the set.
- A set is a collection of distinct elements.

Examples

$$A = \{1, 3, 5, 7, 9\},$$

$$B = \{x \mid x \text{ is an integer, } 0 \leq x < 10\}$$

$$C = \{x : x \text{ is an odd integer, } 0 < x < 10\}$$

$$A \subseteq C \text{ (A is a subset of C)}$$

$$C \subseteq A \text{ (C is a subset of A)}$$

$$A = C \text{ (A and C are equal, that is the elements of A and C are the same.)}$$

$$A \subseteq B \text{ (A is a subset of B)}$$

$$B \not\subseteq A \text{ (B is not a subset of A)}$$

$$A \subset B \text{ (A is a proper subset of B)}$$

$$B \not\subset A \text{ (B is not a proper subset of A)}$$

$$1 \in A \text{ (1 is an element of A)}$$

$$\{1\} \subseteq A$$

$$\{1\} \subset A$$

Sets can have infinitely many elements

\mathbb{N} = the set of *natural numbers*: 1, 2, 3, . . .

\mathbb{Z} = the set of all integers: . . ., -2, -1, 0, 1, 2, . . .

\mathbb{Q} = the set of rational numbers

\mathbb{R} = the set of real numbers

\mathbb{C} = the set of complex numbers

Observe that $\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}$.

U : All sets under investigation in any application of set theory are assumed to belong to some fixed large set called the *universal set*.

\emptyset : A set with no elements is called the *empty set* or *null set* .

For any set A , we have: $\emptyset \subseteq A \subseteq U$

The handshake problem

Let $S = \{a,b,c,d,e,f,g\}$ denote the set of party goers, and a handshake can be represented as a two element subset of S . (For example $\{a,b\}$ denotes the handshake between Alice and Bob.)

Q. How many two element subsets are there of the set S ?

Generalizing the handshake problem

Suppose that S is a set consisting of n elements.

Q. How many two element subsets are there of the set S ?

The hand shake problem seems frivolous but it is actually a representation of an important mathematical concept. For example if we wanted to know which handshake was the “best” we would have to compare $n(n-1)/2$ of them. Let $n = 35,000,000$ (the population of Canada) we would have to compare 612,499,982,500,000 or roughly 612 trillion hand shakes. (Too much!)

If we test one handshake per second it would take roughly 31,688 Years, 269 Days, 1 Hour. (Too long!)

Problem from SN

1.26 Which of the following sets are equal?

$$A = \{x \mid x^2 - 4x + 3 = 0\},$$

$$B = \{x \mid x^2 - 3x + 2 = 0\},$$

$$C = \{x \mid x \in \mathbb{N}, x < 3\},$$

$$D = \{x \mid x \in \mathbb{N}, x \text{ is odd}, x < 5\},$$

$$E = \{1, 2\},$$

$$F = \{1, 2, 1\},$$

$$G = \{3, 1\},$$


$$H = \{1, 1, 3\}.$$

NOTE: To determine the elements of sets A and B, you need to be able to *factor quadratic equations*. This is a topic that you may or may not be familiar with. For this course it is assumed that you are able to do this factoring or pick it up. All examples that you will see in this course will have integer solutions. Here's a link to a web page with some good tips for factoring quadratic equations: <https://www.mathsisfun.com/algebra/factoring-quadratics.html>

Graph of $x^2 - 4x + 3$.

The function crosses the x -axis at two points $x = 1$, and $x = 3$.


Note: $x^2 - 4x + 3 = (x - 1)(x - 3)$.



Graph of $x^2 - 3x + 2$.

The function crosses the x -axis at two points $x = 1$, and $x = 2$.

Note: $x^2 - 4x + 3 = (x - 1)(x - 2)$.



Sets**Definition: (From Schaum's Notes)**

A set may be viewed as any well-defined collection of objects, called the elements or *members* of the set.

This sentence defines in a mathematical sense the term set and the term element.

Key things to remember about sets.

- Always use curly braces { }.
- The elements are *well-defined*, that is, each element can be distinguished from another.
- A set is an *un-ordered* collection of elements.

Notation

$A = \{ 1, 2, 3 \}$ is a set of 3 elements.

$1 \in A$ (1 is an element of the set A.)

$B = \{1,3,2\}$ implies that $A = B$.

Subset

Let A and B be two sets, where every element of A is also an element of B .

For example:

$A = \{\text{red, black}\}$, $B = \{\text{red, black, green}\}$.

Let a be a symbol that denotes an arbitrary object.

Observe that: if $a \in A$ then $a \in B$.

We can say that A is contained in B , or A is a subset of B .

Definition: Let X and Y be two sets such that $a \in X$ implies $a \in Y$. We then can say that X is a *subset* of Y , and notate it as $X \subseteq Y$.

Suppose X and Y are two sets such that:

$X \subseteq Y$ and $Y \subseteq X$.

That means $a \in X$ implies $a \in Y$

and $a \in Y$ implies $a \in X$. So in fact the sets are equal.

Definition: Let X and Y be two sets.

If $X = Y$ then $X \subseteq Y$ and $Y \subseteq X$.

And if $X \subseteq Y$ and $Y \subseteq X$ then $X=Y$.

These two sentences can be expressed in a single sentence as:

$X = Y$ if and only if $X \subseteq Y$ and $Y \subseteq X$.

Definition: Let X and Y be two sets.

If $X \subseteq Y$ and $X \neq Y$ then we say that

X is a *proper subset* of Y , and notate it as:

$X \subset Y$.

Another way to say this is: X is a proper subset of Y if every element of X is also an element of Y and there exists at least one element of Y that is not an element of X .

Find the definitions and examples in SN for the symbols.

$\not\subseteq$ (not a subset) $\not\subset$ (not a proper subset)

\supseteq (superset) $\not\supseteq$ (not a superset)

\supset (proper superset) $\not\supset$ (not a proper superset)

Disjoint Sets

Let A and B be two sets. If A and B have no elements in common then we say that they are *disjoint*.

Using subset notation we can say that if A and B are disjoint then $A \not\subseteq B$ and $B \not\subseteq A$.

However, if $A \not\subseteq B$ and $B \not\subseteq A$ then A and B may not be disjoint.

U : All sets under investigation in any application of set theory are assumed to belong to some fixed large set called the *universal set*.

\emptyset : A set with no elements is called the *empty set* or *null set*.

The empty set is a subset of every set, and the universal set is a superset of every set.

Using symbols the blue sentence can be expressed as follows:

For any set A , we have: $\emptyset \subseteq A \subseteq U$

Examples

Consider the set $A = \{1, 2, 3\}$.

A is a set consisting of 3 elements.

$\{1\} \subseteq A$, ($\{1\}$ is a subset of A)

$\{1\} \subset A$, ($\{1\}$ is a proper subset of A)

$1 \in A$ (1 is an element of A)

$\{1,2,3\} \subseteq A$

$\{1,2,3\} \not\subseteq A$

$\{1,2\} \subset A$

$\emptyset \subseteq A$ and $\emptyset \subset A$

$A \subseteq \mathbb{N}$ and $A \subset \mathbb{N}$

Examples:

People in a room.

Coins in your pocket.

Note: If you have two (or more) quarters in your pocket then you need to be able to distinguish one from the other if you want to consider the coins as a set. If you have no coins in your pocket then the set of coins in your pockets is the empty set.

Seating Arrangements

There is a large table at the party and Alice wants to experience every possible seating arrangement. How many ways can 7 people sit at a table?



This “seating arrangement question” is equivalent to asking for the number of different ways to order 7 people.

Number of ways to order 1 person? 1.

Number of ways to order 2 people?

(1,2) (2,1). 2×1

Number of ways to order 3 people?

(3,1,2)(3,2,1)(1,3,2)(2,3,1)(1,2,3)(2,1,3). $3 \times 2 \times 1$

Number of ways to order 4 people?

Guess: $4 \times 3 \times 2 \times 1 = 4!$

Permutations

There are $n!$ ways to order n distinct objects

Recall $n!$ (n factorial) is given by the expression:

$$n! = n \times (n-1) \times (n-2) \times \dots \times 1$$

Selection with ordering

How many ways are there to pick 7 people out of a class of 70 and seat them into 7 numbered chairs? (Selection with ordering.)

1st pick has 70 choices.

2nd pick has 69 choices.

3rd pick has 68 choices.

4th pick has 67 choices.

5th pick has 66 choices.

6th pick has 65 choices.

7th pick has 64 choices.

So the number of ways to pick 7 people out of a class of 70 is:

$$70 \times 69 \times 68 \times 67 \times 66 \times 65 \times 64 = 70!/63!$$

Lottery Tickets

Lotto 6-49, choose 6 numbers from 1 to 49.

How many ways are there to choose to these numbers?

A very simplified version of this game is Lotto 1- 49, where players choose 1 number from 49. There are 49 choices.

Note that the probability (the odds) of winning Lotto 1- 49 is $1/49$. (one choice divided by the total number of choices)

Consider Lotto 2 - 49, where you have to pick 2 numbers from 49.

A tempting but wrong guess would be 49×48 choices.

Suppose choice 1 is 42, and choice 2 is 18. That is equivalent to choice 1 is 18 and choice 2 is 42, so 49×48 double counts all possibilities.

The actual answer is $49 \times 48 / 2!$.

Combinations (Selection without ordering)

For lotto 6-49, players choose 6 numbers from 1 to 49.

How many ways are there to choose to these numbers?

Solution:

$$49 \times 48 \times 47 \times 46 \times 45 \times 44 / 6! = 13,983,816.$$

This can also be written as:

and pronounced 49 choose 6.

$$\binom{49}{6} = \frac{49!}{43!6!}$$

What is the probability that any single choice is the winning number?

$$1/13,983,816$$

The current price of a Lotto 6-49 card is \$3.

The “fair” prize for choosing the winning numbers should be \$41,951,448.

The minimum jackpot is \$5,000,000.

There have been occurrences when the Lotto 6-49 jackpot exceeded the fair prize.

The largest jackpot (as of Aug. 2016) was \$64,000,000 on October 17, 2015.

Set Operators

Operators on sets are

union \cup and intersection \cap .

Definitions:

$$A \cup B = \{x \mid x \in A \text{ or } x \in B\}$$

$$A \cap B = \{x \mid x \in A \text{ and } x \in B\}$$

Logical Operators

$$p \wedge q \quad \text{pronounced } p \text{ and } q$$

Both p and q have to be true for the compound proposition p and q to be true.

$$p \vee q \quad \text{pronounced } p \text{ or } q$$

At least one of p or q must be true for the compound proposition p or q to be true.

We can rewrite our definition for set union and set intersection using logical operators as follows:

$$A \cup B = \{x : x \in A \vee x \in B\}$$

$$A \cap B = \{x : x \in A \wedge x \in B\}$$


For example:

Suppose A is the set of guitars and B is the set of red musical instruments.

- An element x is in the set of A union B if it is a guitar or if it is a red musical instrument.
- An element of x is in the set of A intersection B if x is red and x is a guitar.

Venn Diagrams


Useful for providing intuitive insight.



Note the rectangle surrounding the circles denotes the Universe U .


The complement of a set A written A^c is defined as:

$$A^c = \{x \mid x \notin A\}$$




The relative complement of a set B with respect to A , sometimes called the difference

$$A \setminus B = \{x \mid x \in A, x \notin B\}$$




(The relative complement is sometimes written as $A - B$.)

$$A \cup B = \{x : x \in A \vee x \in B\}$$



$$A \cap B = \{x : x \in A \wedge x \in B\}$$




The symmetric difference of sets A and B :

The symmetric difference consists of elements that are in A or in B but not in both.

$$A \oplus B = (A \cup B) \setminus (A \cap B) \quad \text{or} \quad A \oplus B = (A \setminus B) \cup (B \setminus A)$$

$$A \oplus B = (A \cup B) \setminus (A \cap B)$$



$$A \oplus B = (A \setminus B) \cup (B \setminus A)$$

