

Lemmas and Proofs for “Complete and Easy Bidirectional Typechecking for Higher-Rank Polymorphism”

Jana Dunfield

Neelakantan R. Krishnaswami

June 2013*

Contents

A	Declarative Subtyping	6
A.1	Properties of Well-Formedness	6
1	Proposition (Weakening)	6
2	Proposition (Substitution)	6
A.2	Reflexivity	6
3	Lemma (Reflexivity of Declarative Subtyping)	6
A.3	Subtyping Implies Well-Formedness	6
4	Lemma (Well-Formedness)	6
A.4	Substitution	6
5	Lemma (Substitution)	6
A.5	Transitivity	6
6	Lemma (Transitivity of Declarative Subtyping)	6
A.6	Invertibility of $\leq_{\forall R}$	6
7	Lemma (Invertibility)	6
A.7	Non-Circularity and Equality	6
1	Definition (Subterm Occurrence)	6
8	Lemma (Occurrence)	6
9	Lemma (Monotype Equality)	6
2	Definition (Contextual Size)	6
B	Type Assignment	7
10	Lemma (Well-Formedness)	7
1	Theorem (Completeness of Bidirectional Typing)	7
11	Lemma (Subtyping Coercion)	7
12	Lemma (Application Subtyping)	7
2	Theorem (Soundness of Bidirectional Typing)	7
C	Robustness of Typing	7
13	Lemma (Type Substitution)	7
3	Definition (Context Subtyping)	7
14	Lemma (Subsumption)	7
3	Theorem (Substitution)	7
4	Theorem (Inverse Substitution)	7
5	Theorem (Annotation Removal)	8

*Recompiled in 2020 to correct the name of the first author.

D	Properties of Context Extension	8
D.1	Syntactic Properties	8
15	Lemma (Declaration Preservation)	8
16	Lemma (Declaration Order Preservation)	8
17	Lemma (Reverse Declaration Order Preservation)	8
18	Lemma (Substitution Extension Invariance)	8
19	Lemma (Extension Equality Preservation)	8
20	Lemma (Reflexivity)	8
21	Lemma (Transitivity)	8
4	Definition (Softness)	8
22	Lemma (Right Softness)	8
23	Lemma (Evar Input)	8
24	Lemma (Extension Order)	8
25	Lemma (Extension Weakening)	9
26	Lemma (Solution Admissibility for Extension)	9
27	Lemma (Solved Variable Addition for Extension)	9
28	Lemma (Unsolved Variable Addition for Extension)	9
29	Lemma (Parallel Admissibility)	9
30	Lemma (Parallel Extension Solution)	9
31	Lemma (Parallel Variable Update)	9
D.2	Instantiation Extends	9
32	Lemma (Instantiation Extension)	9
D.3	Subtyping Extends	9
33	Lemma (Subtyping Extension)	9
E	Decidability of Instantiation	9
34	Lemma (Left Unsolvedness Preservation)	9
35	Lemma (Left Free Variable Preservation)	9
36	Lemma (Instantiation Size Preservation)	9
7	Theorem (Decidability of Instantiation)	9
F	Decidability of Algorithmic Subtyping	10
F.1	Lemmas for Decidability of Subtyping	10
37	Lemma (Monotypes Solve Variables)	10
38	Lemma (Monotype Monotonicity)	10
39	Lemma (Substitution Decreases Size)	10
40	Lemma (Monotype Context Invariance)	10
F.2	Decidability of Subtyping	10
8	Theorem (Decidability of Subtyping)	10
G	Decidability of Typing	10
9	Theorem (Decidability of Typing)	10
H	Soundness of Subtyping	10
H.1	Lemmas for Soundness	10
41	Lemma (Uvar Preservation)	10
42	Lemma (Variable Preservation)	10
43	Lemma (Substitution Typing)	10
44	Lemma (Substitution for Well-Formedness)	11
45	Lemma (Substitution Stability)	11
46	Lemma (Context Partitioning)	11
47	Lemma (Softness Goes Away)	11
48	Lemma (Filling Completes)	11
49	Lemma (Stability of Complete Contexts)	11
50	Lemma (Finishing Types)	11
51	Lemma (Finishing Completions)	11

52	Lemma (Confluence of Completeness)	11
H.2	Instantiation Soundness	11
10	Theorem (Instantiation Soundness)	11
H.3	Soundness of Subtyping	11
11	Theorem (Soundness of Algorithmic Subtyping)	11
I	Typing Extension	11
54	Lemma (Typing Extension)	11
J	Soundness of Typing	11
12	Theorem (Soundness of Algorithmic Typing)	11
K	Completeness of Subtyping	12
K.1	Instantiation Completeness	12
13	Theorem (Instantiation Completeness)	12
K.2	Completeness of Subtyping	12
14	Theorem (Generalized Completeness of Subtyping)	12
L	Completeness of Typing	12
15	Theorem (Completeness of Algorithmic Typing)	12
	Proofs	13
A'	Declarative Subtyping	13
1	Proof of Proposition (Weakening)	13
2	Proof of Proposition (Substitution)	13
A'.1	Properties of Well-Formedness	13
A'.2	Reflexivity	13
3	Proof of Lemma (Reflexivity of Declarative Subtyping)	13
A'.3	Subtyping Implies Well-Formedness	13
4	Proof of Lemma (Well-Formedness)	13
A'.4	Substitution	13
5	Proof of Lemma (Substitution)	13
A'.5	Transitivity	14
6	Proof of Lemma (Transitivity of Declarative Subtyping)	14
A'.6	Invertibility of $\leq \forall R$	15
7	Proof of Lemma (Invertibility)	15
A'.7	Non-Circularity and Equality	16
8	Proof of Lemma (Occurrence)	16
9	Proof of Lemma (Monotype Equality)	16
B'	Type Assignment	17
10	Proof of Lemma (Well-Formedness)	17
1	Proof of Theorem (Completeness of Bidirectional Typing)	17
11	Proof of Lemma (Subtyping Coercion)	18
12	Proof of Lemma (Application Subtyping)	19
2	Proof of Theorem (Soundness of Bidirectional Typing)	19
C'	Robustness of Typing	21
13	Proof of Lemma (Type Substitution)	21
14	Proof of Lemma (Subsumption)	21
3	Proof of Theorem (Substitution)	24
4	Proof of Theorem (Inverse Substitution)	24
5	Proof of Theorem (Annotation Removal)	26

D'	Properties of Context Extension	27
D'.1	Syntactic Properties	27
15	Proof of Lemma (Declaration Preservation)	27
16	Proof of Lemma (Declaration Order Preservation)	27
17	Proof of Lemma (Reverse Declaration Order Preservation)	28
18	Proof of Lemma (Substitution Extension Invariance)	29
19	Proof of Lemma (Extension Equality Preservation)	30
20	Proof of Lemma (Reflexivity)	31
21	Proof of Lemma (Transitivity)	31
22	Proof of Lemma (Right Softness)	33
23	Proof of Lemma (Evar Input)	33
24	Proof of Lemma (Extension Order)	34
25	Proof of Lemma (Extension Weakening)	36
26	Proof of Lemma (Solution Admissibility for Extension)	36
27	Proof of Lemma (Solved Variable Addition for Extension)	36
28	Proof of Lemma (Unsolved Variable Addition for Extension)	36
29	Proof of Lemma (Parallel Admissibility)	36
30	Proof of Lemma (Parallel Extension Solution)	37
31	Proof of Lemma (Parallel Variable Update)	37
D'.2	Instantiation Extends	37
32	Proof of Lemma (Instantiation Extension)	37
D'.3	Subtyping Extends	39
33	Proof of Lemma (Subtyping Extension)	39
E'	Decidability of Instantiation	39
34	Proof of Lemma (Left Unsolvedness Preservation)	39
35	Proof of Lemma (Left Free Variable Preservation)	40
36	Proof of Lemma (Instantiation Size Preservation)	42
7	Proof of Theorem (Decidability of Instantiation)	43
F'	Decidability of Algorithmic Subtyping	45
F'.1	Lemmas for Decidability of Subtyping	45
37	Proof of Lemma (Monotypes Solve Variables)	45
38	Proof of Lemma (Monotype Monotonicity)	46
39	Proof of Lemma (Substitution Decreases Size)	46
40	Proof of Lemma (Monotype Context Invariance)	47
F'.2	Decidability of Subtyping	47
8	Proof of Theorem (Decidability of Subtyping)	47
G'	Decidability of Typing	48
9	Proof of Theorem (Decidability of Typing)	48
H'	Soundness of Subtyping	49
H'.1	Lemmas for Soundness	49
42	Proof of Lemma (Variable Preservation)	49
43	Proof of Lemma (Substitution Typing)	49
44	Proof of Lemma (Substitution for Well-Formedness)	50
45	Proof of Lemma (Substitution Stability)	51
46	Proof of Lemma (Context Partitioning)	51
49	Proof of Lemma (Stability of Complete Contexts)	51
50	Proof of Lemma (Finishing Types)	53
51	Proof of Lemma (Finishing Completions)	53
52	Proof of Lemma (Confluence of Completeness)	53
H'.2	Instantiation Soundness	53
10	Proof of Theorem (Instantiation Soundness)	53
H'.3	Soundness of Subtyping	55

11	Proof of Theorem (Soundness of Algorithmic Subtyping)	55
I'	Typing Extension	57
54	Proof of Lemma (Typing Extension)	57
J'	Soundness of Typing	58
12	Proof of Theorem (Soundness of Algorithmic Typing)	58
K'	Completeness	63
K'.1	Instantiation Completeness	63
13	Proof of Theorem (Instantiation Completeness)	63
K'.2	Completeness of Subtyping	66
14	Proof of Theorem (Generalized Completeness of Subtyping)	66
L'	Completeness of Typing	71
15	Proof of Theorem (Completeness of Algorithmic Typing)	71

A Declarative Subtyping

A.1 Properties of Well-Formedness

Proposition 1 (Weakening). *If $\Psi \vdash A$ then $\Psi, \Psi' \vdash A$ by a derivation of the same size.*

Proposition 2 (Substitution). *If $\Psi \vdash A$ and $\Psi, \alpha, \Psi' \vdash B$ then $\Psi, \Psi' \vdash [A/\alpha]B$.*

A.2 Reflexivity

Lemma 3 (Reflexivity of Declarative Subtyping). *Subtyping is reflexive: if $\Psi \vdash A$ then $\Psi \vdash A \leq A$.*

A.3 Subtyping Implies Well-Formedness

Lemma 4 (Well-Formedness). *If $\Psi \vdash A \leq B$ then $\Psi \vdash A$ and $\Psi \vdash B$.*

A.4 Substitution

Lemma 5 (Substitution). *If $\Psi \vdash \tau$ and $\Psi, \alpha, \Psi' \vdash A \leq B$ then $\Psi, [\tau/\alpha]\Psi' \vdash [\tau/\alpha]A \leq [\tau/\alpha]B$.*

A.5 Transitivity

Lemma 6 (Transitivity of Declarative Subtyping). *If $\Psi \vdash A \leq B$ and $\Psi \vdash B \leq C$ then $\Psi \vdash A \leq C$.*

A.6 Invertibility of $\leq \forall R$

Lemma 7 (Invertibility).

If \mathcal{D} derives $\Psi \vdash A \leq \forall \beta. B$ then \mathcal{D}' derives $\Psi, \beta \vdash A \leq B$ where $\mathcal{D}' < \mathcal{D}$.

A.7 Non-Circularity and Equality

Definition 1 (Subterm Occurrence).

Let $A \preceq B$ iff A is a subterm of B .

Let $A \prec B$ iff A is a proper subterm of B (that is, $A \preceq B$ and $A \neq B$).

Let $A \rhd B$ iff A occurs in B inside an arrow, that is, there exist B_1, B_2 such that $(B_1 \rightarrow B_2) \preceq B$ and $A \preceq B_k$ for some $k \in \{1, 2\}$.

Lemma 8 (Occurrence).

(i) *If $\Psi \vdash A \leq \tau$ then $\tau \not\prec A$.*

(ii) *If $\Psi \vdash \tau \leq B$ then $\tau \not\prec B$.*

Lemma 9 (Monotype Equality). *If $\Psi \vdash \sigma \leq \tau$ then $\sigma = \tau$.*

Definition 2 (Contextual Size). *The size of A with respect to a context Γ , written $|\Gamma \vdash A|$, is defined by*

$$\begin{aligned} |\Gamma \vdash \alpha| &= 1 \\ |\Gamma[\hat{\alpha}] \vdash \hat{\alpha}| &= 1 \\ |\Gamma[\hat{\alpha} = \tau] \vdash \hat{\alpha}| &= 1 + |\Gamma[\hat{\alpha} = \tau] \vdash \tau| \\ |\Gamma \vdash \forall \alpha. A| &= 1 + |\Gamma, \alpha \vdash A| \\ |\Gamma \vdash A \rightarrow B| &= 1 + |\Gamma \vdash A| + |\Gamma \vdash B| \end{aligned}$$

B Type Assignment

Lemma 10 (Well-Formedness).

If $\Psi \vdash e \Leftarrow A$ or $\Psi \vdash e \Rightarrow A$ or $\Psi \vdash A \bullet e \Rightarrow C$ then $\Psi \vdash A$ (and in the last case, $\Psi \vdash C$).

Theorem 1 (Completeness of Bidirectional Typing).

If $\Psi \vdash e : A$ then there exists e' such that $\Psi \vdash e' \Rightarrow A$ and $|e'| = e$.

Lemma 11 (Subtyping Coercion). If $\Psi \vdash A \leq B$ then there exists f which is $\beta\eta$ -equal to the identity such that $\Psi \vdash f : A \rightarrow B$.

Lemma 12 (Application Subtyping). If $\Psi \vdash A \bullet e \Rightarrow C$ then there exists B such that $\Psi \vdash A \leq B \rightarrow C$ and $\Psi \vdash e \Leftarrow B$ by a smaller derivation.

Theorem 2 (Soundness of Bidirectional Typing). We have that:

- If $\Psi \vdash e \Leftarrow A$, then there is an e' such that $\Psi \vdash e' : A$ and $e' =_{\beta\eta} |e|$.
- If $\Psi \vdash e \Rightarrow A$, then there is an e' such that $\Psi \vdash e' : A$ and $e' =_{\beta\eta} |e|$.

C Robustness of Typing

Lemma 13 (Type Substitution).

Assume $\Psi \vdash \tau$.

- If $\Psi, \alpha, \Psi' \vdash e' \Leftarrow C$ then $\Psi, [\tau/\alpha]\Psi' \vdash [\tau/\alpha]e' \Leftarrow [\tau/\alpha]C$.
- If $\Psi, \alpha, \Psi' \vdash e' \Rightarrow C$ then $\Psi, [\tau/\alpha]\Psi' \vdash [\tau/\alpha]e' \Rightarrow [\tau/\alpha]C$.
- If $\Psi, \alpha, \Psi' \vdash B \bullet e' \Rightarrow C$ then $\Psi, [\tau/\alpha]\Psi' \vdash [\tau/\alpha]B \bullet [\tau/\alpha]e' \Rightarrow [A/\alpha]C$.

Moreover, the resulting derivation contains no more applications of typing rules than the given one. (Internal subtyping derivations, however, may grow.)

Definition 3 (Context Subtyping). We define the judgment $\Psi' \leq \Psi$ with the following rules:

$$\frac{}{\cdot \leq \cdot} \text{CtxSubEmpty} \quad \frac{\Psi' \leq \Psi}{\Psi', \alpha \leq \Psi, \alpha} \text{CtxSubUvar} \quad \frac{\Psi' \leq \Psi \quad \Psi \vdash A' \leq A}{\Psi', x : A' \leq \Psi, x : A} \text{CtxSubVar}$$

Lemma 14 (Subsumption). Suppose $\Psi' \leq \Psi$. Then:

- If $\Psi \vdash e \Leftarrow A$ and $\Psi \vdash A \leq A'$ then $\Psi' \vdash e \Leftarrow A'$.
- If $\Psi \vdash e \Rightarrow A$ then there exists A' such that $\Psi \vdash A' \leq A$ and $\Psi' \vdash e \Rightarrow A'$.
- If $\Psi \vdash C \bullet e \Rightarrow A$ and $\Psi \vdash C' \leq C$ then there exists A' such that $\Psi \vdash A' \leq A$ and $\Psi' \vdash C' \bullet e \Rightarrow A'$.

Theorem 3 (Substitution).

Assume $\Psi \vdash e \Rightarrow A$.

- If $\Psi, x : A \vdash e' \Leftarrow C$ then $\Psi \vdash [e/x]e' \Leftarrow C$.
- If $\Psi, x : A \vdash e' \Rightarrow C$ then $\Psi \vdash [e/x]e' \Rightarrow C$.
- If $\Psi, x : A \vdash B \bullet e' \Rightarrow C$ then $\Psi \vdash B \bullet [e/x]e' \Rightarrow C$.

Theorem 4 (Inverse Substitution).

Assume $\Psi \vdash e \Leftarrow A$.

- If $\Psi \vdash [(e : A)/x]e' \Leftarrow C$ then $\Psi, x : A \vdash e' \Leftarrow C$.

(ii) If $\Psi \vdash [(e : A)/x]e' \Rightarrow C$ then $\Psi, x : A \vdash e' \Rightarrow C$.

(iii) If $\Psi \vdash B \bullet [(e : A)/x]e' \Rightarrow C$ then $\Psi, x : A \vdash B \bullet e' \Rightarrow C$.

Theorem 5 (Annotation Removal). *We have that:*

- If $\Psi \vdash ((\lambda x. e) : A) \Leftarrow C$ then $\Psi \vdash \lambda x. e \Leftarrow C$.
- If $\Psi \vdash (() : A) \Leftarrow C$ then $\Psi \vdash () \Leftarrow C$.
- If $\Psi \vdash e_1 (e_2 : A) \Rightarrow C$ then $\Psi \vdash e_1 e_2 \Rightarrow C$.
- If $\Psi \vdash (x : A) \Rightarrow A$ then $\Psi \vdash x \Rightarrow B$ and $\Psi \vdash B \leq A$.
- If $\Psi \vdash ((e_1 e_2) : A) \Rightarrow A$ then $\Psi \vdash e_1 e_2 \Rightarrow B$ and $\Psi \vdash B \leq A$.
- If $\Psi \vdash ((e : B) : A) \Rightarrow A$ then $\Psi \vdash (e : B) \Rightarrow B$ and $\Psi \vdash B \leq A$.
- If $\Psi \vdash ((\lambda x. e) : \sigma \rightarrow \tau) \Rightarrow \sigma \rightarrow \tau$ then $\Psi \vdash \lambda x. e \Rightarrow \sigma \rightarrow \tau$.

Theorem 6 (Soundness of Eta).

If $\Psi \vdash \lambda x. e \Leftarrow A$ and $x \notin \text{FV}(e)$, then $\Psi \vdash e \Leftarrow A$.

D Properties of Context Extension

D.1 Syntactic Properties

Lemma 15 (Declaration Preservation). *If $\Gamma \longrightarrow \Delta$, and u is a variable or marker $\blacktriangleright_{\hat{\alpha}}$ declared in Γ , then u is declared in Δ .*

Lemma 16 (Declaration Order Preservation). *If $\Gamma \longrightarrow \Delta$ and u is declared to the left of v in Γ , then u is declared to the left of v in Δ .*

Lemma 17 (Reverse Declaration Order Preservation). *If $\Gamma \longrightarrow \Delta$ and u and v are both declared in Γ and u is declared to the left of v in Δ , then u is declared to the left of v in Γ .*

Lemma 18 (Substitution Extension Invariance). *If $\Theta \vdash A$ and $\Theta \longrightarrow \Gamma$ then $[\Gamma]A = [\Gamma](\Theta A)$ and $[\Gamma]A = [\Theta](\Gamma A)$.*

Lemma 19 (Extension Equality Preservation).

If $\Gamma \vdash A$ and $\Gamma \vdash B$ and $[\Gamma]A = [\Gamma]B$ and $\Gamma \longrightarrow \Delta$, then $[\Delta]A = [\Delta]B$.

Lemma 20 (Reflexivity). *If Γ is well-formed, then $\Gamma \longrightarrow \Gamma$.*

Lemma 21 (Transitivity). *If $\Gamma \longrightarrow \Delta$ and $\Delta \longrightarrow \Theta$, then $\Gamma \longrightarrow \Theta$.*

Definition 4 (Softness). *A context Θ is soft iff it consists only of $\hat{\alpha}$ and $\hat{\alpha} = \tau$ declarations.*

Lemma 22 (Right Softness). *If $\Gamma \longrightarrow \Delta$ and Θ is soft (and (Δ, Θ) is well-formed) then $\Gamma \longrightarrow \Delta, \Theta$.*

Lemma 23 (Evar Input).

If $\Gamma, \hat{\alpha} \longrightarrow \Delta$ then $\Delta = (\Delta_0, \Delta_{\hat{\alpha}}, \Theta)$ where $\Gamma \longrightarrow \Delta_0$, and $\Delta_{\hat{\alpha}}$ is either $\hat{\alpha}$ or $\hat{\alpha} = \tau$, and Θ is soft.

Lemma 24 (Extension Order).

(i) *If $\Gamma_L, \alpha, \Gamma_R \longrightarrow \Delta$ then $\Delta = (\Delta_L, \alpha, \Delta_R)$ where $\Gamma_L \longrightarrow \Delta_L$.
Moreover, if Γ_R is soft then Δ_R is soft.*

(ii) *If $\Gamma_L, \blacktriangleright_{\hat{\alpha}}, \Gamma_R \longrightarrow \Delta$ then $\Delta = (\Delta_L, \blacktriangleright_{\hat{\alpha}}, \Delta_R)$ where $\Gamma_L \longrightarrow \Delta_L$.
Moreover, if Γ_R is soft then Δ_R is soft.*

(iii) *If $\Gamma_L, \hat{\alpha}, \Gamma_R \longrightarrow \Delta$ then $\Delta = \Delta_L, \Theta, \Delta_R$ where $\Gamma_L \longrightarrow \Delta_L$ and Θ is either $\hat{\alpha}$ or $\hat{\alpha} = \tau$ for some τ .*

(iv) *If $\Gamma_L, \hat{\alpha} = \tau, \Gamma_R \longrightarrow \Delta$ then $\Delta = \Delta_L, \hat{\alpha} = \tau', \Delta_R$ where $\Gamma_L \longrightarrow \Delta_L$ and $[\Delta_L]\tau = [\Delta_L]\tau'$.*

(v) If $\Gamma_L, x : A, \Gamma_R \longrightarrow \Delta$ then $\Delta = (\Delta_L, x : A', \Delta_R)$ where $\Gamma_L \longrightarrow \Delta_L$ and $[\Delta_L]A = [\Delta_L]A'$.
Moreover, Γ_R is soft if and only if Δ_R is soft.

Lemma 25 (Extension Weakening). If $\Gamma \vdash A$ and $\Gamma \longrightarrow \Delta$ then $\Delta \vdash A$.

Lemma 26 (Solution Admissibility for Extension). If $\Gamma_L \vdash \tau$ then $\Gamma_L, \hat{\alpha}, \Gamma_R \longrightarrow \Gamma_L, \hat{\alpha} = \tau, \Gamma_R$.

Lemma 27 (Solved Variable Addition for Extension). If $\Gamma_L \vdash \tau$ then $\Gamma_L, \Gamma_R \longrightarrow \Gamma_L, \hat{\alpha} = \tau, \Gamma_R$.

Lemma 28 (Unsolved Variable Addition for Extension). We have that $\Gamma_L, \Gamma_R \longrightarrow \Gamma_L, \hat{\alpha}, \Gamma_R$.

Lemma 29 (Parallel Admissibility).

If $\Gamma_L \longrightarrow \Delta_L$ and $\Gamma_L, \Gamma_R \longrightarrow \Delta_L, \Delta_R$ then:

(i) $\Gamma_L, \hat{\alpha}, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha}, \Delta_R$

(ii) If $\Delta_L \vdash \tau'$ then $\Gamma_L, \hat{\alpha}, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha} = \tau', \Delta_R$.

(iii) If $\Gamma_L \vdash \tau$ and $\Delta_L \vdash \tau'$ and $[\Delta_L]\tau = [\Delta_L]\tau'$, then $\Gamma_L, \hat{\alpha} = \tau, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha} = \tau', \Delta_R$.

Lemma 30 (Parallel Extension Solution).

If $\Gamma_L, \hat{\alpha}, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha} = \tau', \Delta_R$ and $\Gamma_L \vdash \tau$ and $[\Delta_L]\tau = [\Delta_L]\tau'$ then $\Gamma_L, \hat{\alpha} = \tau, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha} = \tau', \Delta_R$.

Lemma 31 (Parallel Variable Update).

If $\Gamma_L, \hat{\alpha}, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha} = \tau_0, \Delta_R$ and $\Gamma_L \vdash \tau_1$ and $\Delta_L \vdash \tau_2$ and $[\Delta_L]\tau_0 = [\Delta_L]\tau_1 = [\Delta_L]\tau_2$ then $\Gamma_L, \hat{\alpha} = \tau_1, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha} = \tau_2, \Delta_R$.

D.2 Instantiation Extends

Lemma 32 (Instantiation Extension).

If $\Gamma \vdash \hat{\alpha} : \leq \tau \dashv \Delta$ or $\Gamma \vdash \tau \leq \hat{\alpha} \dashv \Delta$ then $\Gamma \longrightarrow \Delta$.

D.3 Subtyping Extends

Lemma 33 (Subtyping Extension).

If $\Gamma \vdash A < : B \dashv \Delta$ then $\Gamma \longrightarrow \Delta$.

E Decidability of Instantiation

Lemma 34 (Left Unsolvedness Preservation).

If $\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_\Gamma \vdash \hat{\alpha} : \leq A \dashv \Delta$ or $\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_\Gamma \vdash A \leq \hat{\alpha} \dashv \Delta$, and $\hat{\beta} \in \text{unsolved}(\Gamma_0)$, then $\hat{\beta} \in \text{unsolved}(\Delta)$.

Lemma 35 (Left Free Variable Preservation). If $\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_\Gamma \vdash \hat{\alpha} : \leq A \dashv \Delta$ or $\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_\Gamma \vdash A \leq \hat{\alpha} \dashv \Delta$, and $\Gamma \vdash B$ and $\hat{\alpha} \notin \text{FV}([\Gamma]B)$ and $\hat{\beta} \in \text{unsolved}(\Gamma_0)$ and $\hat{\beta} \notin \text{FV}([\Gamma]B)$, then $\hat{\beta} \notin \text{FV}([\Delta]B)$.

Lemma 36 (Instantiation Size Preservation). If $\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_\Gamma \vdash \hat{\alpha} : \leq A \dashv \Delta$ or $\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_\Gamma \vdash A \leq \hat{\alpha} \dashv \Delta$, and $\Gamma \vdash B$ and $\hat{\alpha} \notin \text{FV}([\Gamma]B)$, then $|\text{FV}([\Gamma]B)| = |\text{FV}([\Delta]B)|$, where $|C|$ is the plain size of the term C .

This lemma lets us show decidability by taking the size of the type argument as the induction metric.

Theorem 7 (Decidability of Instantiation). If $\Gamma = \Gamma_0[\hat{\alpha}]$ and $\Gamma \vdash A$ such that $[\Gamma]A = A$ and $\hat{\alpha} \notin \text{FV}(A)$, then:

(1) Either there exists Δ such that $\Gamma_0[\hat{\alpha}] \vdash \hat{\alpha} : \leq A \dashv \Delta$, or not.

(2) Either there exists Δ such that $\Gamma_0[\hat{\alpha}] \vdash A \leq \hat{\alpha} \dashv \Delta$, or not.

F Decidability of Algorithmic Subtyping

F.1 Lemmas for Decidability of Subtyping

Lemma 37 (Monotypes Solve Variables). *If $\Gamma \vdash \hat{\alpha} : \leq \tau \dashv \Delta$ or $\Gamma \vdash \tau \leq \hat{\alpha} : \dashv \Delta$, then if $[\Gamma]\tau = \tau$ and $\hat{\alpha} \notin \text{FV}([\Gamma]\tau)$, then $|\text{unsolved}(\Gamma)| = |\text{unsolved}(\Delta)| + 1$.*

Lemma 38 (Monotype Monotonicity). *If $\Gamma \vdash \tau_1 < : \tau_2 \dashv \Delta$ then $|\text{unsolved}(\Delta)| \leq |\text{unsolved}(\Gamma)|$.*

Lemma 39 (Substitution Decreases Size). *If $\Gamma \vdash A$ then $|\Gamma \vdash [\Gamma]A| \leq |\Gamma \vdash A|$.*

Lemma 40 (Monotype Context Invariance).

If $\Gamma \vdash \tau < : \tau' \dashv \Delta$ where $[\Gamma]\tau = \tau$ and $[\Gamma]\tau' = \tau'$ and $|\text{unsolved}(\Gamma)| = |\text{unsolved}(\Delta)|$ then $\Gamma = \Delta$.

F.2 Decidability of Subtyping

Theorem 8 (Decidability of Subtyping).

Given a context Γ and types A, B such that $\Gamma \vdash A$ and $\Gamma \vdash B$ and $[\Gamma]A = A$ and $[\Gamma]B = B$, it is decidable whether there exists Δ such that $\Gamma \vdash A < : B \dashv \Delta$.

G Decidability of Typing

Theorem 9 (Decidability of Typing).

- (i) Synthesis: *Given a context Γ and a term e , it is decidable whether there exist a type A and a context Δ such that $\Gamma \vdash e \Rightarrow A \dashv \Delta$.*
- (ii) Checking: *Given a context Γ , a term e , and a type B such that $\Gamma \vdash B$, it is decidable whether there is a context Δ such that $\Gamma \vdash e \Leftarrow B \dashv \Delta$.*
- (iii) Application: *Given a context Γ , a term e , and a type A such that $\Gamma \vdash A$, it is decidable whether there exist a type C and a context Δ such that $\Gamma \vdash A \bullet e \Rightarrow C \dashv \Delta$.*

H Soundness of Subtyping

Definition 5 (Filling). *The filling of a context $|\Gamma|$ solves all unsolved variables:*

$$\begin{aligned}
 |\cdot| &= \cdot \\
 |\Gamma, x : A| &= |\Gamma|, x : A \\
 |\Gamma, \alpha| &= |\Gamma|, \alpha \\
 |\Gamma, \hat{\alpha} = \tau| &= |\Gamma|, \hat{\alpha} = \tau \\
 |\Gamma, \blacktriangleright \hat{\alpha}| &= |\Gamma|, \blacktriangleright \hat{\alpha} \\
 |\Gamma, \hat{\alpha}| &= |\Gamma|, \hat{\alpha} = 1
 \end{aligned}$$

H.1 Lemmas for Soundness

Lemma 41 (Uvar Preservation).

If $\alpha \in \Omega$ and $\Delta \longrightarrow \Omega$ then $\alpha \in [\Omega]\Delta$.

Proof. By induction on Ω , following the definition of context application. □

Lemma 42 (Variable Preservation).

If $(x : A) \in \Delta$ or $(x : A) \in \Omega$ and $\Delta \longrightarrow \Omega$ then $(x : [\Omega]A) \in [\Omega]\Delta$.

Lemma 43 (Substitution Typing). *If $\Gamma \vdash A$ then $\Gamma \vdash [\Gamma]A$.*

Lemma 44 (Substitution for Well-Formedness). *If $\Omega \vdash A$ then $[\Omega]\Omega \vdash [\Omega]A$.*

Lemma 45 (Substitution Stability).

For any well-formed complete context (Ω, Ω_Z) , if $\Omega \vdash A$ then $[\Omega]A = [\Omega, \Omega_Z]A$.

Lemma 46 (Context Partitioning).

If $\Delta, \blacktriangleright_{\hat{\alpha}}, \Theta \longrightarrow \Omega, \blacktriangleright_{\hat{\alpha}}, \Omega_Z$ then there is a Ψ such that $[\Omega, \blacktriangleright_{\hat{\alpha}}, \Omega_Z](\Delta, \blacktriangleright_{\hat{\alpha}}, \Theta) = [\Omega]\Delta, \Psi$.

Lemma 47 (Softness Goes Away).

If $\Delta, \Theta \longrightarrow \Omega, \Omega_Z$ where $\Delta \longrightarrow \Omega$ and Θ is soft, then $[\Omega, \Omega_Z](\Delta, \Theta) = [\Omega]\Delta$.

Proof. By induction on Θ , following the definition of $[\Omega]\Gamma$. □

Lemma 48 (Filling Completes). *If $\Gamma \longrightarrow \Omega$ and (Γ, Θ) is well-formed, then $\Gamma, \Theta \longrightarrow \Omega, |\Theta|$.*

Proof. By induction on Θ , following the definition of $|-|$ and applying the rules for \longrightarrow . □

Lemma 49 (Stability of Complete Contexts).

If $\Gamma \longrightarrow \Omega$ then $[\Omega]\Gamma = [\Omega]\Omega$.

Lemma 50 (Finishing Types).

If $\Omega \vdash A$ and $\Omega \longrightarrow \Omega'$ then $[\Omega]A = [\Omega']A$.

Lemma 51 (Finishing Completions).

If $\Omega \longrightarrow \Omega'$ then $[\Omega]\Omega = [\Omega']\Omega'$.

Lemma 52 (Confluence of Completeness).

If $\Delta_1 \longrightarrow \Omega$ and $\Delta_2 \longrightarrow \Omega$ then $[\Omega]\Delta_1 = [\Omega]\Delta_2$.

H.2 Instantiation Soundness

Theorem 10 (Instantiation Soundness).

Given $\Delta \longrightarrow \Omega$ and $[\Gamma]B = B$ and $\hat{\alpha} \notin \text{FV}(B)$:

(1) *If $\Gamma \vdash \hat{\alpha} : \leq B \dashv \Delta$ then $[\Omega]\Delta \vdash [\Omega]\hat{\alpha} \leq [\Omega]B$.*

(2) *If $\Gamma \vdash B \leq \hat{\alpha} \dashv \Delta$ then $[\Omega]\Delta \vdash [\Omega]B \leq [\Omega]\hat{\alpha}$.*

H.3 Soundness of Subtyping

Theorem 11 (Soundness of Algorithmic Subtyping).

If $\Gamma \vdash A <: B \dashv \Delta$ where $[\Gamma]A = A$ and $[\Gamma]B = B$ and $\Delta \longrightarrow \Omega$ then $[\Omega]\Delta \vdash [\Omega]A \leq [\Omega]B$.

Corollary 53 (Soundness, Pretty Version). *If $\Psi \vdash A <: B \dashv \Delta$, then $\Psi \vdash A \leq B$.*

I Typing Extension

Lemma 54 (Typing Extension).

If $\Gamma \vdash e \leftarrow A \dashv \Delta$ or $\Gamma \vdash e \Rightarrow A \dashv \Delta$ or $\Gamma \vdash A \bullet e \Rightarrow C \dashv \Delta$ then $\Gamma \longrightarrow \Delta$.

J Soundness of Typing

Theorem 12 (Soundness of Algorithmic Typing). *Given $\Delta \longrightarrow \Omega$:*

(i) *If $\Gamma \vdash e \leftarrow A \dashv \Delta$ then $[\Omega]\Delta \vdash e \leftarrow [\Omega]A$.*

(ii) *If $\Gamma \vdash e \Rightarrow A \dashv \Delta$ then $[\Omega]\Delta \vdash e \Rightarrow [\Omega]A$.*

(iii) *If $\Gamma \vdash A \bullet e \Rightarrow C \dashv \Delta$ then $[\Omega]\Delta \vdash [\Omega]A \bullet e \Rightarrow [\Omega]C$.*

K Completeness of Subtyping

K.1 Instantiation Completeness

Theorem 13 (Instantiation Completeness).

Given $\Gamma \longrightarrow \Omega$ and $A = [\Gamma]A$ and $\hat{\alpha} \in \text{unsolved}(\Gamma)$ and $\hat{\alpha} \notin \text{FV}(A)$:

- (1) If $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq [\Omega]A$
then there are Δ, Ω' such that $\Omega \longrightarrow \Omega'$ and $\Delta \longrightarrow \Omega'$ and $\Gamma \vdash \hat{\alpha} \leq A \dashv \Delta$.
- (2) If $[\Omega]\Gamma \vdash [\Omega]A \leq [\Omega]\hat{\alpha}$
then there are Δ, Ω' such that $\Omega \longrightarrow \Omega'$ and $\Delta \longrightarrow \Omega'$ and $\Gamma \vdash A \leq \hat{\alpha} \dashv \Delta$.

K.2 Completeness of Subtyping

Theorem 14 (Generalized Completeness of Subtyping). If $\Gamma \longrightarrow \Omega$ and $\Gamma \vdash A$ and $\Gamma \vdash B$ and $[\Omega]\Gamma \vdash [\Omega]A \leq [\Omega]B$ then there exist Δ and Ω' such that $\Delta \longrightarrow \Omega'$ and $\Omega \longrightarrow \Omega'$ and $\Gamma \vdash [\Gamma]A <: [\Gamma]B \dashv \Delta$.

Corollary 55 (Completeness of Subtyping). If $\Psi \vdash A \leq B$ then there is a Δ such that $\Psi \vdash A <: B \dashv \Delta$.

L Completeness of Typing

Theorem 15 (Completeness of Algorithmic Typing). Given $\Gamma \longrightarrow \Omega$ and $\Gamma \vdash A$:

- (i) If $[\Omega]\Gamma \vdash e \Leftarrow [\Omega]A$
then there exist Δ and Ω'
such that $\Delta \longrightarrow \Omega'$ and $\Omega \longrightarrow \Omega'$ and $\Gamma \vdash e \Leftarrow [\Gamma]A \dashv \Delta$.
- (ii) If $[\Omega]\Gamma \vdash e \Rightarrow A$
then there exist Δ, Ω' , and A'
such that $\Delta \longrightarrow \Omega'$ and $\Omega \longrightarrow \Omega'$ and $\Gamma \vdash e \Rightarrow A' \dashv \Delta$ and $A = [\Omega']A'$.
- (iii) If $[\Omega]\Gamma \vdash [\Omega]A \bullet e \Rightarrow C$
then there exist Δ, Ω' , and C'
such that $\Delta \longrightarrow \Omega'$ and $\Omega \longrightarrow \Omega'$ and $\Gamma \vdash [\Gamma]A \bullet e \Rightarrow C' \dashv \Delta$ and $C = [\Omega']C'$.

Proofs

In the rest of this document, we prove the results stated above, with the same sectioning.

A' Declarative Subtyping

Proposition 1 (Weakening). *If $\Psi \vdash A$ then $\Psi, \Psi' \vdash A$ by a derivation of the same size.*

Proposition 2 (Substitution). *If $\Psi \vdash A$ and $\Psi, \alpha, \Psi' \vdash B$ then $\Psi, \Psi' \vdash [A/\alpha]B$.*

The proofs of these two propositions are routine inductions.

A'.1 Properties of Well-Formedness

A'.2 Reflexivity

Lemma 3 (Reflexivity of Declarative Subtyping). *Subtyping is reflexive: if $\Psi \vdash A$ then $\Psi \vdash A \leq A$.*

Proof. By induction on A .

• **Case** $A = 1$: Apply rule $\leq\text{Unit}$.

• **Case** $A = \alpha$: Apply rule $\leq\text{Var}$.

• **Case** $A = A_1 \rightarrow A_2$:

$\Psi \vdash A_1 \leq A_1$ By i.h.

$\Psi \vdash A_2 \leq A_2$ By i.h.

$\Psi \vdash A_1 \rightarrow A_2 \leq A_1 \rightarrow A_2$ By $\leq\rightarrow$

• **Case** $A = \forall\alpha. A_0$:

$\Psi, \alpha \vdash A_0 \leq A_0$ By i.h.

$\Psi, \alpha \vdash \alpha$ By DeclUvarWF

$\Psi, \alpha \vdash [\alpha/\alpha]A_0 \leq A_0$ By def. of substitution

$\Psi, \alpha \vdash \forall\alpha. A_0 \leq A_0$ By $\leq\forall\text{L}$

$\Psi \vdash \forall\alpha. A_0 \leq \forall\alpha. A_0$ By $\leq\forall\text{R}$ □

A'.3 Subtyping Implies Well-Formedness

Lemma 4 (Well-Formedness). *If $\Psi \vdash A \leq B$ then $\Psi \vdash A$ and $\Psi \vdash B$.*

Proof. By induction on the given derivation. All 5 cases are straightforward. □

A'.4 Substitution

Lemma 5 (Substitution). *If $\Psi \vdash \tau$ and $\Psi, \alpha, \Psi' \vdash A \leq B$ then $\Psi, [\tau/\alpha]\Psi' \vdash [\tau/\alpha]A \leq [\tau/\alpha]B$.*

Proof. By induction on the given derivation.

• **Case** $\frac{\beta \in (\Psi, \alpha, \Psi')}{\Psi, \alpha, \Psi' \vdash \beta \leq \beta} \leq\text{Var}$

It is given that $\Psi \vdash \tau$.

Either $\beta = \alpha$ or $\beta \neq \alpha$. In the former case: We need to show $\Psi, \Psi' \vdash [\tau/\alpha]\alpha \leq [\tau/\alpha]\alpha$, that is, $\Psi, \Psi' \vdash \tau \leq \tau$, which follows by Lemma 3 (Reflexivity of Declarative Subtyping). In the latter case: We need to show $\Psi, \Psi' \vdash [\tau/\alpha]\beta \leq [\tau/\alpha]\beta$, that is, $\Psi, \Psi' \vdash \beta \leq \beta$. Since $\beta \in (\Psi, \alpha, \Psi')$ and $\beta \neq \alpha$, we have $\beta \in (\Psi, \Psi')$, so applying $\leq\text{Var}$ gives the result.

- **Case**

$$\frac{}{\Psi, \alpha, \Psi' \vdash 1 \leq 1} \leq \text{Unit}$$

For all τ , substituting $[\tau/\alpha]1 = 1$, and applying $\leq \text{Unit}$ gives the result.

- **Case**
$$\frac{\Psi, \alpha, \Psi' \vdash B_1 \leq A_1 \quad \Psi, \alpha, \Psi' \vdash A_2 \leq B_2}{\Psi, \alpha, \Psi' \vdash A_1 \rightarrow A_2 \leq B_1 \rightarrow B_2} \leq \rightarrow$$

$$\begin{array}{l} \Psi, \alpha, \Psi' \vdash B_1 \leq A_1 \\ \Psi, \Psi' \vdash [\tau/\alpha]B_1 \leq [\tau/\alpha]A_1 \end{array} \quad \begin{array}{l} \text{Subderivation} \\ \text{By i.h.} \end{array}$$

$$\begin{array}{l} \Psi, \alpha, \Psi' \vdash A_2 \leq B_2 \\ \Psi, \Psi' \vdash [\tau/\alpha]A_2 \leq [\tau/\alpha]B_2 \end{array} \quad \begin{array}{l} \text{Subderivation} \\ \text{By i.h.} \end{array}$$

$$\begin{array}{l} \Psi, \Psi' \vdash ([\tau/\alpha]A_1) \rightarrow ([\tau/\alpha]A_2) \leq ([\tau/\alpha]B_1) \rightarrow ([\tau/\alpha]B_2) \\ \Psi, \Psi' \vdash [\tau/\alpha](A_1 \rightarrow A_2) \leq [\tau/\alpha](B_1 \rightarrow B_2) \end{array} \quad \begin{array}{l} \text{By } \leq \rightarrow \\ \text{By definition of subst.} \end{array}$$

- **Case**
$$\frac{\Psi, \alpha, \Psi' \vdash \sigma \quad \Psi, \alpha, \Psi' \vdash [\sigma/\beta]A_0 \leq B}{\Psi, \alpha, \Psi' \vdash \forall \beta. A_0 \leq B} \leq \forall L$$

$$\begin{array}{l} \Psi, \alpha, \Psi' \vdash [\sigma/\beta]A_0 \leq B \\ \Psi, \Psi' \vdash [\tau/\alpha][\sigma/\beta]A_0 \leq [\tau/\alpha]B \\ \Psi, \Psi' \vdash [[\tau/\alpha]\sigma / \beta][\tau/\alpha]A_0 \leq [\tau/\alpha]B \end{array} \quad \begin{array}{l} \text{Subderivation} \\ \text{By i.h.} \\ \text{By distributivity of substitution} \end{array}$$

$$\begin{array}{l} \Psi, \alpha, \Psi' \vdash \sigma \\ \Psi \vdash \tau \\ \Psi, \Psi' \vdash [\tau/\alpha]\sigma \end{array} \quad \begin{array}{l} \text{Premise} \\ \text{Given} \\ \text{By Proposition 2} \end{array}$$

$$\begin{array}{l} \Psi, \Psi' \vdash \forall \beta. [\tau/\alpha]A_0 \leq [\tau/\alpha]B \\ \Psi, \Psi' \vdash [\tau/\alpha](\forall \beta. A_0) \leq [\tau/\alpha]B \end{array} \quad \begin{array}{l} \text{By } \leq \forall L \\ \text{By definition of substitution} \end{array}$$

- **Case**
$$\frac{\Psi, \alpha, \Psi', \beta \vdash A \leq B_0}{\Psi, \alpha, \Psi' \vdash A \leq \forall \beta. B_0} \leq \forall R$$

$$\begin{array}{l} \Psi, \alpha, \Psi', \beta \vdash A \leq B_0 \\ \Psi, \Psi', \beta \vdash [\tau/\alpha]A \leq [\tau/\alpha]B_0 \\ \Psi, \Psi' \vdash [\tau/\alpha]A \leq \forall \beta. [\tau/\alpha]B_0 \end{array} \quad \begin{array}{l} \text{Subderivation} \\ \text{By i.h.} \\ \text{By } \leq \forall R \end{array}$$

$$\Psi, \Psi' \vdash [\tau/\alpha]A \leq [\tau/\alpha](\forall \beta. B_0) \quad \text{By definition of substitution} \quad \square$$

A'.5 Transitivity

To prove transitivity, we use a metric that adapts ideas from a proof of cut elimination by Pfenning (1995).

Lemma 6 (Transitivity of Declarative Subtyping). *If $\Psi \vdash A \leq B$ and $\Psi \vdash B \leq C$ then $\Psi \vdash A \leq C$.*

Proof. By induction with the following metric:

$$\langle \# \forall(B), \mathcal{D}_1 + \mathcal{D}_2 \rangle$$

where $\langle \dots \rangle$ denotes lexicographic order, the first part $\# \forall(B)$ is the number of quantifiers in B , and the second part is the (simultaneous) size of the derivations $\mathcal{D}_1 :: \Psi \vdash A \leq B$ and $\mathcal{D}_2 :: \Psi \vdash B \leq C$. We need to consider the number of quantifiers first in one case: when $\leq \forall R$ concluded \mathcal{D}_1 and $\leq \forall L$ concluded \mathcal{D}_2 , because in that case, the derivations on which the i.h. must be applied are not necessarily smaller.

- **Case**
$$\frac{\alpha \in \Psi}{\Psi \vdash \alpha \leq \alpha} \leq \text{Var} \quad \frac{\alpha \in \Psi}{\Psi \vdash \alpha \leq \alpha} \leq \text{Var}$$

Apply rule $\leq \text{Var}$.

- **Case $\leq\text{Unit} / \leq\text{Unit}$:** Similar to the $\leq\text{Var} / \leq\text{Var}$ case.

$$\bullet \text{ Case } \frac{\Psi \vdash B_1 \leq A_1 \quad \Psi \vdash A_2 \leq B_2}{\Psi \vdash A_1 \rightarrow A_2 \leq B_1 \rightarrow B_2} \leq\rightarrow \quad \frac{\Psi \vdash C_1 \leq B_1 \quad \Psi \vdash B_2 \leq C_2}{\Psi \vdash B_1 \rightarrow B_2 \leq C_1 \rightarrow C_2} \leq\rightarrow$$

By i.h. on the 3rd and 1st subderivations, $\Psi \vdash C_1 \leq A_1$.

By i.h. on the 2nd and 4th subderivations, $\Psi \vdash A_2 \leq C_2$.

By $\leq\rightarrow$, $\Psi \vdash A_1 \rightarrow A_2 \leq C_1 \rightarrow C_2$.

If $\leq\forall\text{L}$ concluded \mathcal{D}_1 :

$$\bullet \text{ Case } \frac{\Psi \vdash \tau \quad \Psi \vdash [\tau/\alpha]A_0 \leq B}{\Psi \vdash \forall\alpha. A_0 \leq B} \leq\forall\text{L}$$

$\Psi \vdash \tau$ Premise
 $\Psi \vdash [\tau/\alpha]A_0 \leq B$ Subderivation
 $\Psi \vdash B \leq C$ Given (\mathcal{D}_2)
 $\Psi \vdash [\tau/\alpha]A_0 \leq C$ By i.h.
 $\Psi \vdash \forall\alpha. A_0 \leq C$ By $\leq\forall\text{L}$

If $\leq\forall\text{R}$ concluded \mathcal{D}_2 :

$$\bullet \text{ Case } \frac{\Psi, \beta \vdash B \leq C}{\Psi \vdash B \leq \forall\beta. C} \leq\forall\text{R}$$

$\Psi \vdash \tau$ Premise
 $\Psi, \beta \vdash B \leq C$ Subderivation
 $\Psi \vdash A \leq B$ Given (\mathcal{D}_1)
 $\Psi, \beta \vdash A \leq B$ By Proposition 1
 $\Psi, \beta \vdash A \leq C$ By i.h.
 $\Psi \vdash A \leq \forall\beta. C$ By $\leq\forall\text{L}$

The only remaining possible case is $\leq\forall\text{R} / \leq\forall\text{L}$.

$$\bullet \text{ Case } \frac{\Psi, \beta \vdash A \leq B_0}{\Psi \vdash A \leq \forall\beta. B_0} \leq\forall\text{R} \quad \frac{\Psi \vdash \tau \quad \Psi \vdash [\tau/\beta]B_0 \leq C}{\Psi \vdash \forall\beta. B_0 \leq C} \leq\forall\text{L}$$

$\Psi, \beta \vdash A \leq B_0$ Subderivation of \mathcal{D}_1
 $\Psi \vdash \tau$ Premise of \mathcal{D}_2
 $\Psi \vdash [\tau/\beta]A \leq [\tau/\beta]B_0$ By Lemma 5 (Substitution)
 $[\tau/\beta]A = A$ β cannot appear in A
 $\Psi \vdash A \leq [\tau/\beta]B_0$ By above equality
 $\Psi \vdash [\tau/\beta]B_0 \leq C$ Subderivation of \mathcal{D}_2
 $\Psi \vdash A \leq C$ By i.h. (one less \forall quantifier in B) □

A'.6 Invertibility of $\leq\forall\text{R}$

Lemma 7 (Invertibility).

If \mathcal{D} derives $\Psi \vdash A \leq \forall\beta. B$ then \mathcal{D}' derives $\Psi, \beta \vdash A \leq B$ where $\mathcal{D}' < \mathcal{D}$.

Proof. By induction on the given derivation \mathcal{D} .

- **Cases $\leq\text{Var}$, $\leq\text{Unit}$, $\leq\rightarrow$:** Impossible: the supertype cannot have the form $\forall\beta. B$.

- **Case** $\frac{\Psi, \beta \vdash A \leq B}{\Psi \vdash A \leq \forall\beta. B} \leq\forall R$

The subderivation is exactly what we need, and is strictly smaller than \mathcal{D} .

- **Case** $\frac{\Psi \vdash \tau \quad \Psi \vdash [\tau/\alpha]A_0 \leq \forall\beta. B}{\Psi \vdash \forall\alpha. A_0 \leq \forall\beta. B} \leq\forall L$

By i.h., \mathcal{D}'_0 derives $\Psi, \beta \vdash [\tau/\alpha]A_0 \leq B$ where $\mathcal{D}'_0 < \mathcal{D}_0$.

By $\leq\forall L$, \mathcal{D}' derives $\Psi, \beta \vdash \forall\alpha. A_0 \leq B$; since $\mathcal{D}'_0 < \mathcal{D}_0$, we have $\mathcal{D}' < \mathcal{D}$. □

A'.7 Non-Circularity and Equality

Lemma 8 (Occurrence).

(i) If $\Psi \vdash A \leq \tau$ then $\tau \not\prec A$.

(ii) If $\Psi \vdash \tau \leq B$ then $\tau \not\prec B$.

Proof. By induction on the given derivation.

- **Cases** $\leq\text{Var}$, $\leq\text{Unit}$: (i), (ii): Here A and B have no subterms at all, so the result is immediate.

- **Case** $\frac{\Psi \vdash B_1 \leq A_1 \quad \Psi \vdash A_2 \leq B_2}{\Psi \vdash A_1 \rightarrow A_2 \leq B_1 \rightarrow B_2} \leq\rightarrow$

(i) Here, $A = A_1 \rightarrow A_2$ and $\tau = B_1 \rightarrow B_2$.

$B_1 \not\prec A_1$ By i.h. (ii)

$B_1 \rightarrow B_2 \not\prec A_1$ Suppose $B_1 \rightarrow B_2 \preceq A_1$. Then $B_1 \succ A_1$: contradiction.

$B_2 \not\prec A_2$ By i.h. (i)

$B_1 \rightarrow B_2 \not\prec A_2$ Similar

Suppose (for a contradiction) that $B_1 \rightarrow B_2 \preceq A_1 \rightarrow A_2$.

Now $B_1 \rightarrow B_2 \preceq A_1$ or $B_1 \rightarrow B_2 \preceq A_2$.

But above, we showed that both were false: contradiction.

Therefore, $B_1 \rightarrow B_2 \not\prec A_1 \rightarrow A_2$.

Therefore, $B_1 \rightarrow B_2 \not\prec A_1 \rightarrow A_2$.

(ii) Here, $A = \tau$ and $B = B_1 \rightarrow B_2$.

Symmetric to the previous case.

- **Case** $\frac{\Psi \vdash \tau' \quad \Psi \vdash [\tau'/\alpha]A_0 \leq \tau}{\Psi \vdash \forall\alpha. A_0 \leq \tau} \leq\forall L$

In part (ii), this case cannot arise, so we prove part (i).

By i.h. (i), $\tau \not\prec [\tau'/\alpha]A_0$.

It follows from the definition of \preceq that $\tau \not\prec \forall\alpha. A_0$.

- **Case** $\frac{\Psi, \beta \vdash \tau \leq B_0}{\Psi \vdash \tau \leq \forall\beta. B_0} \leq\forall R$

In part (i), this case cannot arise, so we prove part (ii).

Similar to the $\leq\forall L$ case. □

Lemma 9 (Monotype Equality). If $\Psi \vdash \sigma \leq \tau$ then $\sigma = \tau$.

Proof. By induction on the given derivation.

- **Case $\leq\text{Var}$:** Immediate.
- **Case $\leq\text{Unit}$:** Immediate.
- **Case $\leq\rightarrow$:**
$$\frac{\Psi \vdash B_1 \leq A_1 \quad \Psi \vdash A_2 \leq B_2}{\Psi \vdash A_1 \rightarrow A_2 \leq B_1 \rightarrow B_2} \leq\rightarrow$$
By i.h. on each subderivation, $B_1 = A_1$ and $A_2 = B_2$. Therefore $A_1 \rightarrow A_2 = B_1 \rightarrow B_2$.
- **Case $\leq\forall\text{L}$:** Here $\sigma = \forall\alpha. A_0$, which is not a monotype, so this case is impossible.
- **Case $\leq\forall\text{R}$:** Here $\tau = \forall\beta. B_0$, which is not a monotype, so this case is impossible. □

B' Type Assignment

Lemma 10 (Well-Formedness).

If $\Psi \vdash e \Leftarrow A$ or $\Psi \vdash e \Rightarrow A$ or $\Psi \vdash A \bullet e \Rightarrow C$ then $\Psi \vdash A$ (and in the last case, $\Psi \vdash C$).

Proof. By induction on the given derivation.

In all cases, we apply the induction hypothesis to all subderivations.

- In the DeclVar and $\text{Decl}\rightarrow\text{I}$ cases, we use our standard assumption that every context appearing in a derivation is well-formed.
- In the $\text{Decl}\rightarrow\text{I}\Rightarrow$ case, we use inversion on the $\Psi \vdash \sigma \rightarrow \tau$ premise.
- In the $\text{Decl}\forall\text{App}$ case, we use the property that if $\Psi \vdash [\tau/\alpha]A_0$ then $\Psi \vdash \forall\alpha. A_0$.
- In the DeclAnno case, we use its premise. □

Theorem 1 (Completeness of Bidirectional Typing).

If $\Psi \vdash e : A$ then there exists e' such that $\Psi \vdash e' \Rightarrow A$ and $|e'| = e$.

Proof. By induction on the derivation of $\Psi \vdash e : A$.

- **Case**
$$\frac{x : A \in \Psi}{\Psi \vdash x : A} \text{AVar}$$

Immediate, by rule DeclVar .

- **Case**
$$\frac{\Psi, x : A \vdash e : B}{\Psi \vdash \lambda x. e : A \rightarrow B} \text{A}\rightarrow\text{I}$$

By inversion, we have $\Psi, x : A \vdash e : B$.

By induction, we have $\Psi, x : A \vdash e' \Rightarrow B$, where $|e'| = e$.

By Lemma 3 (Reflexivity of Declarative Subtyping), $\Psi \vdash B \leq B$.

By rule DeclSub , $\Psi, x : A \vdash e' \Leftarrow B$.

By rule $\text{Decl}\rightarrow\text{I}$, $\Psi \vdash \lambda x. e' \Leftarrow A \rightarrow B$.

By Lemma 10 (Well-Formedness), $\Psi \vdash A \rightarrow B$.

By rule DeclAnno , $\Psi \vdash ((\lambda x. e') : A \rightarrow B) \Rightarrow A \rightarrow B$.

By definition, $|((\lambda x. e') : A \rightarrow B)| = |\lambda x. e'| = \lambda x. |e'| = \lambda x. e$.

- **Case**
$$\frac{\Psi \vdash e_1 : A \rightarrow B \quad \Psi \vdash e_2 : A}{\Psi \vdash e_1 e_2 : B} A \rightarrow E$$

By induction, $\Psi \vdash e'_1 \Rightarrow A \rightarrow B$ and $|e'_1| = e_1$.

By induction, $\Psi \vdash e'_2 \Rightarrow A$ and $|e'_2| = e_2$.

By Lemma 3 (Reflexivity of Declarative Subtyping), $\Psi \vdash A \leq A$.

By rule DeclSub, $\Psi \vdash e'_2 \Leftarrow A$.

By rule Decl \rightarrow App, $\Psi \vdash A \rightarrow B \bullet e'_2 \Rightarrow B$.

By rule Decl \rightarrow E, $\Psi \vdash e'_1 e'_2 \Rightarrow B$.

By definition, $|e'_1 e'_2| = |e'_1| |e'_2| = e_1 e_2$.

- **Case**
$$\frac{\Psi, \alpha \vdash e : A}{\Psi \vdash e : \forall \alpha. A} A \forall I$$

By induction, $\Psi, \alpha \vdash e' \Rightarrow A$ where $|e'| = e$.

By Lemma 3 (Reflexivity of Declarative Subtyping), $\Psi, \alpha \vdash A \leq A$.

By rule DeclSub, $\Psi, \alpha \vdash e' \Leftarrow A$.

By rule Decl \forall I, $\Psi \vdash e' \Leftarrow \forall \alpha. A$.

By Lemma 10 (Well-Formedness), $\Psi \vdash \forall \alpha. A$.

By rule DeclAnno, $\Psi \vdash (e' : \forall \alpha. A) \Rightarrow \forall \alpha. A$.

By definition, $|e' : \forall \alpha. A| = |e'| = e$.

- **Case**
$$\frac{\Psi \vdash e : \forall \alpha. A \quad \Psi \vdash \tau}{\Psi \vdash e : [\tau/\alpha]A} A \forall E$$

By induction, $\Psi \vdash e' \Rightarrow \forall \alpha. A$ where $|e'| = e$.

By Lemma 3 (Reflexivity of Declarative Subtyping), $\Psi \vdash [\tau/\alpha]A \leq [\tau/\alpha]A$.

By $\leq \forall L$, $\Psi \vdash \forall \alpha. A \leq [\tau/\alpha]A$.

By rule DeclSub, $\Psi \vdash e' \Leftarrow [\tau/\alpha]A$.

By Lemma 10 (Well-Formedness), $\Psi \vdash [\tau/\alpha]A$.

By rule DeclAnno, $\Psi \vdash (e' : [\tau/\alpha]A) \Leftarrow [\tau/\alpha]A$.

By definition, $|e' : [\tau/\alpha]A| = |e'| = e$. □

Lemma 11 (Subtyping Coercion). *If $\Psi \vdash A \leq B$ then there exists f which is $\beta\eta$ -equal to the identity such that $\Psi \vdash f : A \rightarrow B$.*

Proof. By induction on the derivation of $\Psi \vdash A \leq B$.

- **Case**
$$\frac{\alpha \in \Psi}{\Psi \vdash \alpha \leq \alpha} \leq \text{Var}$$

Choose $f = \lambda x. x$.

Clearly $\Psi \vdash \lambda x. x : \alpha \rightarrow \alpha$.

- **Case**
$$\frac{}{\Psi \vdash 1 \leq 1} \leq \text{Unit}$$

Choose $f = \lambda x. x$.

Clearly $\Psi \vdash \lambda x. x : 1 \rightarrow 1$.

- **Case**
$$\frac{\Psi \vdash B_1 \leq A_1 \quad \Psi \vdash A_2 \leq B_2}{\Psi \vdash A_1 \rightarrow A_2 \leq B_1 \rightarrow B_2} \leq \rightarrow$$

By induction, we have $g : B_1 \rightarrow A_1$, which is $\beta\eta$ -equal to the identity.

By induction, we have $k : A_2 \rightarrow B_2$, which is $\beta\eta$ -equal to the identity.

Let f be $\lambda h. k \circ h \circ g$.

It is easy to verify that $\Psi \vdash f : (A_1 \rightarrow A_2) \rightarrow (B_1 \rightarrow B_2)$.

Since k and g are identities, $f =_{\beta\eta} \lambda h. h$.

$$\bullet \text{ Case } \frac{\Psi \vdash \tau \quad \Psi \vdash [\tau/\alpha]A \leq B}{\Psi \vdash \forall \alpha. A \leq B} \leq \forall L$$

By induction, $g : [\tau/\alpha]A \rightarrow B$.

Let $f \triangleq \lambda x. g x$.

f is an eta-expansion of g , which is $\beta\eta$ -equal to the identity. Hence f is too.

Also, $\lambda x. g x : (\forall \alpha. A) \rightarrow B$, using the Decl \forall E rule on x .

$$\bullet \text{ Case } \frac{\Psi, \beta \vdash A \leq B}{\Psi \vdash A \leq \forall \beta. B} \leq \forall R$$

By induction, we have g such that $\Psi, \beta \vdash g : A \rightarrow B$.

Let $f \triangleq \lambda x. g x$.

Use the following derivation:

$$\begin{array}{c} \vdots \\ \text{WEAKEN } \frac{}{\Psi, \beta \vdash g : A \rightarrow B} \\ \frac{\Psi, x : A, \beta \vdash g : A \rightarrow B \quad \Psi, x : A, \beta \vdash x : A}{\Psi, x : A, \beta \vdash g x : B} \\ \frac{\Psi, x : A \vdash g x : \forall \beta. B}{\Psi \vdash \lambda x. g x : A \rightarrow \forall \beta. B} \quad \square \end{array}$$

Lemma 12 (Application Subtyping). *If $\Psi \vdash A \bullet e \Rightarrow C$ then there exists B such that $\Psi \vdash A \leq B \rightarrow C$ and $\Psi \vdash e \Leftarrow B$ by a smaller derivation.*

Proof. By induction on the given derivation \mathcal{D} .

$$\bullet \text{ Case } \frac{\Psi \vdash e \Leftarrow B}{\Psi \vdash B \rightarrow C \bullet e \Rightarrow C} \text{Decl} \rightarrow \text{App}$$

$\Leftarrow \mathcal{D}' :: \Psi \vdash e \Leftarrow B$ Subderivation
 $\Leftarrow \mathcal{D}' < \mathcal{D}$ \mathcal{D}' is a subderivation of \mathcal{D}
 $\Leftarrow \Psi \vdash \underbrace{B \rightarrow C}_A \leq B \rightarrow C$ By Lemma 3 (Reflexivity of Declarative Subtyping)

$$\bullet \text{ Case } \frac{\Psi \vdash \tau \quad \Psi \vdash [\tau/\alpha]A_0 \bullet e \Rightarrow C}{\Psi \vdash \forall \alpha. A_0 \bullet e \Rightarrow C} \text{Decl} \forall \text{App}$$

$\Psi \vdash \tau$ Subderivation
 $\Psi \vdash [\tau/\alpha]A_0 \bullet e \Rightarrow C$ Subderivation
 $\Psi \vdash [\tau/\alpha]A_0 \leq B \rightarrow C$ By i.h.
 $\Leftarrow \mathcal{D}' :: \Psi \vdash e \Leftarrow B$ "
 $\Leftarrow \mathcal{D}' < \mathcal{D}$ "
 $\Leftarrow \Psi \vdash \forall \alpha. A_0 \leq B \rightarrow C$ By $\leq \forall L$ □

Theorem 2 (Soundness of Bidirectional Typing). *We have that:*

- If $\Psi \vdash e \Leftarrow A$, then there is an e' such that $\Psi \vdash e' : A$ and $e' =_{\beta\eta} |e|$.
- If $\Psi \vdash e \Rightarrow A$, then there is an e' such that $\Psi \vdash e' : A$ and $e' =_{\beta\eta} |e|$.

Proof. \bullet **Case** $\frac{(x : A) \in \Psi}{\Psi \vdash x \Rightarrow A} \text{Decl} \text{Var}$

By rule AVar, $\Psi \vdash x : A$.

Note $x =_{\beta\eta} x$.

$$\bullet \text{ Case } \frac{\Psi \vdash e \Rightarrow A \quad \Psi \vdash A \leq B}{\Psi \vdash e \Leftarrow B} \text{DeclSub}$$

By induction, $\Psi \vdash e' : A$ and $e' =_{\beta\eta} |e|$.

By Lemma 11 (Subtyping Coercion), $f : A \rightarrow B$ such that $f =_{\beta\eta} \text{id}$.

By $A \rightarrow E$, $\Psi \vdash f e' : B$.

Note $f e' =_{\beta\eta} \text{id } e' =_{\beta\eta} e' =_{\beta\eta} |e|$.

$$\bullet \text{ Case } \frac{\Psi \vdash A \quad \Psi \vdash e \Leftarrow A}{\Psi \vdash (e : A) \Rightarrow A} \text{DeclAnno}$$

By induction, $\Psi \vdash e' : A$ such that $e' =_{\beta\eta} |e|$.

Note $e' =_{\beta\eta} |e| = |e : A|$.

$$\bullet \text{ Case } \frac{}{\Psi \vdash () \Leftarrow 1} \text{Decl1I}$$

By AUnit, $\Psi \vdash () : 1$.

Note $() =_{\beta\eta} ()$.

$$\bullet \text{ Case } \frac{}{\Psi \vdash () \Rightarrow 1} \text{Decl1I}\Rightarrow$$

By AUnit, $\Psi \vdash () : 1$.

Note $() =_{\beta\eta} ()$.

$$\bullet \text{ Case } \frac{\Psi, \alpha \vdash e \Leftarrow A}{\Psi \vdash e \Leftarrow \forall \alpha. A} \text{Decl}\forall I$$

By induction, $\Psi, \alpha \vdash e' : A$ such that $e' =_{\beta\eta} |e|$.

By rule $\forall I$, $\Psi \vdash e' : \forall \alpha. A$.

$$\bullet \text{ Case } \frac{\Psi, x : A \vdash e \Leftarrow B}{\Psi \vdash \lambda x. e \Leftarrow A \rightarrow B} \text{Decl}\rightarrow I$$

By induction, $\Psi, x : A \vdash e' : B$ such that $e' =_{\beta\eta} |e|$.

By $A \rightarrow I$, $\Psi \vdash \lambda x. e' : A \rightarrow B$.

Note $\lambda x. e' =_{\beta\eta} \lambda x. |e| = |\lambda x. e|$.

$$\bullet \text{ Case } \frac{\Psi \vdash \sigma \rightarrow \tau \quad \Psi, x : \sigma \vdash e \Leftarrow \tau}{\Psi \vdash \lambda x. e \Rightarrow \sigma \rightarrow \tau} \text{Decl}\rightarrow I \Rightarrow$$

By induction, $\Psi, x : \sigma \vdash e' : \tau$ such that $e' =_{\beta\eta} |e|$.

By $A \rightarrow I$, $\Psi \vdash \lambda x. e' : \sigma \rightarrow \tau$.

Note $\lambda x. e' =_{\beta\eta} \lambda x. |e| = |\lambda x. e|$.

$$\bullet \text{ Case } \frac{\Psi \vdash e_1 \Rightarrow A \quad \Psi \vdash A \bullet e_2 \Rightarrow C}{\Psi \vdash e_1 e_2 \Rightarrow C} \text{Decl}\rightarrow E$$

By induction, $\Psi \vdash e'_1 : A$ such that $e'_1 =_{\beta\eta} |e_1|$.

By Lemma 12 (Application Subtyping), there is a B such that

1. $\Psi \vdash A \leq B \rightarrow C$, and

2. $\Psi \vdash e_2 \Leftarrow B$, which is no bigger than $\Psi \vdash A \bullet e_2 \Rightarrow C$.

By Lemma 11 (Subtyping Coercion), we have f such that $\Psi \vdash f : A \rightarrow B \rightarrow C$ and $f =_{\beta\eta} \text{id}$.

By induction, we get $\Psi \vdash e'_2 : B$ and $e'_2 =_{\beta\eta} |e_2|$.

By $A \rightarrow E$ twice, $\Psi \vdash f e'_1 e'_2 : C$.

Note $f e'_1 e'_2 =_{\beta\eta} \text{id } e'_1 e'_2 =_{\beta\eta} e'_1 e'_2 =_{\beta\eta} |e_1| |e_2| = |e_1 e_2|$. □

C' Robustness of Typing

Lemma 13 (Type Substitution).

Assume $\Psi \vdash \tau$.

- If $\Psi, \alpha, \Psi' \vdash e' \Leftarrow C$ then $\Psi, [\tau/\alpha]\Psi' \vdash [\tau/\alpha]e' \Leftarrow [\tau/\alpha]C$.
- If $\Psi, \alpha, \Psi' \vdash e' \Rightarrow C$ then $\Psi, [\tau/\alpha]\Psi' \vdash [\tau/\alpha]e' \Rightarrow [\tau/\alpha]C$.
- If $\Psi, \alpha, \Psi' \vdash B \bullet e' \Rightarrow C$ then $\Psi, [\tau/\alpha]\Psi' \vdash [\tau/\alpha]B \bullet [\tau/\alpha]e' \Rightarrow [A/\alpha]C$.

Moreover, the resulting derivation contains no more applications of typing rules than the given one. (Internal subtyping derivations, however, may grow.)

Proof. By induction on the given derivation.

In the DeclVar case, split on whether the variable being typed is in Ψ or Ψ' ; the former is immediate, and in the latter, use the fact that $(x : C) \in \Psi'$ implies $(x : [\tau/\alpha]C) \in [\tau/\alpha]\Psi'$.

In the DeclSub case, use the i.h. and Lemma 5 (Substitution).

In the DeclAnno case, we are substituting in the annotation in the term, as well as in the type; we also need Proposition 2.

In Decl \rightarrow l, Decl \rightarrow r and Decl \forall l, we add to the context in the premise, which is why the statement is generalized for nonempty Ψ' . \square

Lemma 14 (Subsumption). Suppose $\Psi' \leq \Psi$. Then:

- (i) If $\Psi \vdash e \Leftarrow A$ and $\Psi \vdash A \leq A'$ then $\Psi' \vdash e \Leftarrow A'$.
- (ii) If $\Psi \vdash e \Rightarrow A$ then there exists A' such that $\Psi \vdash A' \leq A$ and $\Psi' \vdash e \Rightarrow A'$.
- (iii) If $\Psi \vdash C \bullet e \Rightarrow A$ and $\Psi \vdash C' \leq C$
then there exists A' such that $\Psi \vdash A' \leq A$ and $\Psi' \vdash C' \bullet e \Rightarrow A'$.

Proof. By mutual induction: in (i), by lexicographic induction on the derivation of the checking judgment, then of the subtyping judgment; in (ii), by induction on the derivation of the synthesis judgment; in (iii), by lexicographic induction on the derivation of the application judgment, then of the subtyping judgment.

For part (i), checking:

- **Case** $\frac{\Psi \vdash e \Rightarrow B \quad \Psi \vdash B \leq A}{\Psi \vdash e \Leftarrow A} \text{DeclSub}$
 - $\Psi \vdash e \Rightarrow B$ Subderivation
 - $\Psi' \vdash e \Rightarrow B'$ By i.h.
 - $\Psi \vdash B' \leq B$ "
 - $\Psi \vdash B \leq A$ Subderivation
 - $\Psi \vdash A \leq A'$ Given
 - $\Psi \vdash B' \leq A'$ By Lemma 6 (Transitivity of Declarative Subtyping) (twice)
 - $\Psi' \vdash B' \leq A'$ By weakening
 - $\Psi' \vdash e \Leftarrow A'$ By DeclSub
- **Case** $\frac{}{\Psi \vdash () \Leftarrow 1} \text{Decl1l}$
 - $\Psi' \vdash () \Rightarrow 1$ By Decl1l \Rightarrow
 - $\Psi \vdash 1 \leq A'$ Given
 - $\Psi' \vdash 1 \leq A'$ By weakening
 - $\Psi' \vdash () \Leftarrow A'$ By DeclSub

- **Case** $\frac{\Psi, \alpha \vdash e \Leftarrow A_0}{\Psi \vdash e \Leftarrow \forall \alpha. A_0} \text{Decl}\forall$

We consider cases of $\Psi \vdash \forall \alpha. A_0 \leq A'$:

- **Case** $\frac{\Psi, \beta \vdash \forall \alpha. A_0 \leq B}{\Psi \vdash \forall \alpha. A_0 \leq \forall \beta. B} \leq \forall R$
 - $\Psi, \beta \vdash \forall \alpha. A_0 \leq B$ Subderivation
 - $\Psi \vdash e \Leftarrow \forall \alpha. A_0$ Given
 - $\Psi' \vdash e \Leftarrow B$ By i.h. (i)
 - ☞ $\Psi' \vdash e \Leftarrow \underbrace{\forall \beta. B}_{A'}$ By Decl \forall
- **Case** $\frac{\Psi \vdash \tau \quad \Psi \vdash [\tau/\alpha]A_0 \leq A'}{\Psi \vdash \forall \alpha. A_0 \leq A'} \leq \forall L$
 - $\Psi, \alpha \vdash e \Leftarrow A_0$ Subderivation
 - $\Psi \vdash e \Leftarrow [\tau/\alpha]A_0$ By Lemma 13 (Type Substitution)
 - $\Psi \vdash [\tau/\alpha]A_0 \leq A'$ Subderivation
 - ☞ $\Psi' \vdash e \Leftarrow A'$ By i.h. (i)

- **Case** $\frac{\Psi, x : A_1 \vdash e_0 \Leftarrow A_2}{\Psi \vdash \lambda x. e_0 \Leftarrow A_1 \rightarrow A_2} \text{Decl}\rightarrow$

We consider cases of $\Psi \vdash A_1 \rightarrow A_2 \leq A'$:

- **Case** $\frac{\Psi \vdash B_1 \leq A_1 \quad \Psi \vdash A_2 \leq B_2}{\Psi \vdash A_1 \rightarrow A_2 \leq B_1 \rightarrow B_2} \leq \rightarrow$
 - $\Psi \leq \Psi'$ Given
 - $\Psi \vdash B_1 \leq A_1$ Subderivation
 - $\Psi', x : B_1 \leq \Psi, x : A_1$ By CtxSubVar
 - $\Psi', x : B_1 \vdash e_0 \Leftarrow B_2$ By i.h. (i)
 - ☞ $\Psi' \vdash \lambda x. e_0 \Leftarrow B_1 \rightarrow B_2$ By Decl \rightarrow
- **Case** $\frac{\Psi, \beta \vdash A_1 \rightarrow A_2 \leq B'}{\Psi \vdash A_1 \rightarrow A_2 \leq \forall \beta. B'} \leq \forall R$
 - $\Psi, \beta \vdash A_1 \rightarrow A_2 \leq B'$ Subderivation
 - $\Psi, \beta \vdash \lambda x. e_0 \Leftarrow A_1 \rightarrow A_2$ By weakening
 - $\Psi', \beta \vdash \lambda x. e_0 \Leftarrow B'$ By i.h. (i)
 - ☞ $\Psi' \vdash \lambda x. e_0 \Leftarrow \forall \beta. B'$ By Decl \forall

For part (ii), synthesis:

- **Case** $\frac{(x : A) \in \Psi}{\Psi \vdash x \Rightarrow A} \text{DeclVar}$

By inversion on $\Psi' \leq \Psi$, we have $(x : A') \in \Psi'$ where $\Psi \vdash A' \leq A$.
By DeclVar, $\Psi' \vdash x \Rightarrow A'$.

- **Case** $\frac{\Psi \vdash A \quad \Psi \vdash e_0 \Leftarrow A}{\Psi \vdash (e_0 : A) \Rightarrow A} \text{DeclAnno}$

Let $A' = A$.	
$\Psi \vdash A$	Subderivation
$\Psi' \vdash A$	By weakening
$\Psi \vdash e_0 \Leftarrow A$	Subderivation
$\Psi' \vdash e_0 \Leftarrow A$	By i.h.
$\Psi' \vdash (e_0 : A) \Rightarrow A'$	By DeclAnno and $A' = A$
$\Psi \vdash A' \leq A$	By Lemma 3 (Reflexivity of Declarative Subtyping)

• Case

$\Psi \vdash () \Rightarrow 1$	Decl1 \Rightarrow
Let $A' = 1$.	
$\Psi' \vdash () \Rightarrow 1$	By Decl1 \Rightarrow
$\Psi \vdash 1 \leq 1$	By \leq Unit

• Case

$\Psi \vdash \sigma \rightarrow \tau$	$\Psi, x : \sigma \vdash e_0 \Leftarrow \tau$	
$\Psi \vdash \lambda x. e_0 \Rightarrow \sigma \rightarrow \tau$		Decl \rightarrow I \Rightarrow
Let $A' = \sigma \rightarrow \tau$.		
$\Psi' \leq \Psi$		Given
$\Psi \vdash \sigma \leq \sigma$		By Lemma 3 (Reflexivity of Declarative Subtyping)
$\Psi', x : \sigma \leq \Psi, x : \sigma$		By CtxSubVar
$\Psi, x : \sigma \vdash e_0 \Leftarrow \tau$		Subderivation
$\Psi \vdash \tau \leq \tau$		By Lemma 3 (Reflexivity of Declarative Subtyping)
$\Psi', x : \sigma \vdash e_0 \Leftarrow \tau$		By i.h. (i) with τ
$\Psi \vdash A' \leq \sigma \rightarrow \tau$		By Lemma 3 (Reflexivity of Declarative Subtyping)
$\Psi' \vdash \lambda x. e_0 \Rightarrow A'$		By Decl \rightarrow I \Rightarrow

• Case

$\Psi \vdash e_1 \Rightarrow C$	$\Psi \vdash C \bullet e_2 \Rightarrow A$	
$\Psi \vdash e_1 e_2 \Rightarrow A$		Decl \rightarrow E
$\Psi \vdash e_1 \Rightarrow C$		Subderivation
$\Psi' \vdash e_1 \Rightarrow C'$		By i.h. (ii)
$\Psi \vdash C' \leq C$		"
$\Psi \vdash C \bullet e_2 \Rightarrow A$		Subderivation
$\Psi \vdash A' \leq A$		By i.h. (iii)
$\Psi' \vdash C' \bullet e_2 \Rightarrow A'$		"
$\Psi' \vdash e_1 e_2 \Rightarrow A'$		By Decl \rightarrow E

For part (iii), application:

• Case

$\Psi \vdash \tau$	$\Psi \vdash [\tau/\alpha]C_0 \bullet e \Rightarrow A$	
$\Psi \vdash \forall \alpha. C_0 \bullet e \Rightarrow A$		Decl \forall App
$\Psi \vdash C' \leq \forall \alpha. C_0$		Given
$\Psi, \alpha \vdash C' \leq C_0$		By Lemma 7 (Invertibility)
$\Psi \vdash [\tau/\alpha]C' \leq [\tau/\alpha]C_0$		By Lemma 5 (Substitution)
$\Psi \vdash C' \leq [\tau/\alpha]C_0$		α cannot appear in C'
$\Psi \vdash [\tau/\alpha]C_0 \bullet e \Rightarrow A$		Subderivation
$\Psi' \vdash C' \bullet e \Rightarrow A'$		By i.h. (iii)
$\Psi' \vdash A' \leq A$		"

- **Case**
$$\frac{\Psi \vdash e \Leftarrow C_0}{\Psi \vdash C_0 \rightarrow A \bullet e \Rightarrow A} \text{Decl} \rightarrow \text{App}$$

$$\Psi \vdash C' \leq C_0 \rightarrow A \quad \text{Given}$$
- **Case**
$$\frac{\Psi \vdash C_0 \leq C'_1 \quad \Psi \vdash C'_2 \leq A}{\Psi \vdash C'_1 \rightarrow C'_2 \leq C_0 \rightarrow A} \leq \rightarrow$$

Let $A' = C'_2$.

	$\Psi \vdash e \Leftarrow C_0$	Subderivation
	$\Psi \vdash C_0 \leq C'_1$	Subderivation
	$\Psi' \vdash e \Leftarrow C'_1$	By i.h.
	$\Psi' \vdash C'_1 \rightarrow C'_2 \bullet e \Rightarrow C'_2$	By Decl \rightarrow App
⊢	$\Psi' \vdash C'_1 \rightarrow A' \bullet e \Rightarrow A'$	$A' = C'_2$
	$\Psi \vdash C'_2 \leq A$	Subderivation
⊢	$\Psi \vdash A' \leq A$	$A' = C'_2$
- **Case**
$$\frac{\Psi \vdash \tau \quad \Psi \vdash [\tau/\beta]B \leq C_0 \rightarrow A}{\Psi \vdash \forall\beta. B \leq C_0 \rightarrow A} \leq \forall L$$

	$\Psi \vdash [\tau/\beta]B \leq C_0 \rightarrow A$	Subderivation
	$\Psi' \vdash [\tau/\beta]B \bullet e \Rightarrow A'$	By i.h. (iii)
⊢	$\Psi \vdash A' \leq A$	"
	$\Psi \vdash \tau$	Subderivation
	$\Psi' \vdash \tau$	By weakening
⊢	$\Psi' \vdash \forall\beta. B \bullet e \Rightarrow A'$	By Decl \forall App

Theorem 3 (Substitution).

Assume $\Psi \vdash e \Rightarrow A$.

- (i) If $\Psi, x : A \vdash e' \Leftarrow C$ then $\Psi \vdash [e/x]e' \Leftarrow C$.
- (ii) If $\Psi, x : A \vdash e' \Rightarrow C$ then $\Psi \vdash [e/x]e' \Rightarrow C$.
- (iii) If $\Psi, x : A \vdash B \bullet e' \Rightarrow C$ then $\Psi \vdash B \bullet [e/x]e' \Rightarrow C$.

Proof. By a straightforward mutual induction on the given derivation. □

Theorem 4 (Inverse Substitution).

Assume $\Psi \vdash e \Leftarrow A$.

- (i) If $\Psi \vdash [(e : A)/x]e' \Leftarrow C$ then $\Psi, x : A \vdash e' \Leftarrow C$.
- (ii) If $\Psi \vdash [(e : A)/x]e' \Rightarrow C$ then $\Psi, x : A \vdash e' \Rightarrow C$.
- (iii) If $\Psi \vdash B \bullet [(e : A)/x]e' \Rightarrow C$ then $\Psi, x : A \vdash B \bullet e' \Rightarrow C$.

Proof. By mutual induction on the given derivation.

- (i) We have $\Psi \vdash [(e : A)/x]e' \Leftarrow C$.

- **Case**
$$\frac{\Psi \vdash [(e : A)/x]e' \Rightarrow B \quad \Psi \vdash B \leq C}{\Psi \vdash [(e : A)/x]e' \Leftarrow C} \text{DeclSub}$$

By i.h. (ii), $\Psi, x : A \vdash e' \Rightarrow B$.
By DeclSub, $\Psi, x : A \vdash e' \Leftarrow C$.

- **Case**

$$\frac{}{\Psi \vdash () \Leftarrow \underbrace{1}_C} \text{Decl11}$$

We have $[(e : A)/x]e' = ()$. Therefore $e' = ()$, and the result follows by Decl11.

- **Case** $\frac{\Psi, \alpha \vdash [(e : A)/x]e' \Leftarrow C'}{\Psi \vdash [(e : A)/x]e' \Leftarrow \forall \alpha. C'} \text{Decl}\forall$

By i.h. (i), $\Psi, \alpha, x : A \vdash e' \Leftarrow C'$.
 By exchange, $\Psi, x : A, \alpha \vdash e' \Leftarrow C'$.
 By Decl \forall , $\Psi, x : A \vdash e' \Leftarrow \forall \alpha. C'$.

- **Case** $\frac{\Psi, y : C_1 \vdash e'' \Leftarrow C_2}{\Psi \vdash \lambda y. e'' \Leftarrow C_1 \rightarrow C_2} \text{Decl}\rightarrow$

We have $[(e : A)/x]e' = \lambda y. e''$.
 By the definition of substitution, $e' = \lambda y. e_0$ and $e'' = [(e : A)/x]e_0$.

$$\begin{array}{ll} \Psi, y : C_1 \vdash e'' \Leftarrow C_2 & \text{Subderivation} \\ \Psi, y : C_1 \vdash [(e : A)/x]e_0 \Leftarrow C_2 & \text{By above equality} \\ \Psi, y : C_1, x : A \vdash e_0 \Leftarrow C_2 & \text{By i.h. (i)} \\ \Psi, x : A, y : C_1 \vdash e_0 \Leftarrow C_2 & \text{By exchange} \\ \Psi, x : A \vdash \underbrace{\lambda y. e_0}_{e'} \Leftarrow \underbrace{C_1 \rightarrow C_2}_C & \text{By Decl}\rightarrow \end{array}$$

(ii) We have $\Psi \vdash [(e : A)/x]e' \Rightarrow C$.

- **Case** $e' = x$:

Note $[(e : A)/x]x = (e : A)$.
 Hence $\Psi \vdash (e : A) \Rightarrow C$; by inversion, $C = A$.
 By Lemma 10 (Well-Formedness), $\Psi \vdash C$, which is $\Psi \vdash A$.
 By DeclAnno, $\Psi \vdash (e : A) \Rightarrow A$.
 By DeclVar, $\Psi, x : A \vdash \underbrace{x}_{e'} \Rightarrow A$.

- **Case** $e' \neq x$:

We now proceed by cases on the derivation of $\Psi \vdash [(e : A)/x]e' \Rightarrow C$.

- **Case** $\frac{(y : C) \in \Psi}{\Psi \vdash y \Rightarrow C} \text{DeclVar}$

Since $[(e : A)/x]e' = y$, we know that $e' = y$.
 By DeclVar, $\Psi, x : A \vdash y \Rightarrow C$.

- **Case** $\frac{\Psi \vdash e'' \Leftarrow C}{\Psi \vdash \underbrace{(e'' : C)}_{[(e:A)/x]e'} \Rightarrow C} \text{DeclAnno}$

We know $[(e : A)/x]e' = (e'' : C)$ and $e' \neq x$.
 Hence there is e_0 such that $e' = (e_0 : C)$ and $[(e : A)/x]e_0 = e''$.

$$\begin{array}{ll} \Psi \vdash e'' \Leftarrow C & \text{Subderivation} \\ \Psi \vdash [(e : A)/x]e_0 \Leftarrow C & \text{By above equality} \\ \Psi, x : A \vdash e_0 \Leftarrow C & \text{By i.h. (i)} \\ \Psi, x : A \vdash C & \text{By Lemma 10 (Well-Formedness)} \\ \Psi, x : A \vdash (e_0 : C) \Rightarrow C & \text{By DeclAnno} \\ \Psi, x : A \vdash e' \Rightarrow C & \text{By above equality} \end{array}$$

- **Case**

$$\frac{}{\Psi \vdash () \Rightarrow 1} \text{Decl}1 \Rightarrow$$

Since $[(e : A)/x]e' = ()$, it follows that $e' = ()$.
 By Decl $1 \Rightarrow$, $\Psi, x : A \vdash () \Rightarrow 1$.

$$\text{-- Case } \frac{\Psi \vdash \sigma \rightarrow \tau \quad \Psi, y : \sigma \vdash e'' \Leftarrow \tau}{\Psi \vdash \lambda y. e'' \Rightarrow \sigma \rightarrow \tau} \text{Decl} \rightarrow I \Rightarrow$$

We have $[(e : A)/x]e' = \lambda y. e''$.

By definition of substitution, there exists e_0 such that $e' = \lambda y. e_0$ and $e'' = [(e : A)/x]e_0$.

So $\Psi, y : \sigma \vdash [(e : A)/x]e_0 \Leftarrow \tau$.

By i.h. (i), $\Psi, y : \sigma, x : A \vdash e_0 \Leftarrow \tau$.

By exchange and $\text{Decl} \rightarrow I$, $\Psi, x : A \vdash \lambda y. e_0 \Leftarrow \sigma \rightarrow \tau$.

Hence $\text{Decl} \rightarrow I \Rightarrow$, $\Psi, x : A \vdash e' \Rightarrow \sigma \rightarrow \tau$.

$$\text{-- Case } \frac{\Psi \vdash e_1 \Rightarrow B \quad \Psi \vdash B \bullet e_2 \Rightarrow C}{\Psi \vdash \underbrace{e_1 e_2}_{[(e:A)/x]e'} \Rightarrow C} \text{Decl} \rightarrow E$$

Note that $[(e : A)/x]e' = e_1 e_2$.

So there exist e'_1, e'_2 such that $e' = e'_1 e'_2$ and $[(e : A)/x]e'_k = e_k$ for $k \in \{1, 2\}$.

Applying these equalities to each subderivation gives

$$\Psi \vdash [(e : A)/x]e'_1 \Rightarrow B \text{ and } \Psi \vdash B \bullet [(e : A)/x]e'_2 \Rightarrow C$$

By i.h. (ii) and (iii), $\Psi, x : A \vdash e'_1 \Rightarrow B$ and $\Psi, x : A \vdash B \bullet e'_2 \Rightarrow C$.

By $\text{Decl} \rightarrow E$, $\Psi, x : A \vdash e'_1 e'_2 \Rightarrow C$, which is $\Psi, x : A \vdash e' \Rightarrow C$.

(iii) We have $\Psi \vdash [(e : A)/x]e' \bullet A \Rightarrow C$.

$$\bullet \text{ Case } \frac{\Psi \vdash \tau \quad \Psi \vdash [\tau/\alpha]B \bullet [(e : A)/x]e' \Rightarrow C}{\Psi \vdash \forall \alpha. B \bullet [(e : A)/x]e' \Rightarrow C} \text{Decl} \forall \text{App}$$

Follows by i.h. (iii) and $\text{Decl} \forall \text{App}$.

$$\bullet \text{ Case } \frac{\Psi \vdash [(e : A)/x]e' \Leftarrow B}{\Psi \vdash B \rightarrow C \bullet [(e : A)/x]e' \Rightarrow C} \text{Decl} \rightarrow \text{App}$$

Follows by i.h. (i) and $\text{Decl} \rightarrow \text{App}$. □

Theorem 5 (Annotation Removal). *We have that:*

- If $\Psi \vdash ((\lambda x. e) : A) \Leftarrow C$ then $\Psi \vdash \lambda x. e \Leftarrow C$.
- If $\Psi \vdash (() : A) \Leftarrow C$ then $\Psi \vdash () \Leftarrow C$.
- If $\Psi \vdash e_1 (e_2 : A) \Rightarrow C$ then $\Psi \vdash e_1 e_2 \Rightarrow C$.
- If $\Psi \vdash (x : A) \Rightarrow A$ then $\Psi \vdash x \Rightarrow B$ and $\Psi \vdash B \leq A$.
- If $\Psi \vdash ((e_1 e_2) : A) \Rightarrow A$ then $\Psi \vdash e_1 e_2 \Rightarrow B$ and $\Psi \vdash B \leq A$.
- If $\Psi \vdash ((e : B) : A) \Rightarrow A$ then $\Psi \vdash (e : B) \Rightarrow B$ and $\Psi \vdash B \leq A$.
- If $\Psi \vdash ((\lambda x. e) : \sigma \rightarrow \tau) \Rightarrow \sigma \rightarrow \tau$ then $\Psi \vdash \lambda x. e \Rightarrow \sigma \rightarrow \tau$.

Proof. All of these follow directly from inversion and Lemma 14 (Subsumption). The one exception is the third, which additionally requires a small induction on the application judgment. □

Theorem 6 (Soundness of Eta).

If $\Psi \vdash \lambda x. e \Leftarrow A$ and $x \notin \text{FV}(e)$, then $\Psi \vdash e \Leftarrow A$.

Proof. By induction on the derivation of $\Psi \vdash \lambda x. e \Leftarrow A$. There are three non-impossible cases:

- **Case**
$$\frac{\Psi, x : B \vdash e x \Leftarrow C}{\Psi \vdash \lambda x. e x \Leftarrow B \rightarrow C} \text{Decl}\rightarrow I$$

We have $\Psi, x : B \vdash e x \Leftarrow C$.

By inversion on DeclSub, we get $\Psi, x : B \vdash e x \Rightarrow C'$ and $\Psi \vdash C' \leq C$.

By inversion on Decl \rightarrow E, we get $\Psi, x : B \vdash e \Rightarrow A'$ and $\Psi, x : B \vdash A' \bullet x \Rightarrow C'$.

By thinning, we know that $\Psi \vdash e \Rightarrow A'$.

By Lemma 12 (Application Subtyping), we get $B' \leq \Psi, x : B \vdash A' \leq B' \rightarrow C'$ and $\Psi, x : B \vdash x \Leftarrow B'$.

By inversion, we know that $\Psi, x : B \vdash x \Rightarrow B$ and $\Psi \vdash B \leq B'$.

By $\leq \rightarrow$, $\Psi, x : B \vdash B' \rightarrow C' \leq B \rightarrow C$.

Hence by Lemma 6 (Transitivity of Declarative Subtyping), $\Psi, x : B \vdash A' \leq B \rightarrow C$.

Hence $\Psi \vdash A' \leq B \rightarrow C$.

By DeclSub, $\Psi \vdash e \Leftarrow B \rightarrow C$.

- **Case**
$$\frac{\Psi, \alpha \vdash \lambda x. e x \Leftarrow B}{\Psi \vdash \lambda x. e x \Leftarrow \forall \alpha. B} \text{Decl}\forall I$$

By induction, $\Psi, \alpha \vdash \lambda x. e x \Leftarrow B$.

By Decl \forall I, $\Psi \vdash \lambda x. e x \Leftarrow \forall \alpha. B$.

- **Case**
$$\frac{\Psi \vdash \lambda x. e x \Rightarrow B \quad \Psi \vdash B \leq A}{\Psi \vdash \lambda x. e x \Leftarrow A} \text{DeclSub}$$

We have $\Psi \vdash \lambda x. e x \Rightarrow B$ and $\Psi \vdash B \leq A$.

By inversion on Decl \rightarrow I \Rightarrow , $\Psi, x : \sigma \vdash e x \Leftarrow \tau$ and $B = \sigma \rightarrow \tau$.

By inversion on DeclSub, we get $\Psi, x : \sigma \vdash e x \Rightarrow C_2$ and $\Psi \vdash C_2 \leq \tau$.

By inversion on Decl \rightarrow E, we get $\Psi, x : \sigma \vdash e \Rightarrow C$ and $\Psi, x : \sigma \vdash C \bullet x \Rightarrow C_2$.

By thinning, we know that $\Psi \vdash e \Rightarrow C$.

By Lemma 12 (Application Subtyping), we get C_1 such that $\Psi, x : \sigma \vdash C \leq C_1 \rightarrow C_2$ and $\Psi, x : \sigma \vdash x \Leftarrow C_1$.

By inversion on DeclSub, $\Psi, x : \sigma \vdash x \Rightarrow \sigma$ and $\Psi \vdash \sigma \leq C_1$.

By $\leq \rightarrow$, $\Psi, x : \sigma \vdash C_1 \rightarrow C_2 \leq \sigma \rightarrow \tau$.

Hence by Lemma 6 (Transitivity of Declarative Subtyping), $\Psi, x : \sigma \vdash C \leq \sigma \rightarrow \tau$.

Hence $\Psi \vdash C \leq \sigma \rightarrow \tau$.

Hence by Lemma 6 (Transitivity of Declarative Subtyping), $\Psi \vdash C \leq A$.

By DeclSub, $\Psi \vdash e \Leftarrow A$. □

D' Properties of Context Extension

D'.1 Syntactic Properties

Lemma 15 (Declaration Preservation). *If $\Gamma \longrightarrow \Delta$, and u is a variable or marker $\blacktriangleright_{\alpha}$ declared in Γ , then u is declared in Δ .*

Proof. By a routine induction on $\Gamma \longrightarrow \Delta$. □

Lemma 16 (Declaration Order Preservation). *If $\Gamma \longrightarrow \Delta$ and u is declared to the left of v in Γ , then u is declared to the left of v in Δ .*

Proof. By induction on the derivation of $\Gamma \longrightarrow \Delta$.

- **Case**
$$\frac{}{\cdot \longrightarrow \cdot} \rightarrow ID$$

This case is impossible.

- **Case**
$$\frac{\Gamma \longrightarrow \Delta}{\Gamma, x : A \longrightarrow \Delta, x : A} \longrightarrow \text{Var}$$

There are two cases, depending on whether or not $v = x$.

- Case $v = x$:
Since u is declared to the left of v , u is declared in Γ .
By Lemma 15 (Declaration Preservation), u is declared in Δ .
Hence u is declared to the left of x in $\Delta, x : A$.
- Case $v \neq x$:
Then v is declared in Γ , and u is declared to the left of v in Γ .
By induction, u is declared to the left of v in Δ .
Hence u is declared to the left of v in $\Delta, x : A$.

- **Case**
$$\frac{\Gamma \longrightarrow \Delta}{\Gamma, \alpha \longrightarrow \Delta, \alpha} \longrightarrow \text{Uvar}$$

This case is similar to the $\longrightarrow \text{Var}$ case.

- **Case**
$$\frac{\Gamma \longrightarrow \Delta}{\Gamma, \hat{\alpha} \longrightarrow \Delta, \hat{\alpha}} \longrightarrow \text{Unsolved}$$

This case is similar to the $\longrightarrow \text{Var}$ case.

- **Case**
$$\frac{\Gamma \longrightarrow \Delta \quad [\Delta]\tau = [\Delta]\tau'}{\Gamma, \hat{\alpha} = \tau \longrightarrow \Delta, \hat{\alpha} = \tau'} \longrightarrow \text{Solved}$$

This case is similar to the $\longrightarrow \text{Var}$ case.

- **Case**
$$\frac{\Gamma \longrightarrow \Delta}{\Gamma, \blacktriangleright \hat{\alpha} \longrightarrow \Delta, \blacktriangleright \hat{\alpha}} \longrightarrow \text{Marker}$$

This case is similar to the $\longrightarrow \text{Var}$ case.

- **Case**
$$\frac{\Gamma \longrightarrow \Delta}{\Gamma, \hat{\alpha} \longrightarrow \Delta, \hat{\alpha} = \tau} \longrightarrow \text{Solve}$$

This case is similar to the $\longrightarrow \text{Var}$ case.

- **Case**
$$\frac{\Gamma \longrightarrow \Delta}{\Gamma \longrightarrow \Delta, \hat{\alpha}} \longrightarrow \text{Add}$$

By induction, u is declared to the left of v in Δ .
Therefore u is declared to the left of v in $\Delta, \hat{\alpha}$.

- **Case**
$$\frac{\Gamma \longrightarrow \Delta}{\Gamma \longrightarrow \Delta, \hat{\alpha} = \tau} \longrightarrow \text{AddSolved}$$

By induction, u is declared to the left of v in Δ .

Therefore u is declared to the left of v in $\Delta, \hat{\alpha} = \tau$. □

Lemma 17 (Reverse Declaration Order Preservation). *If $\Gamma \longrightarrow \Delta$ and u and v are both declared in Γ and u is declared to the left of v in Δ , then u is declared to the left of v in Γ .*

Proof. It is given that u and v are declared in Γ . Either u is declared to the left of v in Γ , or v is declared to the left of u . Suppose the latter (for a contradiction). By Lemma 16 (Declaration Order Preservation), v is declared to the left of u in Δ . But we know that u is declared to the left of v in Δ : contradiction. Therefore u is declared to the left of v in Γ . □

Lemma 18 (Substitution Extension Invariance). *If $\Theta \vdash A$ and $\Theta \longrightarrow \Gamma$ then $[\Gamma]A = [\Gamma](\Theta A)$ and $[\Gamma]A = [\Theta](\Gamma A)$.*

Proof. To show that $[\Gamma]A = [\Theta](\Gamma A)$, observe that $\Theta \vdash A$, and that by definition of $\Theta \longrightarrow \Gamma$, every solved variable in Θ is solved in Γ . Therefore $[\Theta](\Gamma A) = [\Gamma]A$, since $\text{unsolved}(\Gamma A)$ contains no variables that Θ solves.

To show that $[\Gamma]A = [\Gamma](\Theta A)$, we proceed by induction on $|\Gamma \vdash A|$.

- **Case** $\frac{\alpha \in \Theta}{\Theta \vdash \alpha}$

Note that $[\Gamma]\alpha = \alpha = [\Theta]\alpha$, so $[\Gamma]\alpha = [\Gamma](\Theta)\alpha$.

- **Case** $\frac{\Theta \vdash A \quad \Theta \vdash B}{\Theta \vdash A \rightarrow B}$

By induction, $[\Gamma]A = [\Gamma](\Theta)A$.

By induction, $[\Gamma]B = [\Gamma](\Theta)B$.

Then

$$\begin{aligned} [\Gamma](A \rightarrow B) &= [\Gamma]A \rightarrow [\Gamma]B && \text{By definition of substitution} \\ &= [\Gamma](\Theta)A \rightarrow [\Gamma](\Theta)B && \text{By induction hypothesis (twice)} \\ &= [\Gamma](\Theta)(A \rightarrow B) && \text{By definition of substitution} \\ &= [\Gamma](\Theta)(A \rightarrow B) && \text{By definition of substitution} \end{aligned}$$

- **Case** $\frac{\Theta, \alpha \vdash A}{\Theta \vdash \forall \alpha. A}$

By inversion, we have $\Theta, \alpha \vdash A$.

By rule $\longrightarrow \text{Uvar}$, $\Theta, \alpha \longrightarrow \Gamma, \alpha$.

By induction, $[\Gamma, \alpha]A = [\Gamma, \alpha](\Theta, \alpha)A$.

By definition, $[\Gamma]A = [\Gamma](\Theta)A$.

Then

$$\begin{aligned} [\Gamma]\forall \alpha. A &= \forall \alpha. [\Gamma]A && \text{By definition} \\ &= \forall \alpha. [\Gamma](\Theta)A && \text{By conclusion above} \\ &= [\Gamma](\forall \alpha. (\Theta)A) && \text{By definition} \\ &= [\Gamma](\Theta)(\forall \alpha. A) && \text{By definition} \\ &= [\Gamma, \alpha](\Theta, \alpha)(\forall \alpha. A) && \text{By definition} \end{aligned}$$

- **Case**

$$\frac{\Theta_0, \hat{\alpha}, \Theta_1 \vdash \hat{\alpha}}{\Theta}$$

Note that $[\Theta]\hat{\alpha} = \hat{\alpha}$.

Hence $[\Gamma](\Theta)\hat{\alpha} = [\Gamma]\hat{\alpha}$.

- **Case**

$$\frac{\Theta_0, \hat{\alpha} = \tau, \Theta_1 \vdash \hat{\alpha}}{\Theta}$$

From $\Theta \longrightarrow \Gamma$, By a nested induction we get $\Gamma = \Gamma_0, \hat{\alpha} = \tau', \Gamma_1$, and $[\Gamma]\tau' = [\Gamma]\tau$.

Note that $|\Theta \vdash \tau| < |\Theta \vdash \hat{\alpha}|$.

By induction, $[\Gamma]\tau = [\Gamma](\Theta)\tau$.

Hence

$$\begin{aligned} [\Gamma]\hat{\alpha} &= [\Gamma]\tau' && \text{By definition} \\ &= [\Gamma]\tau && \text{From the extension judgment} \\ &= [\Gamma](\Theta)\tau && \text{From the induction hypothesis} \\ &= [\Gamma](\Theta)\hat{\alpha} && \text{By definition} \end{aligned}$$

□

Lemma 19 (Extension Equality Preservation).

If $\Gamma \vdash A$ and $\Gamma \vdash B$ and $[\Gamma]A = [\Gamma]B$ and $\Gamma \longrightarrow \Delta$, then $[\Delta]A = [\Delta]B$.

Proof. By induction on the derivation of $\Gamma \longrightarrow \Delta$.

• **Case**

$$\frac{\Gamma \longrightarrow \Delta}{\Gamma \longrightarrow \Delta} \rightarrow \text{ID}$$

We have $[\Gamma]A = [\Gamma]B$, but $\Gamma = \Delta$, so $[\Delta]A = [\Delta]B$.

• **Case**

$$\frac{\Gamma' \longrightarrow \Delta'}{\Gamma', x : C \longrightarrow \Delta', x : C} \rightarrow \text{Var}$$

We have $[\Gamma', x : C]A = [\Gamma', x : C]B$.

By definition of substitution, $[\Gamma']A = [\Gamma']B$.

By i.h., $[\Delta']A = [\Delta']B$.

By definition of substitution, $[\Delta', x : C]A = [\Delta', x : C]B$.

• **Case**

$$\frac{\Gamma' \longrightarrow \Delta'}{\Gamma', \alpha \longrightarrow \Delta', \alpha} \rightarrow \text{Uvar}$$

We have $[\Gamma', \alpha]A = [\Gamma', \alpha]B$.

By definition of substitution, $[\Gamma']A = [\Gamma']B$.

By i.h., $[\Delta']A = [\Delta']B$.

By definition of substitution, $[\Delta', \alpha]A = [\Delta', \alpha]B$.

• **Case**

$$\frac{\Gamma' \longrightarrow \Delta'}{\Gamma', \hat{\alpha} \longrightarrow \Delta', \hat{\alpha}} \rightarrow \text{Unsolved}$$

Similar to the $\rightarrow \text{Uvar}$ case.

• **Case**

$$\frac{\Gamma' \longrightarrow \Delta'}{\Gamma', \blacktriangleright_{\hat{\alpha}} \longrightarrow \Delta', \blacktriangleright_{\hat{\alpha}}} \rightarrow \text{Marker}$$

Similar to the $\rightarrow \text{Uvar}$ case.

• **Case**

$$\frac{\Gamma \longrightarrow \Delta'}{\Gamma \longrightarrow \Delta', \hat{\alpha}} \rightarrow \text{Add}$$

We have $[\Gamma]A = [\Gamma]B$.

By i.h., $[\Delta']A = [\Delta']B$.

By definition of substitution, $[\Delta', \hat{\alpha}]A = [\Delta', \hat{\alpha}]B$.

• **Case**

$$\frac{\Gamma \longrightarrow \Delta'}{\Gamma \longrightarrow \Delta', \hat{\alpha} = \tau} \rightarrow \text{AddSolved}$$

We have $[\Gamma]A = [\Gamma]B$.

By i.h., $[\Delta']A = [\Delta']B$.

We implicitly assume that Δ is well-formed, so $\hat{\alpha} \notin \text{dom}(\Delta')$.

Since $\Gamma \longrightarrow \Delta'$ and $\hat{\alpha} \notin \text{dom}(\Delta')$, it follows that $\hat{\alpha} \notin \text{dom}(\Gamma)$.

We have $\Gamma \vdash A$ and $\Gamma \vdash B$, so $\hat{\alpha} \notin (\text{FV}(A) \cup \text{FV}(B))$.

Therefore, by definition of substitution, $[\Delta', \hat{\alpha} = \tau]A = [\Delta', \hat{\alpha} = \tau]B$.

- **Case** $\frac{\Gamma' \longrightarrow \Delta' \quad [\Delta']\tau = [\Delta']\tau'}{\Gamma', \hat{\alpha} = \tau \longrightarrow \Delta', \hat{\alpha} = \tau'} \longrightarrow \text{Solved}$

We have $[\Gamma', \hat{\alpha} = \tau]A = [\Gamma', \hat{\alpha} = \tau]B$.

By definition, $[\Gamma', \hat{\alpha} = \tau]A = [\Gamma', \hat{\alpha} = \tau]\tau$, but we implicitly assume that Γ is well-formed, so $\hat{\alpha} \notin \text{FV}(\tau)$, so actually $[\Gamma', \hat{\alpha} = \tau]A = [\Gamma']\tau$.

Combined with similar reasoning for B , we get

$$[\Gamma'][\tau/\hat{\alpha}]A = [\Gamma'][\tau/\hat{\alpha}]B$$

By i.h., $[\Delta'][\tau/\hat{\alpha}]A = [\Delta'][\tau/\hat{\alpha}]B$.

By distributivity of substitution, $[[\Delta']\tau/\hat{\alpha}][\Delta']A = [[\Delta']\tau/\hat{\alpha}][\Delta']B$.

Using the premise $[\Delta']\tau = [\Delta']\tau'$, we get $[[\Delta']\tau'/\hat{\alpha}][\Delta']A = [[\Delta']\tau'/\hat{\alpha}][\Delta']B$.

By distributivity of substitution (in the other direction), $[\Delta'][\tau'/\hat{\alpha}]A = [\Delta'][\tau'/\hat{\alpha}]B$.

It follows from the definition of substitution that $[\Delta', \hat{\alpha} = \tau']A = [\Delta', \hat{\alpha} = \tau']B$.

- **Case** $\frac{\Gamma' \longrightarrow \Delta'}{\Gamma', \hat{\alpha} \longrightarrow \Delta', \hat{\alpha} = \tau} \longrightarrow \text{Solve}$

We have $[\Gamma', \hat{\alpha}]A = [\Gamma', \hat{\alpha}]B$.

By definition of substitution, $[\Gamma']A = [\Gamma']B$.

By i.h., $[\Delta'][\tau/\hat{\alpha}]A = [\Delta'][\tau/\hat{\alpha}]B$.

It follows from the definition of substitution that $[\Delta', \hat{\alpha} = \tau]A = [\Delta', \hat{\alpha} = \tau]B$. □

Lemma 20 (Reflexivity). *If Γ is well-formed, then $\Gamma \longrightarrow \Gamma$.*

Proof. By induction on the structure of Γ .

- **Case** $\Gamma = \cdot$: Apply rule $\longrightarrow \text{ID}$.
- **Case** $\Gamma = (\Gamma', \alpha)$: By i.h., $\Gamma' \longrightarrow \Gamma'$. By rule $\longrightarrow \text{Uvar}$, we get $\Gamma', \alpha \longrightarrow \Gamma', \alpha$.
- **Case** $\Gamma = (\Gamma', \hat{\alpha})$: By i.h., $\Gamma' \longrightarrow \Gamma'$. By rule $\longrightarrow \text{Unsolved}$, we get $\Gamma', \hat{\alpha} \longrightarrow \Gamma', \hat{\alpha}$.
- **Case** $\Gamma = (\Gamma', \hat{\alpha} = \tau)$:
By i.h., $\Gamma' \longrightarrow \Gamma'$.
Clearly, $[\Gamma']\tau = [\Gamma']\tau$, so we can apply $\longrightarrow \text{Solved}$ to get $\Gamma', \hat{\alpha} = \tau \longrightarrow \Gamma', \hat{\alpha} = \tau$.
- **Case** $\Gamma = (\Gamma', \blacktriangleright_{\hat{\alpha}})$: By i.h., $\Gamma' \longrightarrow \Gamma'$. By rule $\longrightarrow \text{Marker}$, we get $\Gamma', \blacktriangleright_{\hat{\alpha}} \longrightarrow \Gamma', \blacktriangleright_{\hat{\alpha}}$. □

Lemma 21 (Transitivity). *If $\Gamma \longrightarrow \Delta$ and $\Delta \longrightarrow \Theta$, then $\Gamma \longrightarrow \Theta$.*

Proof. By induction on the derivation of $\Delta \longrightarrow \Theta$.

- **Case** $\longrightarrow \text{ID}$:
In this case $\Theta = \Delta$.
Hence $\Gamma \longrightarrow \Delta$ suffices.
- **Case** $\frac{\Delta' \longrightarrow \Theta'}{\Delta', \alpha \longrightarrow \Theta', \alpha} \longrightarrow \text{Uvar}$

We have $\Delta = (\Delta', \alpha)$ and $\Theta = (\Theta', \alpha)$.

By inversion on $\Gamma \longrightarrow \Delta$, we have $\Gamma = (\Gamma', \alpha)$ and $\Gamma' \longrightarrow \Delta'$.

By i.h., $\Gamma' \longrightarrow \Theta'$.

Applying rule $\longrightarrow \text{Uvar}$ gives $\Gamma', \alpha \longrightarrow \Theta', \alpha$.

• **Case**
$$\frac{\Delta' \longrightarrow \Theta'}{\Delta', \hat{\alpha} \longrightarrow \Theta', \hat{\alpha}} \longrightarrow \text{Uvar}$$

We have $\Delta = (\Delta', \hat{\alpha})$ and $\Theta = (\Theta', \hat{\alpha})$.

Either of two rules could have derived $\Gamma \longrightarrow \Delta$:

– **Case**
$$\frac{\Gamma' \longrightarrow \Delta'}{\Gamma', \hat{\alpha} \longrightarrow \Delta', \hat{\alpha}} \longrightarrow \text{Unsolved}$$

Here we have $\Gamma = (\Gamma', \hat{\alpha})$ and $\Gamma' \longrightarrow \Delta'$.

By i.h., $\Gamma' \longrightarrow \Theta'$.

Applying rule $\longrightarrow \text{Unsolved}$ gives $\Gamma', \hat{\alpha} \longrightarrow \Theta', \hat{\alpha}$.

– **Case**
$$\frac{\Gamma \longrightarrow \Delta'}{\Gamma \longrightarrow \Delta', \hat{\alpha}} \longrightarrow \text{Add}$$

By i.h., $\Gamma \longrightarrow \Theta'$.

By rule $\longrightarrow \text{Add}$, we get $\Gamma \longrightarrow \Theta', \hat{\alpha}$.

• **Case**
$$\frac{\Delta' \longrightarrow \Theta' \quad [\Theta']\tau_1 = [\Theta']\tau_2}{\Delta', \hat{\alpha} = \tau_1 \longrightarrow \Theta', \hat{\alpha} = \tau_2} \longrightarrow \text{Solved}$$

In this case $\Delta = (\Delta', \hat{\alpha} = \tau_1)$ and $\Theta = (\Theta', \hat{\alpha} = \tau_2)$.

One of three rules must have derived $\Gamma \longrightarrow \Delta', \hat{\alpha} = \tau$:

– **Case**
$$\frac{\Gamma' \longrightarrow \Delta' \quad [\Delta']\tau_0 = [\Delta']\tau_1}{\Gamma', \hat{\alpha} = \tau_0 \longrightarrow \Delta', \hat{\alpha} = \tau_1} \longrightarrow \text{Solved}$$

Here, $\Gamma = (\Gamma', \hat{\alpha} = \tau_0)$ and $\Delta = (\Delta', \hat{\alpha} = \tau_1)$.

By i.h., we have $\Gamma' \longrightarrow \Theta'$.

The premises of the respective \longrightarrow derivations give us $[\Delta']\tau_0 = [\Delta']\tau_1$ and $[\Theta']\tau_1 = [\Theta']\tau_2$.

We know that $\Gamma' \vdash \tau_0$ and $\Delta' \vdash \tau_1$ and $\Theta' \vdash \tau_2$.

By extension weakening (Lemma 25 (Extension Weakening)), $\Theta' \vdash \tau_0$.

By extension weakening (Lemma 25 (Extension Weakening)), $\Theta' \vdash \tau_1$.

Since $[\Delta']\tau_0 = [\Delta']\tau_1$, we know that $[\Theta'][\Delta']\tau_0 = [\Theta'][\Delta']\tau_1$.

By Lemma 18 (Substitution Extension Invariance), $[\Theta'][\Delta']\tau_0 = [\Theta']\tau_0$.

By Lemma 18 (Substitution Extension Invariance), $[\Theta'][\Delta']\tau_1 = [\Theta']\tau_1$.

So $[\Theta']\tau_0 = [\Theta']\tau_1$.

Hence by transitivity of equality, $[\Theta']\tau_0 = [\Theta']\tau_1 = [\Theta']\tau_2$.

By rule $\longrightarrow \text{Solved}$, $\Gamma', \hat{\alpha} = \tau \longrightarrow \Theta', \hat{\alpha} = \tau_2$.

– **Case**
$$\frac{\Gamma \longrightarrow \Delta'}{\Gamma \longrightarrow \Delta', \hat{\alpha} = \tau_1} \longrightarrow \text{AddSolved}$$

By induction, we have $\Gamma \longrightarrow \Theta'$.

By rule $\longrightarrow \text{AddSolved}$, we get $\Gamma \longrightarrow \Theta', \hat{\alpha} = \tau_2$.

– **Case**
$$\frac{\Gamma' \longrightarrow \Delta'}{\Gamma', \hat{\alpha} \longrightarrow \Delta', \hat{\alpha} = \tau_1} \longrightarrow \text{Solve}$$

We have $\Gamma = (\Gamma', \hat{\alpha})$.

By induction, $\Gamma' \longrightarrow \Theta'$.

By rule $\longrightarrow \text{Solve}$, we get $\Gamma', \hat{\alpha} \longrightarrow \Theta', \hat{\alpha} = \tau_2$.

• **Case**
$$\frac{\Delta' \longrightarrow \Theta'}{\Delta', \blacktriangleright_{\hat{\alpha}} \longrightarrow \Theta', \blacktriangleright_{\hat{\alpha}}} \longrightarrow \text{Marker}$$

In this case we know $\Delta = (\Delta', \blacktriangleright_{\hat{\alpha}})$ and $\Theta = (\Theta', \blacktriangleright_{\hat{\alpha}})$.

Since $\Delta = (\Delta', \blacktriangleright_{\hat{\alpha}})$, only $\longrightarrow \text{Marker}$ could derive $\Gamma \longrightarrow \Delta$, so by inversion, $\Gamma = (\Gamma', \blacktriangleright_{\hat{\alpha}})$ and $\Gamma' \longrightarrow \Delta'$.

By induction, we have $\Gamma' \longrightarrow \Theta'$.

Applying rule $\longrightarrow \text{Marker}$ gives $\Gamma', \blacktriangleright_{\hat{\alpha}} \longrightarrow \Theta', \blacktriangleright_{\hat{\alpha}}$.

• **Case**
$$\frac{\Delta \longrightarrow \Theta'}{\Delta \longrightarrow \Theta', \hat{\alpha}} \longrightarrow \text{Add}$$

In this case, we have $\Theta = (\Theta', \hat{\alpha})$.

By induction, we get $\Gamma \longrightarrow \Theta'$.

By rule $\longrightarrow \text{Add}$, we get $\Gamma \longrightarrow \Theta', \hat{\alpha}$.

• **Case**
$$\frac{\Delta \longrightarrow \Theta'}{\Delta \longrightarrow \Theta', \hat{\alpha} = \tau} \longrightarrow \text{AddSolved}$$

In this case, we have $\Theta = (\Theta', \hat{\alpha} = \tau)$.

By induction, we get $\Gamma \longrightarrow \Theta'$.

By rule $\longrightarrow \text{AddSolved}$, we get $\Gamma \longrightarrow \Theta', \hat{\alpha} = \tau$.

• **Case**
$$\frac{\Delta' \longrightarrow \Theta'}{\Delta', \hat{\alpha} \longrightarrow \Theta', \hat{\alpha} = \tau} \longrightarrow \text{Solve}$$

In this case, we have $\Delta = (\Delta', \hat{\alpha})$ and $\Theta = (\Theta', \hat{\alpha} = \tau)$.

One of two rules could have derived $\Gamma \longrightarrow \Delta', \hat{\alpha}$:

– **Case**
$$\frac{\Gamma' \longrightarrow \Delta'}{\Gamma', \hat{\alpha} \longrightarrow \Delta', \hat{\alpha}} \longrightarrow \text{Unsolved}$$

In this case, we have $\Gamma = (\Gamma', \hat{\alpha})$ and $\Gamma' \longrightarrow \Delta'$ and $\Delta' \longrightarrow \Theta'$.

By induction, we have $\Gamma' \longrightarrow \Theta'$.

By rule $\longrightarrow \text{Solve}$, we get $\Gamma', \hat{\alpha} \longrightarrow \Theta', \hat{\alpha} = \tau$.

– **Case**
$$\frac{\Gamma \longrightarrow \Delta'}{\Gamma \longrightarrow \Delta', \hat{\alpha}} \longrightarrow \text{Add}$$

In this case, we have $\Gamma \longrightarrow \Delta'$ and $\Delta' \longrightarrow \Theta'$.

By induction, we have $\Gamma \longrightarrow \Theta'$.

By rule $\longrightarrow \text{Solve}$, we get $\Gamma \longrightarrow \Theta', \hat{\alpha} = \tau$. □

Lemma 22 (Right Softness). *If $\Gamma \longrightarrow \Delta$ and Θ is soft (and (Δ, Θ) is well-formed) then $\Gamma \longrightarrow \Delta, \Theta$.*

Proof. By induction on Θ , applying rules $\longrightarrow \text{Add}$ and $\longrightarrow \text{AddSolved}$ as needed. □

Lemma 23 (Evar Input).

If $\Gamma, \hat{\alpha} \longrightarrow \Delta$ then $\Delta = (\Delta_0, \Delta_{\hat{\alpha}}, \Theta)$ where $\Gamma \longrightarrow \Delta_0$, and $\Delta_{\hat{\alpha}}$ is either $\hat{\alpha}$ or $\hat{\alpha} = \tau$, and Θ is soft.

Proof. By induction on the given derivation.

• **Cases** $\longrightarrow \text{ID}$, $\longrightarrow \text{Var}$, $\longrightarrow \text{Uvar}$, $\longrightarrow \text{Solved}$, $\longrightarrow \text{Marker}$:

Impossible: the left-hand context cannot have the form $\Gamma, \hat{\alpha}$.

• **Case**
$$\frac{\Gamma \longrightarrow \Delta_0}{\Gamma, \hat{\alpha} \longrightarrow \underbrace{\Delta_0, \hat{\alpha}}_{\Delta}} \longrightarrow \text{Unsolved}$$

Let $\Theta = \cdot$, which is vacuously soft. Therefore $\Delta = (\Delta_0, \hat{\alpha}) = (\Delta_0, \hat{\alpha}, \Theta)$; the subderivation is the rest of the result.

• **Case**
$$\frac{\Gamma \longrightarrow \Delta_0}{\Gamma, \hat{\alpha} \longrightarrow \underbrace{\Delta_0, \hat{\alpha} = \tau}_{\Delta}} \longrightarrow \text{Solve}$$

Let $\Theta = \cdot$, which is vacuously soft. Therefore $\Delta = (\Delta_0, \hat{\alpha}) = (\Delta_0, \hat{\alpha} = \tau, \Theta)$; the subderivation is the rest of the result.

• **Case**
$$\frac{\Gamma, \hat{\alpha} \longrightarrow \Delta_0}{\Gamma, \hat{\alpha} \longrightarrow \underbrace{\Delta_0, \hat{\beta}}_{\Delta}} \longrightarrow \text{Add}$$

Suppose $\hat{\beta} = \hat{\alpha}$.

We have $\Gamma, \hat{\alpha} \longrightarrow \Delta_0$. By Lemma 15 (Declaration Preservation), $\hat{\alpha}$ is declared in Δ_0 .

But then $(\Delta_0, \hat{\beta}) = (\Delta_0, \hat{\alpha})$ with multiple $\hat{\alpha}$ declarations,

which violates the implicit assumption that Δ is well-formed. Contradiction.

Therefore $\hat{\beta} \neq \hat{\alpha}$.

By i.h., $\Delta' = (\Delta_0, \Delta_{\hat{\alpha}}, \Theta')$ where $\Gamma \longrightarrow \Delta_0$ and Θ' is soft.

Let $\Theta = (\Theta', \hat{\beta})$. Therefore $(\Delta', \hat{\beta}) = (\Delta_0, \Delta_{\hat{\alpha}}, \Theta', \hat{\beta})$. As Θ' is soft, $(\Theta', \hat{\beta})$ is soft. Since $\Delta = (\Delta', \hat{\beta})$, this gives $\Delta = (\Delta_0, \Delta_{\hat{\alpha}}, \Theta)$.

• **Case** $\longrightarrow \text{AddSolved}$: Similar to the case for $\longrightarrow \text{Add}$. □

Lemma 24 (Extension Order).

(i) If $\Gamma_L, \alpha, \Gamma_R \longrightarrow \Delta$ then $\Delta = (\Delta_L, \alpha, \Delta_R)$ where $\Gamma_L \longrightarrow \Delta_L$.
Moreover, if Γ_R is soft then Δ_R is soft.

(ii) If $\Gamma_L, \blacktriangleright_{\hat{\alpha}}, \Gamma_R \longrightarrow \Delta$ then $\Delta = (\Delta_L, \blacktriangleright_{\hat{\alpha}}, \Delta_R)$ where $\Gamma_L \longrightarrow \Delta_L$.
Moreover, if Γ_R is soft then Δ_R is soft.

(iii) If $\Gamma_L, \hat{\alpha}, \Gamma_R \longrightarrow \Delta$ then $\Delta = \Delta_L, \Theta, \Delta_R$ where $\Gamma_L \longrightarrow \Delta_L$ and Θ is either $\hat{\alpha}$ or $\hat{\alpha} = \tau$ for some τ .

(iv) If $\Gamma_L, \hat{\alpha} = \tau, \Gamma_R \longrightarrow \Delta$ then $\Delta = \Delta_L, \hat{\alpha} = \tau', \Delta_R$ where $\Gamma_L \longrightarrow \Delta_L$ and $[\Delta_L]\tau = [\Delta_L]\tau'$.

(v) If $\Gamma_L, x : A, \Gamma_R \longrightarrow \Delta$ then $\Delta = (\Delta_L, x : A', \Delta_R)$ where $\Gamma_L \longrightarrow \Delta_L$ and $[\Delta_L]A = [\Delta_L]A'$.
Moreover, Γ_R is soft if and only if Δ_R is soft.

Proof. (i) By induction on the derivation of $\Gamma_L, \alpha, \Gamma_R \longrightarrow \Delta$.

• **Case**

$$\frac{}{\cdot \longrightarrow \cdot} \longrightarrow \text{ID}$$

This case is impossible since $(\Gamma_L, \alpha, \Gamma_R)$ cannot have the form \cdot .

• **Cases** $\longrightarrow \text{Uvar}$:

We have two cases, depending on whether or not the rightmost variable is α .

– **Case**
$$\frac{\Gamma \longrightarrow \Delta'}{\Gamma, \alpha \longrightarrow \Delta', \alpha} \longrightarrow \text{Uvar}$$

Let $\Delta_L = \Delta'$, and let $\Delta_R = \cdot$ (which is soft).

We have $\Gamma \longrightarrow \Delta'$, which is $\Gamma_L \longrightarrow \Delta_L$.

$$\text{-- Case } \frac{\Gamma_L, \alpha, \Gamma'_R \longrightarrow \Delta'}{\Gamma_L, \alpha, \underbrace{\Gamma'_R, \beta}_{\Gamma_R} \longrightarrow \underbrace{\Delta', \beta}_{\Delta}} \longrightarrow \text{Uvar}$$

By i.h., $\Delta' = (\Delta_L, \alpha, \Delta'_R)$ where $\Gamma_L \longrightarrow \Delta_L$.

Hence $\Delta = (\Delta_L, \alpha, \Delta'_R, \beta)$.

(Since $\beta \in \Gamma_R$, it cannot be the case that Γ_R is soft.)

$$\bullet \text{ Case } \frac{\Gamma_L, \alpha, \Gamma'_R \longrightarrow \Delta'}{\Gamma_L, \alpha, \underbrace{\Gamma'_R, x : A}_{\Gamma_R} \longrightarrow \underbrace{\Delta', x : A}_{\Delta}} \longrightarrow \text{Var}$$

By i.h., $\Delta' = (\Delta_L, \alpha, \Delta'_R)$ where $\Gamma_L \longrightarrow \Delta_L$.

Hence $\Delta = (\Delta_L, \alpha, \Delta'_R, x : A)$.

(Since $x : A \in \Gamma_R$, it cannot be the case that Γ_R is soft.)

$$\bullet \text{ Case } \frac{\Gamma_L, \alpha, \Gamma'_R \longrightarrow \Delta'}{\Gamma_L, \alpha, \underbrace{\Gamma'_R, \hat{\alpha}}_{\Gamma_R} \longrightarrow \underbrace{\Delta', \hat{\alpha}}_{\Delta}} \longrightarrow \text{Unsolved}$$

By i.h., $\Delta' = (\Delta_L, \alpha, \Delta'_R)$ where $\Gamma_L \longrightarrow \Delta_L$.

Hence $\Delta = (\Delta_L, \alpha, \Delta'_R, \hat{\alpha})$.

(If Γ_R is soft, by i.h. Δ'_R is soft, so $\Delta_R = (\Delta'_R, \hat{\alpha})$ is soft.)

$$\bullet \text{ Case } \frac{\Gamma_L, \alpha, \Gamma'_R \longrightarrow \Delta'}{\Gamma_L, \alpha, \underbrace{\Gamma'_R, \blacktriangleright \beta}_{\Gamma'_R} \longrightarrow \underbrace{\Delta', \blacktriangleright \beta}_{\Delta}} \longrightarrow \text{Marker}$$

By i.h., $\Delta' = (\Delta_L, \alpha, \Delta'_R)$ where $\Gamma_L \longrightarrow \Delta_L$.

Hence $\Delta = (\Delta_L, \alpha, \Delta'_R, \blacktriangleright \beta)$.

(Since $\blacktriangleright \beta \in \Gamma_R$, it cannot be the case that Γ_R is soft.)

$$\bullet \text{ Case } \frac{\Gamma_L, \alpha, \Gamma'_R \longrightarrow \Delta' \quad [\Delta']\tau = [\Delta']\tau'}{\Gamma_L, \alpha, \underbrace{\Gamma'_R, \hat{\alpha} = \tau}_{\Gamma_R} \longrightarrow \underbrace{\Delta', \hat{\alpha} = \tau'}_{\Delta'}} \longrightarrow \text{Solved}$$

By i.h., $\Delta' = (\Delta_L, \alpha, \Delta'_R)$ where $\Gamma_L \longrightarrow \Delta_L$.

Hence $\Delta = (\Delta_L, \alpha, \Delta'_R, \hat{\alpha} = \tau')$.

(If Γ_R is soft, by i.h. Δ'_R is soft, so $\Delta_R = (\Delta'_R, \hat{\alpha} = \tau)$ is soft.)

$$\bullet \text{ Case } \frac{\Gamma_L, \alpha, \Gamma'_R \longrightarrow \Delta'}{\Gamma_L, \alpha, \underbrace{\Gamma'_R, \hat{\alpha}}_{\Gamma_R} \longrightarrow \underbrace{\Delta', \hat{\alpha} = \tau'}_{\Delta}} \longrightarrow \text{Solve}$$

By i.h., $\Delta' = (\Delta_L, \alpha, \Delta'_R)$ where $\Gamma_L \longrightarrow \Delta_L$.

Therefore $\Delta = (\Delta_L, \alpha, \Delta'_R, \hat{\alpha} = \tau)$.

(If Γ_R is soft, by i.h. Δ'_R is soft, so $\Delta_R = (\Delta'_R, \hat{\alpha} = \tau)$ is soft.)

$$\bullet \text{ Case } \frac{\Gamma_L, \alpha, \Gamma_R \longrightarrow \Delta'}{\Gamma_L, \alpha, \Gamma_R \longrightarrow \underbrace{\Delta', \hat{\alpha}}_{\Delta}} \longrightarrow \text{Add}$$

By i.h., $\Delta' = (\Delta_L, \alpha, \Delta'_R)$ where $\Gamma_L \longrightarrow \Delta_L$.

Therefore $\Delta = (\Delta_L, \alpha, \Delta'_R, \hat{\alpha})$.

(If Γ_R is soft, by i.h. Δ'_R is soft, so $\Delta_R = (\Delta'_R, \hat{\alpha})$ is soft.)

$$\bullet \text{ Case } \frac{\Gamma_L, \alpha, \Gamma_R \longrightarrow \Delta'}{\Gamma_L, \alpha, \Gamma_R \longrightarrow \Delta', \hat{\alpha} = \tau} \longrightarrow \text{AddSolved}$$

In this case, we know that $\Delta = (\Delta', \hat{\alpha} = \tau)$.
 By i.h., $\Delta' = (\Delta_L, \alpha, \Delta'_R)$ where $\Gamma_L \longrightarrow \Delta_L$.
 Hence $\Delta = (\Delta_L, \alpha, \Delta'_R, \hat{\alpha} = \tau)$.
 (If Γ_R is soft, by i.h. Δ'_R is soft, so $\Delta_R = (\Delta'_R, \hat{\alpha} = \tau)$ is soft.)

- (ii) Similar to the proof of (i), except that the \longrightarrow Marker and \longrightarrow Uvar cases are swapped.
- (iii) Similar to (i), with $\Theta = \hat{\alpha}$ in the \longrightarrow Unsolved case and $\Theta = (\hat{\alpha} = \tau)$ in the \longrightarrow Solve case.
- (iv) Similar to (iii).
- (v) Similar to (i), but using the equality premise of \longrightarrow Var. □

Lemma 25 (Extension Weakening). *If $\Gamma \vdash A$ and $\Gamma \longrightarrow \Delta$ then $\Delta \vdash A$.*

Proof. By a straightforward induction on $\Gamma \vdash A$.

In the UvarWF case, we use Lemma 24 (Extension Order) (i). In the EvarWF case, use Lemma 24 (Extension Order) (iii). In the SolvedEvarWF case, use Lemma 24 (Extension Order) (iv).

In the other cases, apply the i.h. to all subderivations, then apply the rule. □

Lemma 26 (Solution Admissibility for Extension). *If $\Gamma_L \vdash \tau$ then $\Gamma_L, \hat{\alpha}, \Gamma_R \longrightarrow \Gamma_L, \hat{\alpha} = \tau, \Gamma_R$.*

Proof. By induction on Γ_R .

- Case $\Gamma_R = \cdot$:
 By Lemma 20 (Reflexivity) (reflexivity), $\Gamma_L \longrightarrow \Gamma_L$.
 Applying rule \longrightarrow Solve gives $\Gamma_L, \hat{\alpha} \longrightarrow \Gamma_L, \hat{\alpha} = \tau$.
- Case $\Gamma_R = (\Gamma'_R, x : A)$:
 By i.h., $\Gamma_L, \hat{\alpha}, \Gamma'_R \longrightarrow \Gamma_L, \hat{\alpha} = \tau, \Gamma'_R$.
 Applying rule \longrightarrow Var gives $\Gamma_L, \hat{\alpha}, \Gamma'_R, x : A \longrightarrow \Gamma_L, \hat{\alpha} = \tau, \Gamma'_R, x : A$.
- Case $\Gamma_R = (\Gamma'_R, \alpha)$: By i.h. and rule \longrightarrow Uvar.
- Case $\Gamma_R = (\Gamma'_R, \hat{\beta})$: By i.h. and rule \longrightarrow Add.
- Case $\Gamma_R = (\Gamma'_R, \hat{\beta} = \tau')$: By i.h. and rule \longrightarrow AddSolved.
- Case $\Gamma_R = (\Gamma'_R, \blacktriangleright \hat{\beta})$: By i.h. and rule \longrightarrow Marker. □

Lemma 27 (Solved Variable Addition for Extension). *If $\Gamma_L \vdash \tau$ then $\Gamma_L, \Gamma_R \longrightarrow \Gamma_L, \hat{\alpha} = \tau, \Gamma_R$.*

Proof. By induction on Γ_R . The proof is exactly the same as the proof of Lemma 26 (Solution Admissibility for Extension), except that in the $\Gamma_R = \cdot$, we apply rule \longrightarrow AddSolved instead of \longrightarrow Solve. □

Lemma 28 (Unsolved Variable Addition for Extension). *We have that $\Gamma_L, \Gamma_R \longrightarrow \Gamma_L, \hat{\alpha}, \Gamma_R$.*

Proof. By induction on Γ_R . The proof is exactly the same as the proof of Lemma 26 (Solution Admissibility for Extension), except that in the $\Gamma_R = \cdot$ case, we apply rule \longrightarrow Add instead of \longrightarrow Solve. □

Lemma 29 (Parallel Admissibility).

If $\Gamma_L \longrightarrow \Delta_L$ and $\Gamma_L, \Gamma_R \longrightarrow \Delta_L, \Delta_R$ then:

- (i) $\Gamma_L, \hat{\alpha}, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha}, \Delta_R$
- (ii) *If $\Delta_L \vdash \tau'$ then $\Gamma_L, \hat{\alpha}, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha} = \tau', \Delta_R$.*
- (iii) *If $\Gamma_L \vdash \tau$ and $\Delta_L \vdash \tau'$ and $[\Delta_L]\tau = [\Delta_L]\tau'$, then $\Gamma_L, \hat{\alpha} = \tau, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha} = \tau', \Delta_R$.*

Proof. By induction on Δ_R . As always, we assume that all contexts mentioned in the statement of the lemma are well-formed. Hence, $\hat{\alpha} \notin \text{dom}(\Gamma_L) \cup \text{dom}(\Gamma_R) \cup \text{dom}(\Delta_L) \cup \text{dom}(\Delta_R)$.

- (i) We proceed by cases of Δ_R . Observe that in all the extension rules, the right-hand context gets smaller, so as we enter subderivations of $\Gamma_L, \Gamma_R \longrightarrow \Delta_L, \Delta_R$, the context Δ_R becomes smaller.

The only tricky part of the proof is that to apply the i.h., we need $\Gamma_L \longrightarrow \Delta_L$. So we need to make sure that as we drop items from the right of Γ_R and Δ_R , we don't go too far and start decomposing Γ_L or Δ_L ! It's easy to avoid decomposing Δ_L : when $\Delta_R = \cdot$, we don't need to apply the i.h. anyway. To avoid decomposing Γ_L , we need to reason by contradiction, using Lemma 15 (Declaration Preservation).

- **Case $\Delta_R = \cdot$:**
We have $\Gamma_L \longrightarrow \Delta_L$. Applying \longrightarrow Unsolved to that derivation gives the result.
- **Case $\Delta_R = (\Delta'_R, \hat{\beta})$:** We have $\hat{\beta} \neq \hat{\alpha}$ by the well-formedness assumption.
The concluding rule of $\Gamma_L, \Gamma_R \longrightarrow \Delta_L, \Delta'_R, \hat{\beta}$ must have been \longrightarrow Unsolved or \longrightarrow Add. In both cases, the result follows by i.h. and applying \longrightarrow Unsolved or \longrightarrow Add.
Note: In \longrightarrow Add, the left-hand context doesn't change, so we clearly maintain $\Gamma_L \longrightarrow \Delta_L$. In \longrightarrow Unsolved, we can correctly apply the i.h. because $\Gamma_R \neq \cdot$. Suppose, for a contradiction, that $\Gamma_R = \cdot$. Then $\Gamma_L = (\Gamma'_L, \hat{\beta})$. It was given that $\Gamma_L \longrightarrow \Delta_L$, that is, $\Gamma'_L, \hat{\beta} \longrightarrow \Delta_L$. By Lemma 15 (Declaration Preservation), Δ_L has a declaration of $\hat{\beta}$. But then $\Delta = (\Delta_L, \Delta'_R, \hat{\beta})$ is not well-formed: contradiction. Therefore $\Gamma_R \neq \cdot$.
- **Case $\Delta_R = (\Delta'_R, \hat{\beta} = \tau)$:** We have $\hat{\beta} \neq \hat{\alpha}$ by the well-formedness assumption.
The concluding rule must have been \longrightarrow Solved, \longrightarrow Solve or \longrightarrow AddSolved. In each case, apply the i.h. and then the corresponding rule. (In \longrightarrow Solved and \longrightarrow Solve, use Lemma 15 (Declaration Preservation) to show $\Gamma_R \neq \cdot$.)
- **Case $\Delta_R = (\Delta'_R, \alpha)$:** The concluding rule must have been \longrightarrow Uvar. The result follows by i.h. and applying \longrightarrow Uvar.
- **Case $\Delta_R = (\Delta'_R, \blacktriangleright \hat{\beta})$:** Similar to the previous case, with rule \longrightarrow Marker.
- **Case $\Delta_R = (\Delta'_R, x : A)$:** Similar to the previous case, with rule \longrightarrow Var.

(ii) Similar to part (i), except that when $\Delta_R = \cdot$, apply rule \longrightarrow Solve.

(iii) Similar to part (i), except that when $\Delta_R = \cdot$, apply rule \longrightarrow Solved, using the given equality to satisfy the second premise. \square

Lemma 30 (Parallel Extension Solution).

If $\Gamma_L, \hat{\alpha}, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha} = \tau', \Delta_R$ and $\Gamma_L \vdash \tau$ and $[\Delta_L]\tau = [\Delta_L]\tau'$ then $\Gamma_L, \hat{\alpha} = \tau, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha} = \tau', \Delta_R$.

Proof. By induction on Δ_R .

In the case where $\Delta_R = (\Delta'_R, \hat{\alpha} = \tau')$, we know that rule \longrightarrow Solve must have concluded the derivation (we can use Lemma 15 (Declaration Preservation) to get a contradiction that rules out \longrightarrow AddSolved); then we have a subderivation $\Gamma_L \longrightarrow \Delta_L$, to which we can apply \longrightarrow Solved. \square

Lemma 31 (Parallel Variable Update).

If $\Gamma_L, \hat{\alpha}, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha} = \tau_0, \Delta_R$ and $\Gamma_L \vdash \tau_1$ and $\Delta_L \vdash \tau_2$ and $[\Delta_L]\tau_0 = [\Delta_L]\tau_1 = [\Delta_L]\tau_2$ then $\Gamma_L, \hat{\alpha} = \tau_1, \Gamma_R \longrightarrow \Delta_L, \hat{\alpha} = \tau_2, \Delta_R$.

Proof. By induction on Δ_R . Similar to the proof of Lemma 30 (Parallel Extension Solution), but applying \longrightarrow Solved at the end. \square

D'.2 Instantiation Extends

Lemma 32 (Instantiation Extension).

If $\Gamma \vdash \hat{\alpha} \preceq \tau \dashv \Delta$ or $\Gamma \vdash \tau \preceq \hat{\alpha} \dashv \Delta$ then $\Gamma \longrightarrow \Delta$.

Proof. By induction on the given instantiation derivation.

- **Case**
$$\frac{\Gamma \vdash \tau}{\Gamma, \hat{\alpha}, \Gamma' \vdash \hat{\alpha} : \leq \tau \dashv \Gamma, \hat{\alpha} = \tau, \Gamma'} \text{InstLSolve}$$

By Lemma 26 (Solution Admissibility for Extension), $\Gamma, \hat{\alpha}, \Gamma' \longrightarrow \Gamma, \hat{\alpha} = \tau, \Gamma'$.

- **Case**
$$\frac{}{\Gamma[\hat{\alpha}][\hat{\beta}] \vdash \hat{\alpha} : \leq \hat{\beta} \dashv \Gamma[\hat{\alpha}][\hat{\beta} = \hat{\alpha}]} \text{InstLReach}$$

$\Gamma[\hat{\alpha}][\hat{\beta}] = \Gamma_0, \hat{\alpha}, \Gamma_1, \hat{\beta}, \Gamma_2$ for some $\Gamma_0, \Gamma_1, \Gamma_2$.

By the definition of well-formedness, $\Gamma_0, \hat{\alpha}, \Gamma_1 \vdash \hat{\alpha}$.

Therefore, by Lemma 26 (Solution Admissibility for Extension), $\Gamma_0, \hat{\alpha}, \Gamma_1, \hat{\beta}, \Gamma_2 \longrightarrow \Gamma_0, \hat{\alpha}, \Gamma_1, \hat{\beta} = \hat{\alpha}, \Gamma_2$.

- **Case**
$$\frac{\Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2] \vdash A_1 \leq \hat{\alpha}_1 \dashv \Gamma' \quad \Gamma' \vdash \hat{\alpha}_2 : \leq [\Gamma']A_2 \dashv \Delta}{\Gamma[\hat{\alpha}] \vdash \hat{\alpha} : \leq A_1 \rightarrow A_2 \dashv \Delta} \text{InstLArr}$$

By Lemma 28 (Unsolved Variable Addition for Extension), we can insert an (unsolved) $\hat{\alpha}_2$, giving $\Gamma[\hat{\alpha}] \longrightarrow \Gamma[\hat{\alpha}_2, \hat{\alpha}]$.

By Lemma 28 (Unsolved Variable Addition for Extension) again, $\Gamma[\hat{\alpha}_2, \hat{\alpha}] \longrightarrow \Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha}]$.

By Lemma 26 (Solution Admissibility for Extension), we can solve $\hat{\alpha}$, giving $\Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha}] \longrightarrow \Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2]$.

Then by transitivity (Lemma 21 (Transitivity)), $\Gamma[\hat{\alpha}] \longrightarrow \Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2]$.

By i.h. on the first subderivation, $\Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2] \longrightarrow \Gamma'$.

By i.h. on the second subderivation, $\Gamma' \longrightarrow \Delta$.

By transitivity (Lemma 21 (Transitivity)), $\Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2] \longrightarrow \Delta$.

By transitivity (Lemma 21 (Transitivity)), $\Gamma[\hat{\alpha}] \longrightarrow \Delta$.

- **Case**
$$\frac{\Gamma[\hat{\alpha}], \beta \vdash \hat{\alpha} : \leq B \dashv \Delta, \beta, \Delta'}{\Gamma[\hat{\alpha}] \vdash \hat{\alpha} : \leq \forall \beta. B \dashv \Delta} \text{InstLAllR}$$

By induction, $\Gamma[\hat{\alpha}], \beta \longrightarrow \Delta, \beta, \Delta'$.

By Lemma 24 (Extension Order) (i), we have $\Gamma[\hat{\alpha}] \longrightarrow \Delta$.

- **Case**
$$\frac{\Gamma \vdash \tau}{\Gamma, \hat{\alpha}, \Gamma' \vdash \tau \leq \hat{\alpha} \dashv \Gamma, \hat{\alpha} = \tau, \Gamma'} \text{InstRSolve}$$

By Lemma 26 (Solution Admissibility for Extension), we can solve $\hat{\alpha}$, giving $\Gamma, \hat{\alpha}, \Gamma' \longrightarrow \Gamma, \hat{\alpha} = \tau, \Gamma'$.

- **Case**
$$\frac{}{\Gamma[\hat{\alpha}][\hat{\beta}] \vdash \hat{\beta} \leq \hat{\alpha} \dashv \Gamma[\hat{\alpha}][\hat{\beta} = \hat{\alpha}]} \text{InstRReach}$$

$\Gamma[\hat{\alpha}][\hat{\beta}] = \Gamma_0, \hat{\alpha}, \Gamma_1, \hat{\beta}, \Gamma_2$ for some $\Gamma_0, \Gamma_1, \Gamma_2$.

By the definition of well-formedness, $\Gamma_0, \hat{\alpha}, \Gamma_1 \vdash \hat{\alpha}$.

Hence by Lemma 26 (Solution Admissibility for Extension), we can solve $\hat{\beta}$, giving $\Gamma_0, \hat{\alpha}, \Gamma_1, \hat{\beta}, \Gamma_2 \longrightarrow \Gamma_0, \hat{\alpha}, \Gamma_1, \hat{\beta} = \hat{\alpha}, \Gamma_2$.

- **Case**
$$\frac{\Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2] \vdash \hat{\alpha}_1 : \leq A_1 \dashv \Gamma' \quad \Gamma' \vdash [\Gamma']A_2 \leq \hat{\alpha}_2 \dashv \Delta}{\Gamma[\hat{\alpha}] \vdash A_1 \rightarrow A_2 \leq \hat{\alpha} \dashv \Delta} \text{InstRArr}$$

Because the contexts here are the same as in InstLArr, this is the same as the InstLArr case.

- **Case**
$$\frac{\Gamma[\hat{\alpha}], \blacktriangleright_{\hat{\beta}}, \hat{\beta} \vdash [\hat{\beta}/\beta]B \leq \hat{\alpha} \dashv \Delta, \blacktriangleright_{\hat{\beta}}, \Delta'}{\Gamma[\hat{\alpha}] \vdash \forall \beta. B \leq \hat{\alpha} \dashv \Delta} \text{InstRAIll}$$

By i.h., $\Gamma[\hat{\alpha}], \blacktriangleright_{\hat{\beta}}, \hat{\beta} \longrightarrow \Delta, \blacktriangleright_{\hat{\beta}}, \Delta'$.

By Lemma 24 (Extension Order) (ii), $\Gamma[\hat{\alpha}] \longrightarrow \Delta$. □

D'.3 Subtyping Extends

Lemma 33 (Subtyping Extension).

If $\Gamma \vdash A <: B \dashv \Delta$ then $\Gamma \longrightarrow \Delta$.

Proof. By induction on the given derivation.

For cases $<:\text{Var}$, $<:\text{Unit}$, $<:\text{Exvar}$, we have $\Delta = \Gamma$, so Lemma 20 (Reflexivity) suffices.

- **Case**
$$\frac{\Gamma \vdash B_1 <: A_1 \dashv \Theta \quad \Theta \vdash [\Omega]A_2 <: [\Omega]B_2 \dashv \Delta}{\Gamma \vdash A_1 \rightarrow A_2 <: B_1 \rightarrow B_2 \dashv \Delta} <:\rightarrow$$

By IH on each subderivation, $\Gamma \longrightarrow \Theta$ and $\Theta \longrightarrow \Delta$.

By Lemma 21 (Transitivity) (transitivity), $\Gamma \longrightarrow \Delta$, which was to be shown.

- **Case**
$$\frac{\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha} \vdash [\hat{\alpha}/\alpha]A <: B \dashv \Delta, \blacktriangleright_{\hat{\alpha}}, \Theta}{\Gamma \vdash \forall \alpha. A <: B \dashv \Delta} <:\forall L$$

By IH, $\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha} \longrightarrow \Delta, \blacktriangleright_{\hat{\alpha}}, \Theta$.

By Lemma 24 (Extension Order) (ii) with $\Gamma_L = \Gamma$ and $\Gamma'_L = \Delta$ and $\Gamma_R = \hat{\alpha}$ and $\Gamma'_R = \Theta$, we obtain

$$\Gamma \longrightarrow \Delta$$

- **Case**
$$\frac{\Gamma, \beta \vdash A <: B \dashv \Delta, \beta, \Theta}{\Gamma \vdash A <: \forall \beta. B \dashv \Delta} <:\forall R$$

By IH, we have $\Gamma, \beta \longrightarrow \Delta, \beta, \Theta$.

By Lemma 24 (Extension Order) (i), we obtain $\Gamma \longrightarrow \Delta$, which was to be shown.

- **Cases $<:\text{InstantiateL}$, $<:\text{InstantiateR}$:** In each of these rules, the premise has the same input and output contexts as the conclusion, so Lemma 32 (Instantiation Extension) suffices. \square

E' Decidability of Instantiation

Lemma 34 (Left Unsolvedness Preservation).

If $\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_{\Gamma} \vdash \hat{\alpha} : \preceq A \dashv \Delta$ or $\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_{\Gamma} \vdash A \preceq \hat{\alpha} \dashv \Delta$, and $\hat{\beta} \in \text{unsolved}(\Gamma_0)$, then $\hat{\beta} \in \text{unsolved}(\Delta)$.

Proof. By induction on the given derivation.

- **Case**
$$\frac{\Gamma_0 \vdash \tau}{\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_{\Gamma} \vdash \hat{\alpha} : \preceq \tau \dashv \Gamma_0, \hat{\alpha} = \tau, \Gamma_1} \text{InstLSolve}$$

Immediate, since to the left of $\hat{\alpha}$, the contexts Δ and Γ are the same.

- **Case**
$$\frac{}{\Gamma[\hat{\alpha}][\hat{\beta}] \vdash \hat{\alpha} : \preceq \hat{\beta} \dashv \Gamma[\hat{\alpha}][\hat{\beta} = \hat{\alpha}]} \text{InstLReach}$$

Immediate, since to the left of $\hat{\alpha}$, the contexts Δ and Γ are the same.

- **Case**
$$\frac{\Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2] \vdash A_1 \preceq \hat{\alpha}_1 \dashv \Gamma' \quad \Gamma' \vdash \hat{\alpha}_2 : \preceq [\Gamma']A_2 \dashv \Delta}{\Gamma[\hat{\alpha}] \vdash \hat{\alpha} : \preceq A_1 \rightarrow A_2 \dashv \Delta} \text{InstLArr}$$

We have $\hat{\beta} \in \text{unsolved}(\Gamma_0)$. Therefore $\hat{\beta} \in \text{unsolved}(\Gamma_0, \hat{\alpha}_2)$.

Clearly, $\hat{\alpha}_2 \in \text{unsolved}(\Gamma_0, \hat{\alpha}_2)$.

We have two subderivations:

$$\Gamma_0, \hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2, \Gamma_1 \vdash A_1 \preceq \hat{\alpha}_1 \dashv \Gamma' \quad (1)$$

$$\Gamma' \vdash \hat{\alpha}_2 \preceq [\Gamma']A_2 \dashv \Delta \quad (2)$$

By induction on (1), $\hat{\beta} \in \text{unsolved}(\Gamma')$.

Also by induction on (1), with $\hat{\alpha}_2$ playing the role of $\hat{\beta}$, we get $\hat{\alpha}_2 \in \text{unsolved}(\Gamma')$.

Since $\hat{\beta} \in \Gamma_0$, it is declared to the left of $\hat{\alpha}_2$ in $\Gamma_0, \hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2, \Gamma_1$.

Hence by Lemma 16 (Declaration Order Preservation), $\hat{\beta}$ is declared to the left of $\hat{\alpha}_2$ in Γ' . That is, $\Gamma' = (\Gamma'_0, \hat{\alpha}_2, \Gamma'_1)$, where $\hat{\beta} \in \text{unsolved}(\Gamma'_0)$.

By induction on (2), $\hat{\beta} \in \text{unsolved}(\Delta)$.

• **Case**
$$\frac{\Gamma_0, \hat{\alpha}, \Gamma_1, \beta \vdash \hat{\alpha} \preceq B \dashv \Delta, \beta, \Delta'}{\Gamma_0, \hat{\alpha}, \Gamma_1 \vdash \hat{\alpha} \preceq \forall \beta. B \dashv \Delta} \text{InstLAllR}$$

We have $\hat{\beta} \in \text{unsolved}(\Gamma_0)$.

By induction, $\hat{\beta} \in \text{unsolved}(\Delta, \beta, \Delta')$.

Note that $\hat{\beta}$ is declared to the left of β in $\Gamma_0, \hat{\alpha}, \Gamma_1, \beta$.

By Lemma 16 (Declaration Order Preservation), $\hat{\beta}$ is declared to the left of β in (Δ, β, Δ') , that is, in Δ . Since $\hat{\beta} \in \text{unsolved}(\Delta, \beta, \Delta')$, we have $\hat{\beta} \in \text{unsolved}(\Delta)$.

• **Cases** InstRSolve, InstRReach: Similar to the InstLSolve and InstLReach cases.

• **Case**
$$\frac{\Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2] \vdash \hat{\alpha}_1 \preceq A_1 \dashv \Gamma' \quad \Gamma' \vdash [\Gamma']A_2 \preceq \hat{\alpha}_2 \dashv \Delta}{\Gamma[\hat{\alpha}] \vdash A_1 \rightarrow A_2 \preceq \hat{\alpha} \dashv \Delta} \text{InstRArr}$$

Similar to the InstLArr case.

• **Case**
$$\frac{\Gamma[\hat{\alpha}], \blacktriangleright_{\hat{\gamma}}, \hat{\gamma} \vdash [\hat{\gamma}/\beta]B \preceq \hat{\alpha} \dashv \Delta, \blacktriangleright_{\hat{\gamma}}, \Delta'}{\Gamma[\hat{\alpha}] \vdash \forall \beta. B \preceq \hat{\alpha} \dashv \Delta} \text{InstRAIll}$$

We have $\hat{\beta} \in \text{unsolved}(\Gamma_0)$.

By induction, $\hat{\beta} \in \text{unsolved}(\Delta, \blacktriangleright_{\hat{\gamma}}, \Delta')$.

Note that $\hat{\beta}$ is declared to the left of $\blacktriangleright_{\hat{\gamma}}$ in $\Gamma_0, \hat{\alpha}, \Gamma_1, \blacktriangleright_{\hat{\gamma}}, \hat{\gamma}$.

By Lemma 16 (Declaration Order Preservation), $\hat{\beta}$ is declared to the left of $\blacktriangleright_{\hat{\gamma}}$ in $\Delta, \blacktriangleright_{\hat{\gamma}}, \Delta'$.

Hence $\hat{\beta}$ is declared in Δ , and we know it is in $\text{unsolved}(\Delta, \blacktriangleright_{\hat{\gamma}}, \Delta')$, so $\hat{\beta} \in \text{unsolved}(\Delta)$. \square

Lemma 35 (Left Free Variable Preservation). *If $\overbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}^{\Gamma} \vdash \hat{\alpha} \preceq A \dashv \Delta$ or $\overbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}^{\Gamma} \vdash A \preceq \hat{\alpha} \dashv \Delta$, and $\Gamma \vdash B$ and $\hat{\alpha} \notin \text{FV}([\Gamma]B)$ and $\hat{\beta} \in \text{unsolved}(\Gamma_0)$ and $\hat{\beta} \notin \text{FV}([\Gamma]B)$, then $\hat{\beta} \notin \text{FV}([\Delta]B)$.*

Proof. By induction on the given instantiation derivation.

• **Case**
$$\frac{\Gamma_0 \vdash \tau}{\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_{\Gamma} \vdash \hat{\alpha} \preceq \tau \dashv \underbrace{\Gamma_0, \hat{\alpha} = \tau, \Gamma_1}_{\Delta}} \text{InstLSolve}$$

We have $\hat{\alpha} \notin \text{FV}([\Gamma]B)$. Since Δ differs from Γ only in $\hat{\alpha}$, it must be the case that $[\Gamma]B = [\Delta]B$. It is given that $\hat{\beta} \notin \text{FV}([\Gamma]B)$, so $\hat{\beta} \notin \text{FV}([\Delta]B)$.

• **Case**
$$\frac{\Gamma'[\hat{\alpha}][\hat{\gamma}] \vdash \hat{\alpha} \preceq \hat{\gamma} \dashv \Gamma'[\hat{\alpha}][\hat{\gamma} = \hat{\alpha}]}{\Gamma[\hat{\alpha}] \vdash \hat{\alpha} \preceq \hat{\gamma} \dashv \Delta} \text{InstLReach}$$

Since Δ differs from Γ only in solving $\hat{\gamma}$ to $\hat{\alpha}$, applying Δ to a type will not introduce a $\hat{\beta}$. We have $\hat{\beta} \notin \text{FV}([\Gamma]B)$, so $\hat{\beta} \notin \text{FV}([\Delta]B)$.

- **Case**
$$\frac{\Gamma_0 \vdash \tau}{\Gamma_0, \hat{\alpha}, \Gamma_1 \vdash \tau \preceq: \hat{\alpha} \dashv \Gamma_0, \hat{\alpha} = \tau, \Gamma_1} \text{InstRSolve}$$

Similar to the InstLSolve case.

- **Case**
$$\frac{}{\Gamma'[\hat{\alpha}][\hat{\gamma}] \vdash \hat{\gamma} \preceq: \hat{\alpha} \dashv \Gamma'[\hat{\alpha}][\hat{\gamma} = \hat{\alpha}]} \text{InstRReach}$$

Similar to the InstLReach case.

- **Case**
$$\frac{\overbrace{\Gamma_0, \hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2, \Gamma_1 \vdash A_1 \preceq: \hat{\alpha}_1 \dashv \Delta \quad \Delta \vdash \hat{\alpha}_2 \preceq: [\Delta]A_2 \dashv \Delta}^{\Gamma'} \quad \underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1 \vdash \hat{\alpha} \preceq: A_1 \rightarrow A_2 \dashv \Delta}_{\Gamma}}{\Gamma_0, \hat{\alpha}, \Gamma_1 \vdash \hat{\alpha} \preceq: A_1 \rightarrow A_2 \dashv \Delta} \text{InstLArr}$$

We have $\Gamma \vdash B$ and $\hat{\alpha} \notin \text{FV}([\Gamma]B)$ and $\hat{\beta} \notin \text{FV}([\Gamma]B)$.

By weakening, we get $\Gamma' \vdash B$; since $\hat{\alpha} \notin \text{FV}([\Gamma]B)$ and Γ' only adds a solution for $\hat{\alpha}$, it follows that $[\Gamma']B = [\Gamma]B$.

Therefore $\hat{\alpha}_1 \notin \text{FV}([\Gamma']B)$ and $\hat{\alpha}_2 \notin \text{FV}([\Gamma']B)$ and $\hat{\beta} \notin \text{FV}([\Gamma']B)$.

Since we have $\hat{\beta} \in \Gamma_0$, we also have $\hat{\beta} \in (\Gamma_0, \hat{\alpha}_2)$.

By induction on the first premise, $\hat{\beta} \notin \text{FV}([\Delta]B)$.

Also by induction on the first premise, with $\hat{\alpha}_2$ playing the role of $\hat{\beta}$, we have $\hat{\alpha}_2 \notin \text{FV}([\Delta]B)$.

Note that $\hat{\alpha}_2 \in \text{unsolved}(\Gamma_0, \hat{\alpha}_2)$.

By Lemma 34 (Left Unsolvedness Preservation), $\hat{\alpha}_2 \in \text{unsolved}(\Delta)$.

Therefore Δ has the form $(\Delta_0, \hat{\alpha}_2, \Delta_1)$.

Since $\hat{\beta} \neq \hat{\alpha}_2$, we know that $\hat{\beta}$ is declared to the left of $\hat{\alpha}_2$ in $\Gamma_0, \hat{\alpha}_2$, so by Lemma 16 (Declaration Order Preservation), $\hat{\beta}$ is declared to the left of $\hat{\alpha}_2$ in Δ . Hence $\hat{\beta} \in \Delta_0$.

Furthermore, by Lemma 32 (Instantiation Extension), we have $\Gamma' \rightarrow \Delta$.

Then by Lemma 25 (Extension Weakening), we have $\Delta \vdash B$. Using induction on the second premise, $\hat{\beta} \notin \text{FV}([\Delta]B)$.

- **Case**
$$\frac{\Gamma_0, \hat{\alpha}, \Gamma_1, \gamma \vdash \hat{\alpha} \preceq: C \dashv \Delta, \gamma, \Delta'}{\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1 \vdash \hat{\alpha} \preceq: \forall \gamma. C \dashv \Delta}_{\Gamma}} \text{InstLAllR}$$

We have $\Gamma \vdash B$ and $\hat{\alpha} \notin \text{FV}([\Gamma]B)$ and $\hat{\beta} \in \Gamma_0$ and $\hat{\beta} \notin \text{FV}([\Gamma]B)$.

By weakening, $\Gamma, \gamma \vdash B$; by the definition of substitution, $[\Gamma, \gamma]B = [\Gamma]B$.

Substituting equals for equals, $\hat{\alpha} \notin \text{FV}([\Gamma, \gamma]B)$ and $\hat{\beta} \notin \text{FV}([\Gamma, \gamma]B)$.

By induction, $\hat{\beta} \notin \text{FV}([\Delta, \gamma, \Delta']B)$.

Since $\hat{\beta}$ is declared to the left of γ in (Γ, γ) , we can use Lemma 16 (Declaration Order Preservation) to show that $\hat{\beta}$ is declared to the left of γ in $(\Delta, \gamma, \Delta')$, that is, in Δ .

We have $\Gamma \vdash B$, so $\gamma \notin \text{FV}(B)$. Thus each free variable u in B is in Γ , to the left of γ in (Γ, γ) .

Therefore, by Lemma 16 (Declaration Order Preservation), each free variable u in B is in Δ .

Therefore $[\Delta, \gamma, \Delta']B = [\Delta]B$.

Earlier, we obtained $\hat{\beta} \notin \text{FV}([\Delta, \gamma, \Delta']B)$, so substituting equals for equals, $\hat{\beta} \notin \text{FV}([\Delta]B)$.

- **Case**
$$\frac{\Gamma_0, \hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2, \Gamma_1 \vdash \hat{\alpha}_1 \preceq: A_1 \dashv \Delta \quad \Gamma' \vdash [\Delta]A_2 \preceq: \hat{\alpha}_2 \dashv \Delta}{\Gamma_0, \hat{\alpha}, \Gamma_1 \vdash A_1 \rightarrow A_2 \preceq: \hat{\alpha} \dashv \Delta} \text{InstRArr}$$

Similar to the InstLArr case.

- **Case**
$$\frac{\Gamma[\hat{\alpha}], \blacktriangleright_{\hat{\gamma}}, \hat{\gamma} \vdash [\hat{\gamma}/\gamma]C \preceq: \hat{\alpha} \dashv \Delta, \blacktriangleright_{\hat{\gamma}}, \Delta'}{\Gamma[\hat{\alpha}] \vdash \forall \gamma. C \preceq: \hat{\alpha} \dashv \Delta} \text{InstRAIII}$$

We have $\Gamma \vdash B$ and $\hat{\alpha} \notin \text{FV}([\Gamma]B)$ and $\hat{\beta} \in \Gamma_0$ and $\hat{\beta} \notin \text{FV}([\Gamma]B)$.

By weakening, $\Gamma, \blacktriangleright_{\hat{\gamma}}, \hat{\gamma} \vdash B$; by the definition of substitution, $[\Gamma, \blacktriangleright_{\hat{\gamma}}, \hat{\gamma}]B = [\Gamma]B$.

Substituting equals for equals, $\hat{\alpha} \notin \text{FV}([\Gamma, \blacktriangleright_{\hat{\gamma}}, \hat{\gamma}]B)$ and $\hat{\beta} \notin \text{FV}([\Gamma, \blacktriangleright_{\hat{\gamma}}, \hat{\gamma}]B)$.

By induction, $\hat{\beta} \notin \text{FV}([\Delta, \blacktriangleright_{\hat{\gamma}}, \Delta']B)$.

Note that $\hat{\beta}$ is declared to the left of $\blacktriangleright_{\hat{\gamma}}$ in $\Gamma, \blacktriangleright_{\hat{\gamma}}, \hat{\gamma}$.

By Lemma 16 (Declaration Order Preservation), $\hat{\beta}$ is declared to the left of $\blacktriangleright_{\hat{\gamma}}$ in $\Delta, \blacktriangleright_{\hat{\gamma}}, \Delta'$.

So $\hat{\beta}$ is declared in Δ .

Now, note that each free variable u in B is in Γ , which is to the left of $\blacktriangleright_{\hat{\gamma}}$ in $\Gamma, \blacktriangleright_{\hat{\gamma}}, \hat{\gamma}$.

Therefore, by Lemma 16 (Declaration Order Preservation), each free variable u in B is in Δ .

Therefore $[\Delta, \blacktriangleright_{\hat{\gamma}}, \Delta']B = [\Delta]B$.

Earlier, we obtained $\hat{\beta} \notin \text{FV}([\Delta, \blacktriangleright_{\hat{\gamma}}, \Delta']B)$, so substituting equals for equals, $\hat{\beta} \notin \text{FV}([\Delta]B)$. \square

Lemma 36 (Instantiation Size Preservation). *If $\Gamma_0, \hat{\alpha}, \Gamma_1 \vdash \hat{\alpha} : \leq A \dashv \Delta$ or $\Gamma_0, \hat{\alpha}, \Gamma_1 \vdash A \leq \hat{\alpha} \dashv \Delta$, and $\Gamma \vdash B$ and $\hat{\alpha} \notin \text{FV}([\Gamma]B)$, then $||\Gamma]B| = ||[\Delta]B|$, where $|C|$ is the plain size of the term C .*

Proof. By induction on the given derivation.

- **Case**
$$\frac{\Gamma_0 \vdash \tau}{\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_{\Gamma} \vdash \hat{\alpha} : \leq \tau \dashv \Gamma_0, \hat{\alpha} = \tau, \Gamma_1} \text{InstLSolve}$$

Since Δ differs from Γ only in solving $\hat{\alpha}$, and we know $\hat{\alpha} \notin \text{FV}([\Gamma]B)$, we have $[\Delta]B = [\Gamma]B$; therefore $||[\Delta]B| = ||[\Gamma]B|$.

- **Case**
$$\frac{}{\Gamma[\hat{\alpha}][\hat{\beta}] \vdash \hat{\alpha} : \leq \hat{\beta} \dashv \Gamma[\hat{\alpha}][\hat{\beta} = \hat{\alpha}]} \text{InstLReach}$$

Here, Δ differs from Γ only in solving $\hat{\beta}$ to $\hat{\alpha}$. However, $\hat{\alpha}$ has the same size as $\hat{\beta}$, so even if $\hat{\beta} \in \text{FV}([\Gamma]B)$, we have $||[\Delta]B| = ||[\Gamma]B|$.

- **Case**
$$\frac{\overbrace{\Gamma_0, \hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2, \Gamma_1}_{\Gamma'} \vdash A_1 \leq \hat{\alpha}_1 \dashv \Theta \quad \Theta \vdash \hat{\alpha}_2 : \leq [\Theta]A_2 \dashv \Delta}{\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_{\Gamma} \vdash \hat{\alpha} : \leq A_1 \rightarrow A_2 \dashv \Delta} \text{InstLArr}$$

We have $\Gamma \vdash B$ and $\hat{\alpha} \notin \text{FV}([\Gamma]B)$. Since $\hat{\alpha}_1, \hat{\alpha}_2 \notin \text{dom}(\Gamma)$, we have $\hat{\alpha}, \hat{\alpha}_1, \hat{\alpha}_2 \notin \text{FV}([\Gamma]B)$. It follows that $[\Gamma']B = [\Gamma]B$.

By weakening, $\Gamma' \vdash B$.

By induction on the first premise, $||[\Gamma']B| = ||[\Theta]B|$.

By Lemma 16 (Declaration Order Preservation), since $\hat{\alpha}_2$ is declared to the left of $\hat{\alpha}_1$ in Γ' , we have that $\hat{\alpha}_2$ is declared to the left of $\hat{\alpha}_1$ in Θ .

By Lemma 34 (Left Unsolvedness Preservation), since $\hat{\alpha}_2 \in \text{unsolved}(\Gamma')$, it is unsolved in Θ : that is, $\Theta = (\Theta_0, \hat{\alpha}_2, \Theta_1)$.

By Lemma 32 (Instantiation Extension), we have $\Gamma' \longrightarrow \Theta$.

By Lemma 25 (Extension Weakening), $\Theta \vdash B$.

Since $\hat{\alpha}_2 \notin \text{FV}([\Gamma']B)$, Lemma 35 (Left Free Variable Preservation) gives $\hat{\alpha}_2 \notin \text{FV}([\Theta]B)$.

By induction on the second premise, $||[\Theta]B| = ||[\Delta]B|$, and by transitivity of equality, $||[\Gamma]B| = ||[\Delta]B|$.

- **Case**
$$\frac{\Gamma_0, \hat{\alpha}, \Gamma_1, \beta \vdash \hat{\alpha} : \leq A_0 \dashv \Delta, \beta, \Delta'}{\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_{\Gamma} \vdash \hat{\alpha} : \leq \forall \beta. A_0 \dashv \Delta} \text{InstLAllR}$$

We have $\Gamma \vdash B$ and $\hat{\alpha} \notin \text{FV}([\Gamma]B)$.

By weakening, $\Gamma, \beta \vdash B$.

From the definition of substitution, $[\Gamma]B = [\Gamma, \beta]B$. Hence $\hat{\alpha} \notin \text{FV}([\Gamma, \beta]B)$.

The input context of the premise is $(\Gamma_0, \hat{\alpha}, \Gamma_1, \beta)$, which is (Γ, β) , so by induction, $||[\Gamma, \beta]B| = ||[\Delta, \beta, \Delta']B|$.

Suppose u is a free variable in B . Then u is declared in Γ , and so occurs before β in Γ, β .

By Lemma 16 (Declaration Order Preservation), u is declared before β in Δ, β, Δ' .
 So every free variable u in B is declared in Δ .
 Hence $[\Delta, \beta, \Delta']B = [\Delta]B$.
 We have $[\Gamma]B = [\Gamma, \beta]B$, so $||[\Gamma]B|| = ||[\Gamma, \beta]B||$; by transitivity of equality, $||[\Gamma]B|| = ||[\Delta]B||$.

- **Case**
$$\frac{\Gamma_0 \vdash \tau}{\Gamma_0, \hat{\alpha}, \Gamma_1 \vdash \tau \preceq \hat{\alpha} \dashv \Gamma_0, \hat{\alpha} = \tau, \Gamma_1} \text{InstRSolve}$$

Similar to the InstLSolve case.

- **Case**
$$\frac{\Gamma[\hat{\alpha}][\hat{\beta}] \vdash \hat{\beta} \preceq \hat{\alpha} \dashv \Gamma[\hat{\alpha}][\hat{\beta} = \hat{\alpha}]}{\Gamma[\hat{\alpha}][\hat{\beta}] \vdash \hat{\beta} \preceq \hat{\alpha} \dashv \Gamma[\hat{\alpha}][\hat{\beta} = \hat{\alpha}]} \text{InstRReach}$$

Similar to the InstLReach case.

- **Case**
$$\frac{\overbrace{\Gamma_0, \hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2, \Gamma_1 \vdash \hat{\alpha}_1 : \preceq A_1 \dashv \Theta}^{\Gamma'} \quad \Theta \vdash [\Theta]A_2 \preceq \hat{\alpha}_2 \dashv \Delta}{\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1 \vdash A_1 \rightarrow A_2 \preceq \hat{\alpha} \dashv \Delta}_{\Gamma}} \text{InstRArr}}$$

Similar to the InstLArr case.

- **Case**
$$\frac{\Gamma'[\hat{\alpha}], \blacktriangleright_{\hat{\beta}}, \hat{\beta} \vdash [\hat{\beta}/\beta]A_0 \preceq \hat{\alpha} \dashv \Delta, \blacktriangleright_{\hat{\beta}}, \Delta'}{\Gamma'[\hat{\alpha}] \vdash \forall \beta. A_0 \preceq \hat{\alpha} \dashv \Delta} \text{InstRAILL}$$

We have $\Gamma \vdash B$ and $\hat{\alpha} \notin \text{FV}([\Gamma]B)$.

By weakening, $\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta} \vdash B$.

From the definition of substitution, $[\Gamma]B = [\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta}]B$. Hence $\hat{\alpha} \notin \text{FV}([\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta}]B)$.

By induction, $||[\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta}]B|| = ||[\Delta, \blacktriangleright_{\hat{\beta}}, \Delta']B||$.

Suppose u is a free variable in B .

Then u is declared in Γ , and so occurs before $\blacktriangleright_{\hat{\beta}}$ in $\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta}$.

By Lemma 16 (Declaration Order Preservation), u is declared before $\blacktriangleright_{\hat{\beta}}$ in $\Delta, \blacktriangleright_{\hat{\beta}}, \Delta'$.

So every free variable u in B is declared in Δ .

Hence $[\Delta, \blacktriangleright_{\hat{\beta}}, \Delta']B = [\Delta]B$.

Since $[\Gamma]B = [\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta}]B$, we have $||[\Gamma]B|| = ||[\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta}]B||$; by transitivity of equality, $||[\Gamma]B|| = ||[\Delta]B||$. \square

Theorem 7 (Decidability of Instantiation). *If $\Gamma = \Gamma_0[\hat{\alpha}]$ and $\Gamma \vdash A$ such that $[\Gamma]A = A$ and $\hat{\alpha} \notin \text{FV}(A)$, then:*

(1) *Either there exists Δ such that $\Gamma_0[\hat{\alpha}] \vdash \hat{\alpha} : \preceq A \dashv \Delta$, or not.*

(2) *Either there exists Δ such that $\Gamma_0[\hat{\alpha}] \vdash A \preceq \hat{\alpha} \dashv \Delta$, or not.*

Proof. By induction on the derivation of $\Gamma \vdash A$.

(1) $\Gamma \vdash \hat{\alpha} : \preceq A \dashv \Delta$ is decidable.

- **Case**

$$\frac{\Gamma_L, \hat{\alpha}, \Gamma_R \vdash \alpha}{\Gamma'[\alpha]} \text{UvarWF}$$

If $\alpha \in \Gamma_L$, then by UvarWF we have $\Gamma_L \vdash \alpha$, and by rule InstLSolve we have a derivation.

Otherwise no rule matches, and so no derivation exists.

- **Case UnitWF:** By rule InstLSolve.

- **Case**

$$\frac{\Gamma_L, \hat{\alpha}, \Gamma_R \vdash \hat{\beta}}{\Gamma} \text{EvarWF}$$

By inversion, we have $\hat{\beta} \in \Gamma$, and $[\Gamma]\hat{\beta} = \hat{\beta}$. Since $\hat{\alpha} \notin \text{FV}([\Gamma]\hat{\beta}) = \text{FV}(\hat{\beta}) = \{\hat{\beta}\}$, it follows that $\hat{\alpha} \neq \hat{\beta}$: Either $\hat{\beta} \in \Gamma_L$ or $\hat{\beta} \in \Gamma_R$.

If $\hat{\beta} \in \Gamma_L$, then we have a derivation by InstLSolve.

If $\hat{\beta} \in \Gamma_R$, then we have a derivation by InstLReach.

- **Case**

$$\frac{\Gamma'[\hat{\beta} = \tau] \vdash \hat{\beta}}{\Gamma} \text{SolvedEvarWF}$$

It is given that $[\Gamma]\hat{\beta} = \hat{\beta}$, so this case is impossible.

- **Case**

$$\frac{\Gamma \vdash A_1 \quad \Gamma \vdash A_2}{\Gamma_L, \hat{\alpha}, \Gamma_R \vdash A_1 \rightarrow A_2} \text{ArrowWF}$$

By assumption, $[\Gamma](A_1 \rightarrow A_2) = A_1 \rightarrow A_2$ and $\hat{\alpha} \notin \text{FV}([\Gamma](A_1 \rightarrow A_2))$.

If $A_1 \rightarrow A_2$ is a monotype and is well-formed under Γ_L , we can apply InstLSolve.

Otherwise, the only rule with a conclusion matching $A_1 \rightarrow A_2$ is InstLArr.

First, consider whether $\Gamma_L, \hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2, \Gamma_R \vdash A_1 \rightarrow A_2 \preceq \hat{\alpha}_1 \dashv \dashv$ is decidable.

By definition of substitution, $[\Gamma](A_1 \rightarrow A_2) = ([\Gamma]A_1) \rightarrow ([\Gamma]A_2)$. Since $[\Gamma](A_1 \rightarrow A_2) = A_1 \rightarrow A_2$, we have $[\Gamma]A_1 = A_1$ and $[\Gamma]A_2 = A_2$.

By weakening, $\Gamma_L, \hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2, \Gamma_R \vdash A_1 \rightarrow A_2$.

Since $\Gamma \vdash A_1$ and $\Gamma \vdash A_2$, we have $\hat{\alpha}_1, \hat{\alpha}_2 \notin \text{FV}(A_1) \cup \text{FV}(A_2)$.

Since $\hat{\alpha} \notin \text{FV}(A) \supseteq \text{FV}(A_1)$, it follows that $[\Gamma']A_1 = A_1$.

By i.h., either there exists Θ such that $\Gamma_L, \hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2, \Gamma_R \vdash A_1 \preceq \hat{\alpha}_1 \dashv \Theta$, or not.

If not, then no derivation by InstLArr exists.

If so, then we have $\Gamma_L, \hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2, \Gamma_R \vdash \hat{\alpha}_1 \preceq A_1 \dashv \Theta$.

By Lemma 34 (Left Unsolvedness Preservation), we know that $\hat{\alpha}_2 \in \text{unsolved}(\Theta)$.

By Lemma 35 (Left Free Variable Preservation), we know that $\hat{\alpha}_2 \notin \text{FV}([\Theta]A_2)$.

Clearly, $[\Theta]([\Theta]A_2) = [\Theta]A_2$.

Hence by i.h., either there exists Δ such that $\Theta \vdash \hat{\alpha}_2 \preceq [\Theta]A_2 \dashv \Delta$, or not.

If not, then no derivation by InstLArr exists.

If it does, then by rule InstLArr, we have $\Gamma \vdash \hat{\alpha} \preceq A \dashv \Delta$.

- **Case**

$$\frac{\Gamma, \alpha \vdash A_0}{\Gamma \vdash \forall \alpha. A_0} \text{ForallWF}$$

We have $\forall \alpha. A_0 = [\Gamma](\forall \alpha. A_0)$. By definition of substitution, $[\Gamma](\forall \alpha. A_0) = \forall \alpha. [\Gamma]A_0$, so $A_0 = [\Gamma]A_0$.

By definition of substitution, $[\Gamma, \alpha]A_0 = [\Gamma]A_0$.

We have $\hat{\alpha} \notin \text{FV}([\Gamma](\forall \alpha. A_0))$. Therefore $\hat{\alpha} \notin \text{FV}([\Gamma]A_0) = \text{FV}([\Gamma, \alpha]A_0)$.

By i.h., either there exists Θ such that $\Gamma, \alpha \vdash \hat{\alpha} \preceq A_0 \dashv \Theta$, or not.

Suppose $\Gamma, \alpha \vdash \hat{\alpha} \preceq A_0 \dashv \Theta$.

By Lemma 32 (Instantiation Extension), $\Gamma \longrightarrow \Theta$;

by Lemma 24 (Extension Order) (i), $\Theta = \Delta, \alpha, \Delta'$.

Hence by rule InstLAllR, $\Gamma \vdash \hat{\alpha} \preceq \forall \alpha. A_0 \dashv \Delta$.

Suppose not.

Then there is no derivation, since InstLAllR is the only rule matching $\forall \alpha. A_0$.

(2) $\Gamma \vdash A \preceq \hat{\alpha} \dashv \Delta$ is decidable.

- **Case UvarWF:**

Similar to the UvarWF case in part (1), but applying rule InstRSolve instead of InstLSolve.

- **Case UnitWF:** Apply InstRSolve.

- **Case**

$$\frac{\Gamma_L, \hat{\alpha}, \Gamma_R \vdash \hat{\beta}}{\Gamma} \text{EvarWF}$$

Similar to the EvarWF case in part (1), but applying InstRSolve/InstRReach instead of InstLSolve/InstLReach.

- **Case SolvedEvarWF:**

Impossible, for exactly the same reasons as in the SolvedEvarWF case of part (1).

- **Case**

$$\frac{\Gamma \vdash A_1 \quad \Gamma \vdash A_2}{\Gamma_L, \hat{\alpha}, \Gamma_R \vdash A_1 \rightarrow A_2} \text{ArrowWF}$$

As the ArrowWF case of part (1), except applying InstRArr instead of InstLArr.

- **Case**

$$\frac{\Gamma, \beta \vdash B}{\Gamma_L, \hat{\alpha}, \Gamma_R \vdash \forall \beta. B} \text{ForallWF}$$

By assumption, $[\Gamma](\forall \beta. B) = \forall \beta. B$. With the definition of substitution, we get $[\Gamma]B = B$. Hence $[\Gamma]B = B$.

Hence $[\hat{\beta}/\beta][\Gamma]B = [\hat{\beta}/\beta]B$. Since $\hat{\beta}$ is fresh, $[\hat{\beta}/\beta][\Gamma]B = [\Gamma][\hat{\beta}/\beta]B$.

By definition of substitution, $[\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta}][\hat{\beta}/\beta]B = [\Gamma][\hat{\beta}/\beta]B$, which by transitivity of equality is $[\hat{\beta}/\beta]B$.

We have $\hat{\alpha} \notin \text{FV}([\Gamma](\forall \beta. B))$, so $\hat{\alpha} \notin \text{FV}([\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta}][\hat{\beta}/\beta]B)$.

Therefore, by induction, either $\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta} \vdash [\hat{\beta}/\beta]B \preceq: \hat{\alpha} \dashv \Theta$ or not.

Suppose $\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta} \vdash [\hat{\beta}/\beta]B \preceq: \hat{\alpha} \dashv \Theta$.

By Lemma 32 (Instantiation Extension), $\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta} \longrightarrow \Theta$;

by Lemma 24 (Extension Order) (ii), $\Theta = \Delta, \blacktriangleright_{\hat{\beta}}, \Delta'$.

Hence by rule InstRAIIL, $\Gamma \vdash \forall \beta. B \preceq: \hat{\alpha} \dashv \Delta$.

Suppose not.

Then there is no derivation, since InstRAIIL is the only rule matching $\forall \beta. B$. □

F' Decidability of Algorithmic Subtyping

F'.1 Lemmas for Decidability of Subtyping

Lemma 37 (Monotypes Solve Variables). *If $\Gamma \vdash \hat{\alpha} \preceq: \tau \dashv \Delta$ or $\Gamma \vdash \tau \preceq: \hat{\alpha} \dashv \Delta$, then if $[\Gamma]\tau = \tau$ and $\hat{\alpha} \notin \text{FV}([\Gamma]\tau)$, then $|\text{unsolved}(\Gamma)| = |\text{unsolved}(\Delta)| + 1$.*

Proof. By induction on the given derivation.

- **Case**

$$\frac{\Gamma_L \vdash \tau}{\Gamma_L, \hat{\alpha}, \Gamma_R \vdash \hat{\alpha} \preceq: \tau \dashv \underbrace{\Gamma_L, \hat{\alpha} = \tau, \Gamma_R}_{\Delta}} \text{InstLSolve}$$

It is evident that $|\text{unsolved}(\Gamma_L, \hat{\alpha}, \Gamma_R)| = |\text{unsolved}(\Gamma_L, \hat{\alpha} = \tau, \Gamma_R)| + 1$.

- **Case**

$$\frac{}{\Gamma[\hat{\alpha}][\hat{\beta}] \vdash \hat{\alpha} \preceq: \hat{\beta} \dashv \Gamma[\hat{\alpha}][\hat{\beta} = \hat{\alpha}]} \text{InstLReach}$$

Similar to the previous case.

$$\bullet \text{ Case } \frac{\Gamma_0[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2] \vdash \tau_1 \leq \hat{\alpha}_1 \dashv \Theta \quad \Theta \vdash \hat{\alpha}_2 : \leq [\Theta]\tau_2 \dashv \Delta}{\Gamma_0[\hat{\alpha}] \vdash \hat{\alpha} : \leq \tau_1 \rightarrow \tau_2 \dashv \Delta} \text{InstLArr}$$

$$\begin{aligned} |\text{unsolved}(\Gamma_0[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2])| &= |\text{unsolved}(\Gamma_0[\hat{\alpha}])| + 1 && \text{Immediate} \\ |\text{unsolved}(\Gamma_0[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2])| &= |\text{unsolved}(\Theta)| + 1 && \text{By i.h.} \\ |\text{unsolved}(\Gamma)| &= |\text{unsolved}(\Theta)| && \text{Subtracting 1} \\ \text{---} &= |\text{unsolved}(\Delta)| + 1 && \text{By i.h.} \end{aligned}$$

$$\bullet \text{ Case } \frac{\Gamma, \beta \vdash \hat{\alpha} : \leq B \dashv \Delta, \beta, \Delta'}{\Gamma \vdash \hat{\alpha} : \leq \forall \beta. B \dashv \Delta} \text{InstLAllR}$$

This case is impossible, since a monotype cannot have the form $\forall \beta. B$.

• **Cases** InstRSolve, InstRReach: Similar to the InstLSolve and InstLReach cases.

• **Case** InstRArr: Similar to the InstLArr case.

$$\bullet \text{ Case } \frac{\Gamma[\hat{\alpha}], \beta \vdash B \leq \hat{\alpha} \dashv \Delta, \beta, \Delta'}{\Gamma[\hat{\alpha}] \vdash \forall \beta. B \leq \hat{\alpha} \dashv \Delta} \text{InstRAIlL}$$

This case is impossible, since a monotype cannot have the form $\forall \beta. B$. □

Lemma 38 (Monotype Monotonicity). *If $\Gamma \vdash \tau_1 < \tau_2 \dashv \Delta$ then $|\text{unsolved}(\Delta)| \leq |\text{unsolved}(\Gamma)|$.*

Proof. By induction on the given derivation.

• **Cases** $< : \text{Var}$, $< : \text{Exvar}$:

In these rules, $\Delta = \Gamma$, so $\text{unsolved}(\Delta) = \text{unsolved}(\Gamma)$; therefore $|\text{unsolved}(\Delta)| \leq |\text{unsolved}(\Gamma)|$.

• **Case** $< : \rightarrow$: We have an intermediate context Θ .

By inversion, $\tau_1 = \tau_{11} \rightarrow \tau_{12}$ and $\tau_2 = \tau_{21} \rightarrow \tau_{22}$. Therefore, we have monotypes in the first and second premises.

By induction on the first premise, $|\text{unsolved}(\Theta)| \leq |\text{unsolved}(\Gamma)|$. By induction on the second premise, $|\text{unsolved}(\Delta)| \leq |\text{unsolved}(\Theta)|$. By transitivity of \leq , $|\text{unsolved}(\Delta)| \leq |\text{unsolved}(\Gamma)|$, which was to be shown.

• **Cases** $< : \forall L$, $< : \forall R$: We are given a derivation of subtyping on monotypes, so these cases are impossible.

• **Cases** $< : \text{InstantiateL}$, $< : \text{InstantiateR}$: The input and output contexts in the premise exactly match the conclusion, so the result follows by Lemma 37 (Monotypes Solve Variables). □

Lemma 39 (Substitution Decreases Size). *If $\Gamma \vdash A$ then $|\Gamma \vdash [\Gamma]A| \leq |\Gamma \vdash A|$.*

Proof. By induction on $|\Gamma \vdash A|$. If $A = 1$ or $A = \alpha$, or $A = \hat{\alpha}$ and $\hat{\alpha} \in \text{unsolved}(\Gamma)$ then $[\Gamma]A = A$. Therefore, $|\Gamma \vdash [\Gamma]A| = |\Gamma \vdash A|$.

If $A = \hat{\alpha}$ and $(\hat{\alpha} = \tau) \in \Gamma$, then by induction hypothesis, $|\Gamma \vdash [\Gamma]\tau| \leq |\Gamma \vdash \tau|$. Of course $|\Gamma \vdash \tau| \leq |\Gamma \vdash \tau| + 1$. By definition of substitution, $[\Gamma]\tau = [\Gamma]\hat{\alpha}$, so

$$|\Gamma \vdash [\Gamma]\hat{\alpha}| \leq |\Gamma \vdash \tau| + 1$$

By the definition of type size, $|\Gamma \vdash \hat{\alpha}| = |\Gamma \vdash \tau| + 1$, so

$$|\Gamma \vdash [\Gamma]\hat{\alpha}| \leq |\Gamma \vdash \hat{\alpha}|$$

which was to be shown.

If $A = A_1 \rightarrow A_2$, the result follows via the induction hypothesis (twice).

If $A = \forall \alpha. A_0$, the result follows via the induction hypothesis. □

Lemma 40 (Monotype Context Invariance).

If $\Gamma \vdash \tau <: \tau' \dashv \Delta$ where $[\Gamma]\tau = \tau$ and $[\Gamma]\tau' = \tau'$ and $|\text{unsolved}(\Gamma)| = |\text{unsolved}(\Delta)|$ then $\Gamma = \Delta$.

Proof. By induction on the derivation of $\Gamma \vdash \tau <: \tau' \dashv \Delta$.

- **Cases** $<:\text{Var}$, $<:\text{Unit}$, $<:\text{Exvar}$:

In these rules, the output context is the same as the input context, so the result is immediate.

- **Case**
$$\frac{\Gamma \vdash \tau'_1 <: \tau_1 \dashv \Theta \quad \Theta \vdash [\Theta]\tau_2 <: [\Theta]\tau'_2 \dashv \Delta}{\Gamma \vdash \tau_1 \rightarrow \tau_2 <: \tau'_1 \rightarrow \tau'_2 \dashv \Delta} <:\rightarrow$$

We have that $[\Gamma](\tau_1 \rightarrow \tau_2) = \tau_1 \rightarrow \tau_2$. By definition of substitution, $[\Gamma]\tau_1 = \tau_1$ and $[\Gamma]\tau_2 = \tau_2$. Similarly, $[\Gamma]\tau_1 = \tau'_1$ and $[\Gamma]\tau_2 = \tau'_2$.

By i.h., $\Theta = \Gamma$.

Since Θ is predicative, $[\Theta]\tau_2$ and $[\Theta]\tau'_2$ are monotypes.

Substitution is idempotent: $[\Theta][\Theta]\tau_2 = [\Theta]\tau_2$ and $[\Theta][\Theta]\tau'_2 = [\Theta]\tau'_2$.

By i.h., $\Delta = \Theta$. Hence $\Delta = \Gamma$.

- **Cases** $<:\forall L$, $<:\forall R$: Impossible, since τ and τ' are monotypes.

- **Case**
$$\frac{\hat{\alpha} \notin \text{FV}(A) \quad \Gamma_0[\hat{\alpha}] \vdash \hat{\alpha} : \leq A \dashv \Delta}{\Gamma_0[\hat{\alpha}] \vdash \hat{\alpha} <: A \dashv \Delta} <:\text{InstantiateL}$$

By Lemma 37 (Monotypes Solve Variables), $|\text{unsolved}(\Delta)| < |\text{unsolved}(\Gamma_0[\hat{\alpha}])|$, but it is given that $|\text{unsolved}(\Gamma_0[\hat{\alpha}])| = |\text{unsolved}(\Delta)|$, so this case is impossible.

- **Case** $<:\text{InstantiateR}$: Impossible, as for the $<:\text{InstantiateL}$ case. □

F'.2 Decidability of Subtyping

Theorem 8 (Decidability of Subtyping).

Given a context Γ and types A, B such that $\Gamma \vdash A$ and $\Gamma \vdash B$ and $[\Gamma]A = A$ and $[\Gamma]B = B$, it is decidable whether there exists Δ such that $\Gamma \vdash A <: B \dashv \Delta$.

Proof. Let the judgment $\Gamma \vdash A <: B \dashv \Delta$ be measured lexicographically by

(S1) the number of \forall quantifiers in A and B ;

(S2) $|\text{unsolved}(\Gamma)|$, the number of unsolved existential variables in Γ ;

(S3) $|\Gamma \vdash A| + |\Gamma \vdash B|$.

For each subtyping rule, we show that every premise is smaller than the conclusion. The condition that $[\Gamma]A = A$ and $[\Gamma]B = B$ is easily satisfied at each inductive step, using the definition of substitution.

- Rules $<:\text{Var}$, $<:\text{Unit}$ and $<:\text{Exvar}$ have no premises.

- **Case**
$$\frac{\Gamma \vdash B_1 <: A_1 \dashv \Theta \quad \Theta \vdash [\Theta]A_2 <: [\Theta]B_2 \dashv \Delta}{\Gamma \vdash A_1 \rightarrow A_2 <: B_1 \rightarrow B_2 \dashv \Delta} <:\rightarrow$$

If A_2 or B_2 has a quantifier, then the first premise is smaller by (S1). Otherwise, the first premise shares an input context with the conclusion, so it has the same (S2). The types B_1 and A_1 are subterms of the conclusion's types, so the first premise is smaller by (S3).

If B_1 or A_1 has a quantifier, then the second premise is smaller by (S1). Otherwise, by Lemma 38 (Monotype Monotonicity) on the first premise, $|\text{unsolved}(\Theta)| \leq |\text{unsolved}(\Gamma)|$.

- If $|\text{unsolved}(\Theta)| < |\text{unsolved}(\Gamma)|$, then the second premise is smaller by (S2).

– If $|\text{unsolved}(\Theta)| = |\text{unsolved}(\Gamma)|$, we have the same (S2).

However, by Lemma 40 (Monotype Context Invariance), $\Theta = \Gamma$, so $|\Theta \vdash [\Theta]A_2| = |\Gamma \vdash [\Gamma]A_2|$, which by Lemma 39 (Substitution Decreases Size) is less than or equal to $|\Gamma \vdash A_2|$.

By the same logic, $|\Theta \vdash [\Theta]B_2| \leq |\Gamma \vdash B_2|$.

Therefore,

$$|\Theta \vdash [\Theta]A_2| + |\Theta \vdash [\Theta]B_2| \leq |\Gamma \vdash (A_1 \rightarrow A_2)| + |\Gamma \vdash (B_1 \rightarrow B_2)|$$

and the second premise is smaller by (S3).

- **Cases \prec : \forall L, \prec : \forall R:** In each of these rules, the premise has one less quantifier than the conclusion, so the premise is smaller by (S1).
- **Cases \prec :InstantiateL, \prec :InstantiateR:** Follows from Theorem 7. □

G' Decidability of Typing

Theorem 9 (Decidability of Typing).

- (i) **Synthesis:** Given a context Γ and a term e , it is decidable whether there exist a type A and a context Δ such that $\Gamma \vdash e \Rightarrow A \dashv \Delta$.
- (ii) **Checking:** Given a context Γ , a term e , and a type B such that $\Gamma \vdash B$, it is decidable whether there is a context Δ such that $\Gamma \vdash e \Leftarrow B \dashv \Delta$.
- (iii) **Application:** Given a context Γ , a term e , and a type A such that $\Gamma \vdash A$, it is decidable whether there exist a type C and a context Δ such that $\Gamma \vdash A \bullet e \Rightarrow C \dashv \Delta$.

Proof. For rules deriving judgments of the form

$$\begin{aligned} \Gamma \vdash e &\Rightarrow - \dashv - \\ \Gamma \vdash e &\Leftarrow B \dashv - \\ \Gamma \vdash A \bullet e &\Rightarrow - \dashv - \end{aligned}$$

(where we write “ $-$ ” for parts of the judgments that are outputs), the following induction measure on such judgments is adequate to prove decidability:

$$\left\langle e, \begin{array}{l} \Rightarrow \\ \Leftarrow, |\Gamma \vdash B| \\ \Rightarrow, |\Gamma \vdash A| \end{array} \right\rangle$$

where $\langle \dots \rangle$ denotes lexicographic order, and where (when comparing two judgments typing terms of the same size) the synthesis judgment (top line) is considered smaller than the checking judgment (second line), which in turn is considered smaller than the application judgment (bottom line). That is,

$$\Rightarrow \prec \Leftarrow \prec \Rightarrow$$

Note that this measure only uses the input parts of the judgments, leading to a straightforward decidability argument.

We will show that in each rule, every synthesis/checking/application premise is smaller than the conclusion.

- **Case Var:** No premises.

- **Case Sub:** The first premise has the same subject term e as the conclusion, but the judgment is smaller because the measure considers a synthesis judgment to be smaller than a checking judgment.
The second premise is a subtyping judgment, which by Theorem 8 is decidable.
- **Case Anno:**
It is easy to show that the judgment $\Gamma \vdash A$ is decidable.
The second premise types e , but the conclusion types $(e : A)$, so the first part of the measure gets smaller.
- **Case ! :** No premises.
- **Case $\rightarrow\text{!}$:** In the premise, the term is smaller.
- **Case $\rightarrow\text{E}$:** In both premises, the term is smaller.
- **Case $\forall\text{!}$:** Both the premise and conclusion type e , and both are checking; however, $|\Gamma, \alpha \vdash A| < |\Gamma \vdash \forall \alpha. A|$, so the premise is smaller.
- **Case $\rightarrow\text{App}$:** Both the premise and conclusion type e , but the premise is a checking judgment, so the premise is smaller.
- **Case $\text{Subst}\Leftarrow$:** Both the premise and conclusion type e , and both are checking; however, since we can apply this rule only when Γ has a solution for $\hat{\alpha}$ —that is, when $\Gamma = \Gamma_0[\hat{\alpha} = \tau]$ —we have that $|\Gamma \vdash [\Gamma]\hat{\alpha}| < |\Gamma \vdash \hat{\alpha}|$, making the last part of the measure smaller.
- **Case SubstApp :** Similar to $\text{Subst}\Leftarrow$.
- **Case $\forall\text{App}$:** Both the premise and conclusion type e , and both are application judgments; however, by the definition of $|\Gamma \vdash -|$, the size of the type in the premise $[\hat{\alpha}/\alpha]A$ is smaller than $\forall \alpha. A$.
- **Case $\hat{\alpha}\text{App}$:** Both the premise and conclusion type e , but we switch to checking in the premise, so the premise is smaller.
- **Case $\text{!}\Rightarrow$:** No premises.
- **Case $\rightarrow\text{!}\Rightarrow$:** In the premise, the term is smaller. □

H' Soundness of Subtyping

H'.1 Lemmas for Soundness

Lemma 42 (Variable Preservation).

If $(x : A) \in \Delta$ or $(x : A) \in \Omega$ and $\Delta \longrightarrow \Omega$ then $(x : [\Omega]A) \in [\Omega]\Delta$.

Proof. By mutual induction on Δ and Ω .

Suppose $(x : A) \in \Delta$. In the case where $\Delta = (\Delta', x : A)$ and $\Omega = (\Omega', x : A_\Omega)$, inversion on $\Delta \longrightarrow \Omega$ gives $[\Omega']A = [\Omega']A_\Omega$; by the definition of context application, $[\Omega', x : A_\Omega](\Delta', x : A) = [\Omega']\Delta', x : [\Omega']A_\Omega$, which contains $x : [\Omega']A_\Omega$, which is equal to $x : [\Omega']A$. By well-formedness of Ω , we know that $[\Omega']A = [\Omega]A$.

Suppose $(x : A) \in \Omega$. The reasoning is similar, because equality is symmetric. □

Lemma 43 (Substitution Typing). If $\Gamma \vdash A$ then $\Gamma \vdash [\Gamma]A$.

Proof. By induction on $|\Gamma \vdash A|$ (the size of A under Γ).

- **Cases UvarWF, UnitWF:** Here $A = \alpha$ or $A = 1$, so applying Γ to A does not change it: $A = [\Gamma]A$. Since $\Gamma \vdash A$, we have $\Gamma \vdash [\Gamma]A$, which was to be shown.

- **Case EvarWF:** In this case $A = \hat{\alpha}$, but $\Gamma = \Gamma_0[\hat{\alpha}]$, so applying Γ to A does not change it, and we proceed as in the UnitWF case above.
- **Case SolvedEvarWF:** In this case $A = \hat{\alpha}$ and $\Gamma = \Gamma_L, \hat{\alpha} = \tau, \Gamma_R$. Thus $[\Gamma]A = [\Gamma]\alpha = [\Gamma_L]\tau$. We assume contexts are well-formed, so all free variables in τ are declared in Γ_L . Consequently, $|\Gamma_L \vdash \tau| = |\Gamma \vdash \tau|$, which is less than $|\Gamma \vdash \hat{\alpha}|$. We can therefore apply the i.h. to τ , yielding $\Gamma \vdash [\Gamma]\tau$. By the definition of substitution, $[\Gamma]\tau = [\Gamma]\hat{\alpha}$, so we have $\Gamma \vdash [\Gamma]\hat{\alpha}$.
- **Case ArrowWF:** In this case $A = A_1 \rightarrow A_2$. By i.h., $\Gamma \vdash [\Gamma]A_1$ and $\Gamma \vdash [\Gamma]A_2$. By ArrowWF, $\Gamma \vdash ([\Gamma]A_1) \rightarrow ([\Gamma]A_2)$, which by the definition of substitution is $\Gamma \vdash [\Gamma](A_1 \rightarrow A_2)$.
- **Case ForallWF:** In this case $A = \forall\alpha. A_0$. By i.h., $\Gamma, \alpha \vdash [\Gamma, \alpha]A_0$. By the definition of substitution, $[\Gamma, \alpha]A_0 = [\Gamma]A_0$, so by ForallWF, $\Gamma \vdash \forall\alpha. [\Gamma]A_0$, which by the definition of substitution is $\Gamma \vdash [\Gamma](\forall\alpha. A_0)$. \square

Lemma 44 (Substitution for Well-Formedness). *If $\Omega \vdash A$ then $[\Omega]\Omega \vdash [\Omega]A$.*

Proof. By induction on $|\Omega \vdash A|$, the size of A under Ω (Definition 2).

We consider cases of the well-formedness rule concluding the derivation of $\Omega \vdash A$.

- **Case**

$$\frac{}{\Omega \vdash 1} \text{UnitWF}$$

$[\Omega]\Omega \vdash 1$ By DeclUnitWF
 $[\Omega]\Omega \vdash [\Omega]1$ By definition of substitution

- **Case**

$$\frac{\underbrace{\Omega'[\alpha] \vdash \alpha}_{\Omega} \text{UvarWF}}{\Omega \rightarrow \Omega} \text{UvarWF}$$

$\alpha \in [\Omega]\Omega$ By Lemma 41 (Uvar Preservation)
 $[\Omega]\Omega \vdash \alpha$ By DeclUvarWF
 $[\Omega]\Omega \vdash [\Omega]\alpha$ By definition of substitution

- **Case**

$$\frac{\underbrace{\Omega'[\hat{\alpha} = \tau] \vdash \hat{\alpha}}_{\Omega} \text{SolvedEvarWF}}{\Omega \rightarrow \Omega} \text{SolvedEvarWF}$$

$\Omega \vdash \hat{\alpha}$ Given
 $\Omega \rightarrow \Omega$ By Lemma 20 (Reflexivity)
 $\Omega \vdash [\Omega]\hat{\alpha}$ By Lemma 43 (Substitution Typing)
 $|\Omega \vdash [\Omega]\hat{\alpha}| < |\Omega \vdash \hat{\alpha}|$ Follows from definition of type size
 $[\Omega]\Omega \vdash [\Omega][\Omega]\hat{\alpha}$ By i.h.
 $[\Omega][\Omega]\hat{\alpha} = [\Omega]\hat{\alpha}$ By Lemma 18 (Substitution Extension Invariance)
 $[\Omega]\Omega \vdash [\Omega]\hat{\alpha}$ Applying equality

- **Case**

$$\frac{\underbrace{\Omega'[\hat{\alpha}] \vdash \hat{\alpha}}_{\Omega} \text{EvarWF}}{\Omega \rightarrow \Omega} \text{EvarWF}$$

Impossible: the grammar for Ω does not allow unsolved declarations.

- **Case**
- $$\frac{\Omega \vdash A_1 \quad \Omega \vdash A_2}{\Omega \vdash A_1 \rightarrow A_2} \text{ArrowWF}$$

$\Omega \vdash A_1$	Subderivation
$ \Omega \vdash A_1 < \Omega \vdash A_1 \rightarrow A_2 $	Follows from definition of type size
$[\Omega]\Omega \vdash [\Omega]A_1$	By i.h.
$[\Omega]\Omega \vdash [\Omega]A_2$	By similar reasoning on 2nd subderivation
$[\Omega]\Omega \vdash [\Omega]A_1 \rightarrow [\Omega]A_2$	By DeclArrowWF
$[\Omega]\Omega \vdash [\Omega](A_1 \rightarrow A_2)$	By definition of substitution

• **Case** $\frac{\Omega, \alpha \vdash A_0}{\Omega \vdash \forall \alpha. A_0}$ ForallWF

$\Omega, \alpha \vdash A_0$	Subderivation
Let $\Omega' = (\Omega, \alpha)$.	
$ \Omega' \vdash A_0 < \Omega \vdash \forall \alpha. A_0 $	Follows from definition of type size
$[\Omega'](\Omega, \alpha) \vdash [\Omega']A_0$	By i.h.
$[\Omega]\Omega, \alpha \vdash [\Omega']A_0$	By definition of context application
$[\Omega]\Omega, \alpha \vdash [\Omega]A_0$	By definition of substitution
$[\Omega]\Omega \vdash \forall \alpha. [\Omega]A_0$	By DeclForallWF
$[\Omega]\Omega \vdash [\Omega](\forall \alpha. A_0)$	By definition of substitution

□

Lemma 45 (Substitution Stability).

For any well-formed complete context (Ω, Ω_Z) , if $\Omega \vdash A$ then $[\Omega]A = [\Omega, \Omega_Z]A$.

Proof. By induction on Ω_Z . If $\Omega_Z = \cdot$, the result is immediate. Otherwise, use the i.h. and the fact that $\Omega \vdash A$ implies $\text{FV}(A) \cap \text{dom}(\Omega_Z) = \emptyset$. □

Lemma 46 (Context Partitioning).

If $\Delta, \blacktriangleright_{\hat{\alpha}}, \Theta \longrightarrow \Omega, \blacktriangleright_{\hat{\alpha}}, \Omega_Z$ then there is a Ψ such that $[\Omega, \blacktriangleright_{\hat{\alpha}}, \Omega_Z](\Delta, \blacktriangleright_{\hat{\alpha}}, \Theta) = [\Omega]\Delta, \Psi$.

Proof. By induction on the given derivation.

- **Case** $\longrightarrow \text{ID}$: Impossible: $\Delta, \blacktriangleright_{\hat{\alpha}}, \Theta$ cannot have the form \cdot .
- **Case** $\longrightarrow \text{Var}$: We have $\Omega_Z = (\Omega'_Z, x : A)$ and $\Theta = (\Theta', x : A')$. By i.h., there is Ψ' such that $[\Omega, \blacktriangleright_{\hat{\alpha}}, \Omega'_Z](\Delta, \blacktriangleright_{\hat{\alpha}}, \Theta') = [\Omega]\Delta, \Psi'$. Then by the definition of context application, $[\Omega, \blacktriangleright_{\hat{\alpha}}, \Omega'_Z, x : A](\Delta, \blacktriangleright_{\hat{\alpha}}, \Theta', x : A') = [\Omega]\Delta, \Psi', x : [\Omega']A$. Let $\Psi = (\Psi', x : [\Omega']A)$.
- **Case** $\longrightarrow \text{Uvar}$: Similar to the $\longrightarrow \text{Var}$ case, with $\Psi = (\Psi', \alpha)$.
- **Cases** $\longrightarrow \text{Unsolved}$, $\longrightarrow \text{Solve}$, $\longrightarrow \text{Marker}$, $\longrightarrow \text{Add}$, $\longrightarrow \text{AddSolved}$: Broadly similar to the $\longrightarrow \text{Uvar}$ case, but since the rightmost context element is soft it disappears in context application, so we let $\Psi = \Psi'$. □

Lemma 49 (Stability of Complete Contexts).

If $\Gamma \longrightarrow \Omega$ then $[\Omega]\Gamma = [\Omega]\Omega$.

Proof. By induction on the derivation of $\Gamma \longrightarrow \Omega$.

• **Case** $\frac{}{\cdot \longrightarrow \cdot} \longrightarrow \text{ID}$

In this case, $\Omega = \Gamma = \cdot$.

By definition, $[\cdot]\cdot = \cdot$, which gives us the conclusion.

- **Case** $\frac{\Gamma' \longrightarrow \Omega' \quad [\Omega']A_\Gamma = [\Omega']A}{\Gamma', x : A_\Gamma \longrightarrow \Omega', x : A} \longrightarrow \text{Var}$

$[\Omega']\Gamma' = [\Omega']\Omega'$ By i.h.
 $[\Omega']A_\Gamma = [\Omega']A$ Premise

$[\Omega]\Gamma = [\Omega', x : A](\Gamma', x : A_\Gamma)$ Expanding Ω and Γ
 $= [\Omega']\Gamma', x : [\Omega']A_\Gamma$ By definition of context application
(using $[\Omega']A_\Gamma = [\Omega']A$)
 $= [\Omega']\Omega', x : [\Omega']A$ By above equalities
 $= [\Omega]\Omega$ By definition of context application
- **Case** $\frac{\Gamma' \longrightarrow \Omega'}{\Gamma', \alpha \longrightarrow \Omega', \alpha} \longrightarrow \text{Uvar}$

$[\Omega]\Gamma = [\Omega', \alpha](\Gamma', \alpha)$ Expanding Ω and Γ
 $= [\Omega']\Gamma', \alpha$ By definition of context application
 $= [\Omega']\Omega', \alpha$ By i.h.
 $= \Omega', \alpha$ By definition of context application
 $= [\Omega]\Omega$ By $\Omega = (\Omega', \alpha)$
- **Case** $\frac{\Gamma' \longrightarrow \Omega'}{\Gamma', \blacktriangleright \hat{\alpha} \longrightarrow \Omega', \blacktriangleright \hat{\alpha}} \longrightarrow \text{Marker}$

Similar to the $\longrightarrow \text{Uvar}$ case.
- **Case** $\frac{\Gamma \longrightarrow \Omega'}{\Gamma \longrightarrow \Omega', \hat{\alpha} = \tau} \longrightarrow \text{AddSolved}$

$[\Omega]\Gamma = [\Omega', \hat{\alpha} = \tau]\Gamma$ Expanding Ω
 $= [\Omega']\Gamma$ By $\hat{\alpha} \notin \text{dom}(\Gamma)$
 $= [\Omega']\Omega'$ By i.h.
 $= \Omega', \hat{\alpha} = \tau$ By definition of context application
 $= [\Omega]\Omega$ By $\Omega = (\Omega', \hat{\alpha} = \tau)$
- **Case** $\frac{\Gamma' \longrightarrow \Omega' \quad [\Omega']\tau_\Gamma = [\Omega']\tau}{\Gamma', \hat{\alpha} = \tau_\Gamma \longrightarrow \Omega', \hat{\alpha} = \tau} \longrightarrow \text{Solved}$

$[\Omega]\Gamma = [\Omega', \hat{\alpha} = \tau](\Gamma', \hat{\alpha} = \tau_\Gamma)$ Expanding Ω and Γ
 $= [\Omega']\Gamma'$ By definition of context application
 $= [\Omega']\Omega'$ By i.h.
 $= \Omega', \hat{\alpha} = \tau$ By definition of context application
 $= [\Omega]\Omega$ By $\Omega = (\Omega', \hat{\alpha} = \tau)$
- **Case** $\frac{\Gamma' \longrightarrow \Omega'}{\Gamma', \hat{\alpha} \longrightarrow \Omega', \hat{\alpha} = \tau} \longrightarrow \text{Solve}$

$[\Omega]\Gamma = [\Omega', \hat{\alpha} = \tau](\Gamma', \hat{\alpha})$ Expanding Ω and Γ
 $= [\Omega']\Gamma'$ By definition of context application
 $= [\Omega']\Omega'$ By i.h.
 $= \Omega', \hat{\alpha} = \tau$ By definition of context application
 $= [\Omega]\Omega$ By $\Omega = (\Omega', \hat{\alpha} = \tau)$
- **Case** $\frac{\Gamma \longrightarrow \Delta}{\Gamma, \hat{\alpha} \longrightarrow \Delta, \hat{\alpha}} \longrightarrow \text{Unsolved}$

Impossible: Ω cannot have the form $\Delta, \hat{\alpha}$.

- **Case**
$$\frac{\Gamma \longrightarrow \Delta}{\Gamma \longrightarrow \Delta, \hat{\alpha}} \longrightarrow \text{Add}$$

Impossible: Ω cannot have the form $\Delta, \hat{\alpha}$. □

Lemma 50 (Finishing Types).

If $\Omega \vdash A$ and $\Omega \longrightarrow \Omega'$ then $[\Omega]A = [\Omega']A$.

Proof. By Lemma 18 (Substitution Extension Invariance), $[\Omega']A = [\Omega'][\Omega]A$.

If $\text{FEV}(C) = \emptyset$ then $[\Omega']C = C$.

Since Ω is complete and $\Omega \vdash A$, we have $\text{FEV}([\Omega]A) = \emptyset$. Therefore $[\Omega'][\Omega]A = [\Omega]A$. □

Lemma 51 (Finishing Completions).

If $\Omega \longrightarrow \Omega'$ then $[\Omega]\Omega = [\Omega']\Omega'$.

Proof. By induction on the given derivation of $\Omega \longrightarrow \Omega'$.

Only cases $\longrightarrow \text{ID}$, $\longrightarrow \text{Var}$, $\longrightarrow \text{Uvar}$, $\longrightarrow \text{Solved}$, $\longrightarrow \text{Marker}$ and $\longrightarrow \text{AddSolved}$ are possible. In all of these cases, we use the i.h. and the definition of context application; in cases $\longrightarrow \text{Var}$ and $\longrightarrow \text{Solved}$, we also use the equality in the premise of the respective rule. □

Lemma 52 (Confluence of Completeness).

If $\Delta_1 \longrightarrow \Omega$ and $\Delta_2 \longrightarrow \Omega$ then $[\Omega]\Delta_1 = [\Omega]\Delta_2$.

Proof.

- | | |
|---------------------------------------|--|
| $\Delta_1 \longrightarrow \Omega$ | Given |
| $[\Omega]\Delta_1 = [\Omega]\Omega$ | By Lemma 49 (Stability of Complete Contexts) |
| $\Delta_2 \longrightarrow \Omega$ | Given |
| $[\Omega]\Delta_2 = [\Omega]\Omega$ | By Lemma 49 (Stability of Complete Contexts) |
| $[\Omega]\Delta_1 = [\Omega]\Delta_2$ | By transitivity of equality |
-

H'.2 Instantiation Soundness

Theorem 10 (Instantiation Soundness).

Given $\Delta \longrightarrow \Omega$ and $[\Gamma]B = B$ and $\hat{\alpha} \notin \text{FV}(B)$:

(1) If $\Gamma \vdash \hat{\alpha} : \preceq B \dashv \Delta$ then $[\Omega]\Delta \vdash [\Omega]\hat{\alpha} \leq [\Omega]B$.

(2) If $\Gamma \vdash B \preceq \hat{\alpha} \dashv \Delta$ then $[\Omega]\Delta \vdash [\Omega]B \leq [\Omega]\hat{\alpha}$.

Proof. By induction on the given instantiation derivation.

- (1) • **Case**
- $$\frac{\Gamma_0 \vdash \tau}{\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_{\Gamma} \vdash \hat{\alpha} : \preceq \tau \dashv \underbrace{\Gamma_0, \hat{\alpha} = \tau, \Gamma_1}_{\Delta}} \text{InstLSolve}$$

In this case $[\Delta]\hat{\alpha} = [\Delta]\tau$. By reflexivity of subtyping (Lemma 3 (Reflexivity of Declarative Subtyping)), $[\Omega]\Delta \vdash [\Delta]\hat{\alpha} \leq [\Delta]\tau$.

- **Case**

$$\frac{\Gamma[\hat{\alpha}][\hat{\beta}] \vdash \hat{\alpha} : \preceq \hat{\beta} \dashv \Gamma[\hat{\alpha}][\hat{\beta} = \hat{\alpha}]}{\Delta} \text{InstLReach}$$

We have $\Delta = \Gamma[\hat{\alpha}][\hat{\beta} = \hat{\alpha}]$. Therefore $[\Delta]\hat{\alpha} = \hat{\alpha} = [\Delta]\hat{\beta}$.

By reflexivity of subtyping (Lemma 3 (Reflexivity of Declarative Subtyping)), $[\Omega]\Delta \vdash [\Delta]\hat{\alpha} \leq [\Delta]\hat{\beta}$.

• **Case**

$$\frac{\overbrace{\Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2] \vdash A_1 \leq \hat{\alpha}_1 \dashv \Gamma'}^{\Gamma_1} \quad \Gamma' \vdash \hat{\alpha}_2 : \leq [\Gamma']A_2 \dashv \Delta}{\Gamma[\hat{\alpha}] \vdash \hat{\alpha} : \leq A_1 \rightarrow A_2 \dashv \Delta} \text{InstLArr}$$

$$\begin{array}{ll} [\Gamma](A_1 \rightarrow A_2) = [\Gamma_1](A_1 \rightarrow A_2) & \hat{\alpha} \notin \text{FV}(A_1 \rightarrow A_2) \\ \hat{\alpha}_1, \hat{\alpha}_2 \notin \text{FV}(A_1) \cup \text{FV}(A_2) & \hat{\alpha}_1, \hat{\alpha}_2 \text{ fresh} \\ \Gamma' \vdash \hat{\alpha}_2 : \leq [\Gamma']A_2 \dashv \Delta & \text{Subderivation} \\ \Gamma' \longrightarrow \Delta & \text{By Lemma 32 (Instantiation Extension)} \\ \Delta \longrightarrow \Omega & \text{Given} \\ \Gamma' \longrightarrow \Omega & \text{By Lemma 21 (Transitivity)} \\ \\ \Gamma_1 \vdash A_1 \leq \hat{\alpha}_1 \dashv \Gamma' & \text{Subderivation} \\ [\Omega]\Delta \vdash [\Omega]A_1 \leq [\Omega]\hat{\alpha}_1 & \text{By i.h. and Lemma 52 (Confluence of Completeness)} \\ \\ \Gamma' \vdash \hat{\alpha}_2 : \leq [\Gamma']A_2 \dashv \Delta & \text{Subderivation} \\ [\Omega]\Delta \vdash [\Omega][\Gamma']\hat{\alpha}_2 \leq [\Omega][\Gamma']A_2 & \text{By i.h.} \\ \Gamma' \longrightarrow \Omega & \text{Above} \\ [\Omega]\Delta \vdash [\Omega]\hat{\alpha}_2 \leq [\Omega]A_2 & \text{By Lemma 18 (Substitution Extension Invariance)} \\ \\ [\Omega]\Delta \vdash [\Omega](\hat{\alpha}_1 \rightarrow \hat{\alpha}_2) \leq [\Omega]A_1 \rightarrow [\Omega]A_2 & \text{By } \leq \rightarrow \text{ and definition of substitution} \end{array}$$

Since $(\hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2) \in \Gamma_1$ and $\Gamma_1 \longrightarrow \Delta$, we know that $[\Omega]\hat{\alpha} = [\Omega](\hat{\alpha}_1 \rightarrow \hat{\alpha}_2)$.
Therefore $[\Omega]\Delta \vdash [\Omega]\hat{\alpha} \leq [\Omega](A_1 \rightarrow A_2)$.

• **Case**

$$\frac{\Gamma[\hat{\alpha}], \beta \vdash \hat{\alpha} : \leq B_0 \dashv \Delta, \beta, \Delta'}{\Gamma[\hat{\alpha}] \vdash \hat{\alpha} : \leq \forall \beta. B_0 \dashv \Delta} \text{InstLAllR}$$

We have $\Delta \longrightarrow \Omega$ and $[\Gamma[\hat{\alpha}]](\forall \beta. B_0) = \forall \beta. B_0$ and $\hat{\alpha} \notin \text{FV}(\forall \beta. B_0)$.

Hence $\hat{\alpha} \notin \text{FV}(B_0)$ and by definition, $[\Gamma[\hat{\alpha}], \beta]B_0 = B_0$.

By Lemma 48 (Filling Completes), $\Delta, \beta, \Delta' \longrightarrow \Omega, \beta, |\Delta'|$.

By induction, $[\Omega, \beta, |\Delta'|](\Delta, \beta, \Delta') \vdash [\Omega, \beta, |\Delta'|]\hat{\alpha} \leq [\Omega, \beta, |\Delta'|]B_0$.

Each free variable in $\hat{\alpha}$ and B_0 is declared in (Ω, β) , so $\Omega, \beta, |\Delta'|$ behaves as $[\Omega, \beta]$ on $\hat{\alpha}$ and on B_0 , yielding $[\Omega, \beta, |\Delta'|](\Delta, \beta, \Delta') \vdash [\Omega, \beta]\hat{\alpha} \leq [\Omega, \beta]B_0$.

By Lemma 46 (Context Partitioning) and thinning, $[\Omega, \beta](\Delta, \beta) \vdash [\Omega, \beta]\hat{\alpha} \leq [\Omega, \beta]B_0$.

By the definition of context application, $[\Omega]\Delta, \beta \vdash [\Omega, \beta]\hat{\alpha} \leq [\Omega, \beta]B_0$.

By the definition of substitution, $[\Omega]\Delta, \beta \vdash [\Omega]\hat{\alpha} \leq [\Omega]B_0$.

Since $\hat{\alpha}$ is declared to the left of β , we have $\beta \notin \text{FV}([\Omega]\hat{\alpha})$.

Applying rule $\leq \forall L$ gives $[\Omega]\Delta \vdash [\Omega]\hat{\alpha} \leq \forall \beta. [\Omega]B_0$.

(2) • **Case**

$$\frac{\Gamma_0 \vdash \tau}{\underbrace{\Gamma_0, \hat{\alpha}, \Gamma_1}_{\Gamma} \vdash \tau \leq \hat{\alpha} \dashv \underbrace{\Gamma_0, \hat{\alpha} = \tau, \Gamma_1}_{\Gamma'}} \text{InstRSolve}$$

Similar to the InstLSolve case.

• **Case**

$$\frac{\Gamma[\hat{\alpha}][\hat{\beta}] \vdash \hat{\beta} \leq \hat{\alpha} \dashv \underbrace{\Gamma[\hat{\alpha}][\hat{\beta} = \hat{\alpha}]}_{\Gamma'}}{\Gamma[\hat{\alpha}][\hat{\beta}] \vdash \hat{\beta} \leq \hat{\alpha} \dashv \Delta} \text{InstRReach}$$

Similar to the InstLReach case.

• **Case**

$$\frac{\Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2] \vdash \hat{\alpha}_1 : \leq A_1 \dashv \Gamma' \quad \Gamma' \vdash [\Gamma']A_2 \leq \hat{\alpha}_2 \dashv \Delta}{\Gamma[\hat{\alpha}] \vdash A_1 \rightarrow A_2 \leq \hat{\alpha} \dashv \Delta} \text{InstRArr}$$

Similar to the InstLArr case.

• **Case**

$$\frac{\Gamma[\hat{\alpha}], \triangleright_{\hat{\beta}}, \hat{\beta} \vdash [\hat{\beta}/\beta]B_0 \leq \hat{\alpha} \dashv \Delta, \triangleright_{\hat{\beta}}, \Delta'}{\Gamma[\hat{\alpha}] \vdash \forall \beta. B_0 \leq \hat{\alpha} \dashv \Delta} \text{InstRAILL}$$

$$\begin{array}{ll}
[\Gamma[\hat{\alpha}]](\forall\beta. B_0) = \forall\beta. B_0 & \text{Given} \\
[\Gamma[\hat{\alpha}]]B_0 = B_0 & \\
[\Gamma[\hat{\alpha}], \blacktriangleright_{\hat{\beta}}, \hat{\beta}] [\hat{\beta}/\beta]B_0 = [\hat{\beta}/\beta]B_0 & \\
\Delta \longrightarrow \Omega & \text{Given} \\
\Delta, \blacktriangleright_{\hat{\beta}}, \Delta' \longrightarrow \Omega, \blacktriangleright_{\hat{\beta}}, |\Delta'| & \text{By Lemma 48 (Filling Completes)} \\
\hat{\alpha} \notin \text{FV}(\forall\beta. B_0) & \text{Given} \\
\hat{\alpha} \notin \text{FV}(B_0) & \text{By definition of FV}(-)
\end{array}$$

$$\begin{array}{ll}
\Gamma[\hat{\alpha}], \blacktriangleright_{\hat{\beta}}, \hat{\beta} \vdash [\hat{\beta}/\beta]B_0 \leq \hat{\alpha} \dashv \Delta, \blacktriangleright_{\hat{\beta}}, \Delta' & \text{Subderivation} \\
[\Omega, \blacktriangleright_{\hat{\beta}}, |\Delta'|](\Delta, \blacktriangleright_{\hat{\beta}}, \Delta') \vdash [\Omega, \blacktriangleright_{\hat{\beta}}, |\Delta'|][\hat{\beta}/\beta]B_0 \leq [\Omega, \blacktriangleright_{\hat{\beta}}, |\Delta'|]\hat{\alpha} & \text{By i.h.} \\
\Gamma[\hat{\alpha}], \blacktriangleright_{\hat{\beta}}, \hat{\beta} \longrightarrow \Delta, \blacktriangleright_{\hat{\beta}}, \Delta' & \text{By Lemma 32 (Instantiation Extension)}
\end{array}$$

By Lemma 16 (Declaration Order Preservation), $\hat{\alpha}$ is declared before $\blacktriangleright_{\hat{\beta}}$, that is, in Ω .

Thus, $[\Omega, \blacktriangleright_{\hat{\beta}}, |\Delta'|]\hat{\alpha} = [\Omega]\hat{\alpha}$.

By Lemma 23 (Evar Input), we know that Δ' is soft, so by Lemma 47 (Softness Goes Away),

$$[\Omega, \blacktriangleright_{\hat{\beta}}, |\Delta'|](\Delta, \blacktriangleright_{\hat{\beta}}, \Delta') = [\Omega, \blacktriangleright_{\hat{\beta}}](\Delta, \blacktriangleright_{\hat{\beta}}) = [\Omega]\Delta.$$

Applying these equalities to the derivation above gives

$$[\Omega]\Delta \vdash [\Omega, \blacktriangleright_{\hat{\beta}}, |\Delta'|][\hat{\beta}/\beta]B_0 \leq [\Omega]\hat{\alpha}$$

By distributivity of substitution,

$$[\Omega]\Delta \vdash [[\Omega, \blacktriangleright_{\hat{\beta}}, |\Delta'|][\hat{\beta}/\beta]][\Omega, \blacktriangleright_{\hat{\beta}}, |\Delta'|]B_0 \leq [\Omega]\hat{\alpha}$$

Furthermore, $[\Omega, \blacktriangleright_{\hat{\beta}}, |\Delta'|]B_0 = [\Omega]B_0$, since B_0 's free variables are either β or in Ω , giving

$$[\Omega]\Delta \vdash [[\Omega, \blacktriangleright_{\hat{\beta}}, |\Delta'|][\hat{\beta}/\beta]][\Omega]B_0 \leq [\Omega]\hat{\alpha}$$

Now apply $\leq\forall L$ and the definition of substitution to get $[\Omega]\Delta \vdash [\Omega](\forall\beta. B_0) \leq [\Omega]\hat{\alpha}$. \square

H'.3 Soundness of Subtyping

Theorem 11 (Soundness of Algorithmic Subtyping).

If $\Gamma \vdash A <: B \dashv \Delta$ where $[\Gamma]A = A$ and $[\Gamma]B = B$ and $\Delta \longrightarrow \Omega$ then $[\Omega]\Delta \vdash [\Omega]A \leq [\Omega]B$.

Proof. By induction on the derivation of $\Gamma \vdash A <: B \dashv \Delta$.

- **Case**

$$\begin{array}{ll}
\frac{}{\Gamma'[\alpha] \vdash \alpha <: \alpha \dashv \Gamma'[\alpha]} <:\text{Var} \\
\Gamma'[\alpha] \vdash \alpha <: \alpha \dashv \Gamma'[\alpha] & \\
\alpha \in \Delta & \Delta = \Gamma'[\alpha] \\
\alpha \in [\Omega]\Delta & \text{Follows from definition of context application} \\
[\Omega]\Delta \vdash \alpha \leq \alpha & \text{By } \leq\text{Var} \\
[\Omega]\Delta \vdash [\Omega]\alpha \leq [\Omega]\alpha & \text{By def. of substitution}
\end{array}$$

- **Case $<:\text{Unit}$:** Similar to the $<:\text{Var}$ case, applying rule $\leq\text{Unit}$ instead of $\leq\text{Var}$.

- **Case**

$$\begin{array}{ll}
\frac{}{\Gamma_L, \hat{\alpha}, \Gamma_R \vdash \hat{\alpha} <: \hat{\alpha} \dashv \Gamma_L, \hat{\alpha}, \Gamma_R} <:\text{Exvar} \\
[\Omega]\hat{\alpha} \text{ defined} & \text{Follows from definition of context application} \\
[\Omega]\Delta \vdash [\Omega]\hat{\alpha} & \text{Assumption that } [\Omega]\Delta \text{ is well-formed} \\
[\Omega]\Delta \vdash [\Omega]\hat{\alpha} \leq [\Omega]\hat{\alpha} & \text{By Lemma 3 (Reflexivity of Declarative Subtyping)}
\end{array}$$

• **Case**
$$\frac{\Gamma \vdash B_1 <: A_1 \dashv \Theta \quad \Theta \vdash [\Theta]A_2 <: [\Theta]B_2 \dashv \Delta}{\Gamma \vdash \underbrace{A_1 \rightarrow A_2}_A <: \underbrace{B_1 \rightarrow B_2}_B \dashv \Delta} <: \rightarrow$$

$\Gamma \vdash B_1 <: A_1 \dashv \Theta$	Subderivation
$\Delta \longrightarrow \Omega$	Given
$\Theta \longrightarrow \Omega$	By Lemma 21 (Transitivity)
$[\Omega]\Theta \vdash [\Omega]B_1 \leq [\Omega]A_1$	By i.h.
$[\Omega]\Delta \vdash [\Omega]B_1 \leq [\Omega]A_1$	By Lemma 52 (Confluence of Completeness)
$\Theta \vdash [\Theta]A_2 <: [\Theta]B_2 \dashv \Delta$	Subderivation
$[\Omega]\Delta \vdash [\Omega][\Theta]A_2 \leq [\Omega][\Theta]B_2$	By i.h.
$[\Omega][\Theta]A_2 = [\Omega]A_2$	By Lemma 18 (Substitution Extension Invariance)
$[\Omega][\Theta]B_2 = [\Omega]B_2$	By Lemma 18 (Substitution Extension Invariance)
$[\Omega]\Delta \vdash [\Omega]A_2 \leq [\Omega]B_2$	Above equations
$[\Omega]\Delta \vdash ([\Omega]A_1) \rightarrow ([\Omega]A_2) \leq ([\Omega]B_1) \rightarrow ([\Omega]B_2)$	By $\leq \rightarrow$
$[\Omega]\Delta \vdash [\Omega](A_1 \rightarrow A_2) \leq [\Omega](B_1 \rightarrow B_2)$	By def. of substitution

• **Case**
$$\frac{\Gamma, \triangleright_{\hat{\alpha}}, \hat{\alpha} \vdash [\hat{\alpha}/\alpha]A_0 <: B \dashv \Delta, \triangleright_{\hat{\alpha}}, \Theta}{\Gamma \vdash \forall \alpha. A_0 <: B \dashv \Delta} <: \forall L$$

Let $\Omega' = (\Omega, \triangleright_{\hat{\alpha}}, \Theta)$.

$\Gamma, \triangleright_{\hat{\alpha}}, \hat{\alpha} \vdash [\hat{\alpha}/\alpha]A_0 <: B \dashv \Delta, \triangleright_{\hat{\alpha}}, \Theta$	Subderivation
$\Delta \longrightarrow \Omega$	Given
$(\Delta, \triangleright_{\hat{\alpha}}, \Theta) \longrightarrow \Omega'$	By Lemma 48 (Filling Completes)
$[\Omega'](\Delta, \triangleright_{\hat{\alpha}}, \Theta) \vdash [\Omega'][\hat{\alpha}/\alpha]A_0 \leq [\Omega']B$	By i.h.
$[\Omega'](\Delta, \triangleright_{\hat{\alpha}}, \Theta) \vdash [\Omega'][\hat{\alpha}/\alpha]A_0 \leq [\Omega]B$	By $[\Omega']B = [\Omega]B$ (Lemma 45 (Substitution Stability))
$[\Omega'](\Delta, \triangleright_{\hat{\alpha}}, \Theta) \vdash [[\Omega']\hat{\alpha}/\alpha][\Omega']A_0 \leq [\Omega]B$	By distributivity of substitution
$\Gamma, \triangleright_{\hat{\alpha}}, \hat{\alpha} \vdash \hat{\alpha}$	By EvarWF
$\Gamma, \triangleright_{\hat{\alpha}}, \hat{\alpha} \longrightarrow \Delta, \triangleright_{\hat{\alpha}}, \Theta$	By Lemma 33 (Subtyping Extension)
$\Delta, \triangleright_{\hat{\alpha}}, \Theta \vdash \hat{\alpha}$	By Lemma 25 (Extension Weakening)
$(\Delta, \triangleright_{\hat{\alpha}}, \Theta) \longrightarrow \Omega'$	Above
$[\Omega']\Omega' \vdash [\Omega']\hat{\alpha}$	By Lemma 44 (Substitution for Well-Formedness)
$[\Omega'](\Delta, \triangleright_{\hat{\alpha}}, \Theta) \vdash [\Omega']\hat{\alpha}$	By Lemma 49 (Stability of Complete Contexts)
$[\Omega'](\Delta, \triangleright_{\hat{\alpha}}, \Theta) \vdash \forall \alpha. [\Omega']A_0 \leq [\Omega]B$	By $\leq \forall L$
$[\Omega'](\Delta, \triangleright_{\hat{\alpha}}, \Theta) \vdash \forall \alpha. [\Omega, \alpha]A_0 \leq [\Omega]B$	By Lemma 45 (Substitution Stability)
$[\Omega]\Delta \vdash \forall \alpha. [\Omega, \alpha]A_0 \leq [\Omega]B$	By Lemma 46 (Context Partitioning) and thinning
$[\Omega]\Delta \vdash \forall \alpha. [\Omega]A_0 \leq [\Omega]B$	By def. of substitution
$[\Omega]\Delta \vdash [\Omega](\forall \alpha. A_0) \leq [\Omega]B$	By def. of substitution

• **Case**
$$\frac{\Gamma, \alpha \vdash A <: B_0 \dashv \Delta, \alpha, \Theta}{\Gamma \vdash A <: \forall \alpha. B_0 \dashv \Delta} <: \forall R$$

$$\begin{array}{l}
\Gamma, \alpha \vdash A <: B_0 \dashv \Delta, \alpha, \Theta \quad \text{Subderivation} \\
\text{Let } \Omega_Z = |\Theta|. \\
\text{Let } \Omega' = (\Omega, \alpha, \Omega_Z). \\
(\Delta, \alpha, \Theta) \longrightarrow \Omega' \quad \text{By Lemma 48 (Filling Completes)} \\
[\Omega'](\Delta, \alpha, \Theta) \vdash [\Omega']A \leq [\Omega']B_0 \quad \text{By i.h.} \\
[\Omega, \alpha](\Delta, \alpha) \vdash [\Omega, \alpha]A \leq [\Omega, \alpha]B_0 \quad \text{By Lemma 45 (Substitution Stability)} \\
[\Omega, \alpha](\Delta, \alpha) \vdash [\Omega]A \leq [\Omega]B_0 \quad \text{By def. of substitution} \\
[\Omega]\Delta \vdash [\Omega]A \leq \forall \alpha. [\Omega]B_0 \quad \text{By } \leq \forall R \\
[\Omega]\Delta \vdash [\Omega]A \leq [\Omega](\forall \alpha. B_0) \quad \text{By def. of substitution}
\end{array}$$

$$\begin{array}{l}
\bullet \text{ Case } \hat{\alpha} \notin \text{FV}(B) \quad \Gamma \vdash \hat{\alpha} : \leq B \dashv \Delta \\
\frac{\Gamma \vdash \hat{\alpha} <: B \dashv \Delta}{\Gamma \vdash \hat{\alpha} : \leq B \dashv \Delta} \text{<:InstantiateL} \\
\Gamma \vdash \hat{\alpha} : \leq B \dashv \Delta \quad \text{Subderivation} \\
[\Omega]\Delta \vdash [\Omega]\hat{\alpha} \leq [\Omega]B \quad \text{By Theorem 10}
\end{array}$$

• **Case <:InstantiateR:** Similar to the case for <:InstantiateL. □

Corollary 53 (Soundness, Pretty Version). *If $\Psi \vdash A <: B \dashv \Delta$, then $\Psi \vdash A \leq B$.*

Proof. By reflexivity (Lemma 20 (Reflexivity)), $\Psi \longrightarrow \Psi$.

Since Ψ has no existential variables, it is a complete context Ω .

By Theorem 11, $[\Psi]\Psi \vdash [\Psi]A \leq [\Psi]B$.

Since Ψ has no existential variables, $[\Psi]\Psi = \Psi$, and $[\Psi]A = A$, and $[\Psi]B = B$.

Therefore $\Psi \vdash A \leq B$. □

I' Typing Extension

Lemma 54 (Typing Extension).

If $\Gamma \vdash e \leftarrow A \dashv \Delta$ or $\Gamma \vdash e \Rightarrow A \dashv \Delta$ or $\Gamma \vdash A \bullet e \Rightarrow C \dashv \Delta$ then $\Gamma \longrightarrow \Delta$.

Proof. By induction on the given derivation.

• **Cases Var, 1l, 1l \Rightarrow :**

Since $\Delta = \Gamma$, the result follows by Lemma 20 (Reflexivity).

$$\begin{array}{l}
\bullet \text{ Case } \frac{\Gamma \vdash e \Rightarrow B \dashv \Theta \quad \Theta \vdash [\Theta]B <: [\Theta]A \dashv \Delta}{\Gamma \vdash e \leftarrow A \dashv \Delta} \text{Sub} \\
\Gamma \longrightarrow \Theta \quad \text{By i.h.} \\
\Theta \longrightarrow \Delta \quad \text{By Lemma 33 (Subtyping Extension)} \\
\Rightarrow \Gamma \longrightarrow \Delta \quad \text{By Lemma 21 (Transitivity)}
\end{array}$$

$$\begin{array}{l}
\bullet \text{ Case } \frac{\Gamma \vdash A \quad \Gamma \vdash e \leftarrow A \dashv \Delta}{\Gamma \vdash (e : A) \Rightarrow A \dashv \Delta} \text{Anno} \\
\Rightarrow \Gamma \longrightarrow \Delta \quad \text{By i.h.}
\end{array}$$

$$\begin{array}{l}
\bullet \text{ Case } \frac{\Gamma, \alpha \vdash e \leftarrow A_0 \dashv \Delta, \alpha, \Theta}{\Gamma \vdash e \leftarrow \forall \alpha. A_0 \dashv \Delta} \forall I \\
\Gamma, \alpha \longrightarrow \Delta, \alpha, \Theta \quad \text{By i.h.} \\
\Rightarrow \Gamma \longrightarrow \Delta \quad \text{By Lemma 24 (Extension Order) (i)}
\end{array}$$

- **Case**
$$\frac{\Gamma, \hat{\alpha} \vdash [\hat{\alpha}/\alpha]A_0 \bullet e \Rightarrow C \dashv \Delta}{\Gamma \vdash \forall \alpha. A_0 \bullet e \Rightarrow C \dashv \Delta} \forall\text{App}$$

$$\Gamma, \hat{\alpha} \longrightarrow \Delta \quad \text{By i.h.}$$

$$\Gamma \longrightarrow \Gamma, \hat{\alpha} \quad \text{By } \longrightarrow\text{Add}$$

$$\Gamma \longrightarrow \Delta \quad \text{By Lemma 21 (Transitivity)}$$
- **Case**
$$\frac{\Gamma, x : A_1 \vdash e \Leftarrow A_2 \dashv \Delta, x : A_1, \Theta}{\Gamma \vdash \lambda x. e \Leftarrow A_1 \rightarrow A_2 \dashv \Delta} \rightarrow\text{I}$$

$$\Gamma, x : A_1 \longrightarrow \Delta, x : A_1, \Theta \quad \text{By i.h.}$$

$$\Gamma \longrightarrow \Delta \quad \text{By Lemma 24 (Extension Order) (v)}$$
- **Case**
$$\frac{\Gamma \vdash e_1 \Rightarrow B \dashv \Theta \quad \Theta \vdash [\Theta]B \bullet e_2 \Rightarrow A \dashv \Delta}{\Gamma \vdash e_1 e_2 \Rightarrow A \dashv \Delta} \rightarrow\text{E}$$

By the i.h. on each premise, then Lemma 21 (Transitivity).
- **Case**
$$\frac{\Gamma, \hat{\alpha}, \hat{\beta}, x : \hat{\alpha} \vdash e \Leftarrow \hat{\beta} \dashv \Delta, x : \hat{\alpha}, \Theta}{\Gamma \vdash \lambda x. e \Rightarrow \hat{\alpha} \rightarrow \hat{\beta} \dashv \Delta} \rightarrow\text{I} \Rightarrow$$

$$\Gamma, \hat{\alpha}, \hat{\beta}, x : \hat{\alpha} \longrightarrow \Delta, x : \hat{\alpha}, \Theta \quad \text{By i.h.}$$

$$\Gamma, \hat{\alpha}, \hat{\beta} \longrightarrow \Delta \quad \text{By Lemma 24 (Extension Order) (v)}$$

$$\Gamma \longrightarrow \Gamma, \hat{\alpha}, \hat{\beta} \quad \text{By } \longrightarrow\text{Add (twice)}$$

$$\Gamma \longrightarrow \Delta \quad \text{By Lemma 21 (Transitivity)}$$
- **Case**
$$\frac{\Gamma \vdash e \Leftarrow A \dashv \Delta}{\Gamma \vdash A \rightarrow C \bullet e \Rightarrow C \dashv \Delta} \rightarrow\text{App}$$

$$\Gamma \longrightarrow \Delta \quad \text{By i.h.}$$
- **Case**
$$\frac{\Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2] \vdash e \Leftarrow \hat{\alpha}_1 \dashv \Delta}{\Gamma[\hat{\alpha}] \vdash \hat{\alpha} \bullet e \Rightarrow \hat{\alpha}_2 \dashv \Delta} \hat{\alpha}\text{App}$$

$$\Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2] \longrightarrow \Delta \quad \text{By i.h.}$$

$$\Gamma[\hat{\alpha}] \longrightarrow \Gamma[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2] \quad \text{By Lemma 27 (Solved Variable Addition for Extension)}$$

$$\text{then Lemma 29 (Parallel Admissibility) (ii)}$$

$$\Gamma \longrightarrow \Delta \quad \text{By Lemma 21 (Transitivity)} \quad \square$$

J' Soundness of Typing

Theorem 12 (Soundness of Algorithmic Typing). *Given $\Delta \longrightarrow \Omega$:*

- (i) *If $\Gamma \vdash e \Leftarrow A \dashv \Delta$ then $[\Omega]\Delta \vdash e \Leftarrow [\Omega]A$.*
- (ii) *If $\Gamma \vdash e \Rightarrow A \dashv \Delta$ then $[\Omega]\Delta \vdash e \Rightarrow [\Omega]A$.*
- (iii) *If $\Gamma \vdash A \bullet e \Rightarrow C \dashv \Delta$ then $[\Omega]\Delta \vdash [\Omega]A \bullet e \Rightarrow [\Omega]C$.*

Proof. By induction on the given algorithmic typing derivation.

- **Case**
$$\frac{(x : A) \in \Gamma}{\Gamma \vdash x \Rightarrow A \dashv \Gamma} \text{Var}$$

- | | | |
|---|---|-------------------------------------|
| | $(x : A) \in \Gamma$ | Premise |
| | $(x : A) \in \Delta$ | By $\Gamma = \Delta$ |
| | $\Delta \longrightarrow \Omega$ | Given |
| | $(x : [\Omega]A) \in [\Omega]\Gamma$ | By Lemma 42 (Variable Preservation) |
| ☞ | $[\Omega]\Gamma \vdash x \Rightarrow [\Omega]A$ | By DeclVar |
- **Case** $\frac{\Gamma \vdash e \Rightarrow A \dashv \Theta \quad \Theta \vdash [\Theta]A <: [\Theta]B \dashv \Delta}{\Gamma \vdash e \Leftarrow B \dashv \Delta}$ Sub
- | | | |
|---|--|---|
| | $\Gamma \vdash e \Rightarrow A \dashv \Theta$ | Subderivation |
| | $\Theta \vdash [\Theta]A <: [\Theta]B \dashv \Delta$ | Subderivation |
| | $\Theta \longrightarrow \Delta$ | By Lemma 54 (Typing Extension) |
| | $\Delta \longrightarrow \Omega$ | Given |
| | $\Theta \longrightarrow \Omega$ | By Lemma 21 (Transitivity) |
| | $[\Omega]\Theta \vdash e \Rightarrow [\Omega]A$ | By i.h. |
| | $[\Omega]\Theta = [\Omega]\Delta$ | By Lemma 52 (Confluence of Completeness) |
| | $[\Omega]\Delta \vdash e \Rightarrow [\Omega]A$ | By above equalities |
| | $\Theta \vdash [\Theta]A <: [\Theta]B \dashv \Delta$ | Subderivation |
| | $[\Omega]\Delta \vdash [\Omega][\Theta]A \leq [\Omega][\Theta]B$ | By Theorem 11 |
| | $[\Omega][\Theta]A = [\Omega]A$ | By Lemma 18 (Substitution Extension Invariance) |
| | $[\Omega][\Theta]B = [\Omega]B$ | By Lemma 18 (Substitution Extension Invariance) |
| | $[\Omega]\Delta \vdash [\Omega]A \leq [\Omega]B$ | By above equalities |
| ☞ | $[\Omega]\Delta \vdash e \Leftarrow [\Omega]B$ | By DeclSub |
- **Case** $\frac{\Gamma \vdash A \quad \Gamma \vdash e_0 \Leftarrow A \dashv \Delta}{\Gamma \vdash (e_0 : A) \Rightarrow A \dashv \Delta}$ Anno
- | | | |
|---|---|--|
| | $\Gamma \vdash e_0 \Leftarrow A \dashv \Delta$ | Subderivation |
| | $[\Omega]\Delta \vdash e_0 \Leftarrow [\Omega]A$ | By i.h. |
| | $\Gamma \vdash A$ | Subderivation |
| | $\Gamma \longrightarrow \Delta$ | By Lemma 54 (Typing Extension) |
| | $\Delta \longrightarrow \Omega$ | Given |
| | $\Gamma \longrightarrow \Omega$ | By Lemma 21 (Transitivity) |
| | $\Omega \vdash A$ | By Lemma 25 (Extension Weakening) |
| | $[\Omega]\Omega \vdash [\Omega]A$ | By Lemma 44 (Substitution for Well-Formedness) |
| | $[\Omega]\Delta = [\Omega]\Omega$ | By Lemma 49 (Stability of Complete Contexts) |
| | $[\Omega]\Delta \vdash [\Omega]A$ | By above equality |
| | $[\Omega]\Delta \vdash (e_0 : [\Omega]A) \Rightarrow [\Omega]A$ | By DeclAnno |
| | A contains no existential variables | Assumption about source programs |
| | $[\Omega]A = A$ | From definition of substitution |
| ☞ | $[\Omega]\Delta \vdash (e_0 : A) \Rightarrow [\Omega]A$ | By above equality |
- **Case**
- | | | |
|---|---|-------------------------------|
| | $\frac{}{\Gamma \vdash () \Leftarrow 1 \dashv \underbrace{\Gamma}_{\Delta}}$ 1l | |
| | $[\Omega]\Delta \vdash () \Leftarrow 1$ | By Decl1l |
| ☞ | $[\Omega]\Delta \vdash () \Leftarrow [\Omega]1$ | By definition of substitution |
- **Case** $\frac{\Gamma, x : A_1 \vdash e_0 \Leftarrow A_2 \dashv \Delta, x : A_1, \Theta}{\Gamma \vdash \lambda x. e \Leftarrow A_1 \rightarrow A_2 \dashv \Delta}$ \rightarrow l

$\Delta \longrightarrow \Omega$	Given
$\Delta, \alpha \longrightarrow \Omega, \alpha$	By \longrightarrow Uvar
$\Gamma, \alpha \longrightarrow \Delta, \alpha, \Theta$	By Lemma 54 (Typing Extension)
Θ is soft	By Lemma 24 (Extension Order) (i) (with $\Gamma_R = \cdot$, which is soft)
$\underbrace{\Delta, \alpha, \Theta}_{\Delta'} \longrightarrow \underbrace{\Omega, \alpha, \Theta }_{\Omega'}$	By Lemma 48 (Filling Completes)
$\Gamma, \alpha \vdash e \Leftarrow A_0 \dashv \Delta'$	Subderivation
$[\Omega']\Delta' \vdash e \Leftarrow [\Omega']A_0$	By i.h.
$[\Omega']A_0 = [\Omega]A_0$	By Lemma 45 (Substitution Stability)
$[\Omega']\Delta' \vdash e \Leftarrow [\Omega]A_0$	By above equality
$\underbrace{\Delta, \alpha, \Theta}_{\Delta'} \longrightarrow \underbrace{\Omega, \alpha, \Theta }_{\Omega'}$	Above
Θ is soft	Above
$[\Omega']\Delta' = [\Omega]\Delta, \alpha$	By Lemma 47 (Softness Goes Away)
$[\Omega]\Delta, \alpha \vdash e \Leftarrow [\Omega]A_0$	By above equality
$[\Omega]\Delta \vdash e \Leftarrow \forall \alpha. [\Omega]A_0$	By Decl \forall I
☞ $[\Omega]\Delta \vdash e \Leftarrow [\Omega](\forall \alpha. A_0)$	By definition of substitution

• **Case** $\frac{\Gamma, \hat{\alpha} \vdash [\hat{\alpha}/\alpha]A_0 \bullet e \Rightarrow C \dashv \Delta}{\Gamma \vdash \forall \alpha. A_0 \bullet e \Rightarrow C \dashv \Delta} \forall\text{App}$

$\Gamma, \hat{\alpha} \vdash [\hat{\alpha}/\alpha]A_0 \bullet e \Rightarrow C \dashv \Delta$	Subderivation
$\Delta \longrightarrow \Omega$	Given
$[\Omega]\Delta \vdash [\Omega][\hat{\alpha}/\alpha]A_0 \bullet e \Rightarrow [\Omega]C$	By i.h.
$[\Omega]\Delta \vdash [[\Omega]\hat{\alpha}/\alpha] [\Omega]A_0 \bullet e \Rightarrow [\Omega]C$	By distributivity of substitution
$\Gamma, \hat{\alpha} \longrightarrow \Delta$	By Lemma 54 (Typing Extension)
$\Gamma, \hat{\alpha} \longrightarrow \Omega$	By Lemma 21 (Transitivity)
$\Gamma, \hat{\alpha} \vdash \hat{\alpha}$	By EvarWF
$\Omega \vdash \hat{\alpha}$	By Lemma 25 (Extension Weakening)
$[\Omega]\Omega \vdash [\Omega]\hat{\alpha}$	By Lemma 44 (Substitution for Well-Formedness)
$[\Omega]\Omega = [\Omega]\Delta$	By Lemma 49 (Stability of Complete Contexts)
$[\Omega]\Delta \vdash [\Omega]\hat{\alpha}$	By above equality
$[\Omega]\Delta \vdash \forall \alpha. [\Omega]A_0 \bullet e \Rightarrow [\Omega]C$	By Decl \forall App
☞ $[\Omega]\Delta \vdash [\Omega](\forall \alpha. A_0) \bullet e \Rightarrow [\Omega]C$	By definition of substitution

• **Case** $\frac{\Gamma \vdash e \Leftarrow B \dashv \Delta}{\Gamma \vdash B \rightarrow C \bullet e \Rightarrow C \dashv \Delta} \rightarrow\text{App}$

$\Gamma \vdash e \Leftarrow B \dashv \Delta$	Subderivation
$\Delta \longrightarrow \Omega$	Given
$[\Omega]\Delta \vdash e \Leftarrow [\Omega]B$	By i.h.
$[\Omega]\Delta \vdash ([\Omega]B) \rightarrow ([\Omega]C) \bullet e \Rightarrow [\Omega]C$	By Decl \rightarrow App
☞ $[\Omega]\Delta \vdash [\Omega](B \rightarrow C) \bullet e \Rightarrow [\Omega]C$	By definition of substitution

• **Case** $\frac{\Gamma_0[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2] \vdash e \Leftarrow \hat{\alpha}_1 \dashv \Delta}{\underbrace{\Gamma_0[\hat{\alpha}]}_{\Gamma} \vdash \hat{\alpha} \bullet e \Rightarrow \hat{\alpha}_2 \dashv \Delta} \hat{\alpha}\text{App}$

$$\begin{array}{l}
\overbrace{\Gamma_0[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2]}^{\Gamma'} \vdash e \Leftarrow \hat{\alpha}_1 \dashv \Delta \quad \text{Subderivation} \\
\Delta \longrightarrow \Omega \quad \text{Given} \\
[\Omega]\Delta \vdash e \Leftarrow [\Omega]\hat{\alpha}_1 \quad \text{By i.h.} \\
[\Omega]\Delta \vdash ([\Omega]\hat{\alpha}_1) \rightarrow ([\Omega]\hat{\alpha}_2) \bullet e \Rightarrow [\Omega]\hat{\alpha}_2 \quad \text{By Decl} \rightarrow \text{App} \\
\Gamma' \longrightarrow \Delta \quad \text{By Lemma 54 (Typing Extension)} \\
\Delta \longrightarrow \Omega \quad \text{Given} \\
\Gamma' \longrightarrow \Omega \quad \text{By Lemma 21 (Transitivity)} \\
[\Gamma']\hat{\alpha} = [\Gamma'](\hat{\alpha}_1 \rightarrow \hat{\alpha}_2) \quad \text{By definition of } [\Gamma'](-) \\
[\Omega][\Gamma']\hat{\alpha} = [\Omega][\Gamma'](\hat{\alpha}_1 \rightarrow \hat{\alpha}_2) \quad \text{Applying } \Omega \text{ to both sides} \\
[\Omega]\hat{\alpha} = [\Omega](\hat{\alpha}_1 \rightarrow \hat{\alpha}_2) \quad \text{By Lemma 18 (Substitution Extension Invariance), twice} \\
= ([\Omega]\hat{\alpha}_1) \rightarrow ([\Omega]\hat{\alpha}_2) \quad \text{By definition of substitution} \\
\text{☞} \quad [\Omega]\Delta \vdash [\Omega]\hat{\alpha} \bullet e \Rightarrow [\Omega]\hat{\alpha}_2 \quad \text{By above equality}
\end{array}$$

• **Case**

$$\begin{array}{l}
\overline{\Gamma \vdash () \Rightarrow 1 \dashv \underbrace{\Gamma}_{\Delta}} \quad \text{11} \Rightarrow \\
\text{☞} \quad [\Omega]\Delta \vdash () \Rightarrow [\Omega]1 \quad \text{By Decl11} \Rightarrow \text{ and definition of substitution}
\end{array}$$

• **Case**

$$\begin{array}{l}
\frac{\Gamma, \hat{\alpha}, \hat{\beta}, x : \hat{\alpha} \vdash e_0 \Leftarrow \hat{\beta} \dashv \Delta, x : \hat{\alpha}, \Theta}{\Gamma \vdash \lambda x. e_0 \Rightarrow \hat{\alpha} \rightarrow \hat{\beta} \dashv \Delta} \rightarrow \text{1} \Rightarrow \\
\Gamma, \hat{\alpha}, \hat{\beta}, x : \hat{\alpha} \longrightarrow \Delta, x : \hat{\alpha}, \Theta \quad \text{By Lemma 54 (Typing Extension)} \\
\Theta \text{ is soft} \quad \text{By Lemma 24 (Extension Order) (v) (with } \Gamma_R = \cdot, \text{ which is soft)} \\
\Gamma, \hat{\alpha}, \hat{\beta} \longrightarrow \Delta \quad \text{"} \\
\Delta \longrightarrow \Omega \quad \text{Given} \\
\Delta, x : \hat{\alpha} \longrightarrow \Omega, x : [\Omega]\hat{\alpha} \quad \text{By } \longrightarrow \text{Var} \\
\overbrace{\Delta, x : \hat{\alpha}, \Theta}^{\Delta'} \longrightarrow \underbrace{\Omega, x : [\Omega]\hat{\alpha}, |\Theta|}_{\Omega'} \quad \text{By Lemma 48 (Filling Completes)} \\
\Gamma, \hat{\alpha}, \hat{\beta}, x : \hat{\alpha} \vdash e \Leftarrow \hat{\beta} \dashv \Delta, x : \hat{\alpha}, \Theta \quad \text{Subderivation} \\
[\Omega']\Delta' \vdash e_0 \Leftarrow [\Omega']\hat{\beta} \quad \text{By i.h.} \\
[\Omega']\hat{\beta} = [\Omega, x : [\Omega]\hat{\alpha}]\hat{\beta} \quad \text{By Lemma 45 (Substitution Stability)} \\
= [\Omega]\hat{\beta} \quad \text{By definition of substitution} \\
[\Omega']\Delta' = [\Omega, x : [\Omega]\hat{\alpha}](\Delta, x : \hat{\alpha}) \quad \text{By Lemma 47 (Softness Goes Away)} \\
= [\Omega]\Delta, x : [\Omega]\hat{\alpha} \quad \text{By definition of context substitution} \\
[\Omega]\Delta, x : [\Omega]\hat{\alpha} \vdash e_0 \Leftarrow [\Omega]\hat{\beta} \quad \text{By above equalities} \\
\Gamma, \hat{\alpha}, \hat{\beta} \longrightarrow \Delta \quad \text{Above} \\
\Gamma, \hat{\alpha}, \hat{\beta} \longrightarrow \Omega \quad \text{By Lemma 21 (Transitivity)} \\
\Gamma, \hat{\alpha}, \hat{\beta} \vdash \hat{\alpha} \quad \text{By EvarWF} \\
\Omega \vdash \hat{\alpha} \quad \text{By Lemma 25 (Extension Weakening)} \\
[\Omega]\Delta \vdash [\Omega]\hat{\alpha} \quad \text{By Lemma 44 (Substitution for Well-Formedness)} \\
\quad \text{and Lemma 49 (Stability of Complete Contexts)} \\
[\Omega]\Delta \vdash [\Omega]\hat{\beta} \quad \text{By similar reasoning} \\
[\Omega]\Delta \vdash ([\Omega]\hat{\alpha}) \rightarrow ([\Omega]\hat{\beta}) \quad \text{By DeclArrowWF} \\
[\Omega]\hat{\alpha}, [\Omega]\hat{\beta} \text{ monotypes} \quad \Omega \text{ predicative} \\
[\Omega]\Delta \vdash \lambda x. e_0 \Rightarrow ([\Omega]\hat{\alpha}) \rightarrow ([\Omega]\hat{\beta}) \quad \text{By Decl} \rightarrow \text{1} \Rightarrow \\
\text{☞} \quad [\Omega]\Delta \vdash \lambda x. e_0 \Rightarrow [\Omega](\hat{\alpha} \rightarrow \hat{\beta}) \quad \text{By definition of substitution} \quad \square
\end{array}$$

K' Completeness

K'.1 Instantiation Completeness

Theorem 13 (Instantiation Completeness).

Given $\Gamma \longrightarrow \Omega$ and $A = [\Gamma]A$ and $\hat{\alpha} \in \text{unsolved}(\Gamma)$ and $\hat{\alpha} \notin \text{FV}(A)$:

- (1) If $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq [\Omega]A$
then there are Δ, Ω' such that $\Omega \longrightarrow \Omega'$ and $\Delta \longrightarrow \Omega'$ and $\Gamma \vdash \hat{\alpha} : \leq A \dashv \Delta$.
- (2) If $[\Omega]\Gamma \vdash [\Omega]A \leq [\Omega]\hat{\alpha}$
then there are Δ, Ω' such that $\Omega \longrightarrow \Omega'$ and $\Delta \longrightarrow \Omega'$ and $\Gamma \vdash A : \leq \hat{\alpha} \dashv \Delta$.

Proof. By mutual induction on the given declarative subtyping derivation.

(1) We have $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq [\Omega]A$. We now case-analyze the shape of A .

- **Case** $A = \hat{\beta}$:

It is given that $\hat{\alpha} \notin \text{FV}(\hat{\beta})$, so $\hat{\alpha} \neq \hat{\beta}$.

Since $A = \hat{\beta}$, we have $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq [\Omega]\hat{\beta}$.

Since Ω is predicative, $[\Omega]\hat{\alpha} = \tau_1$ and $[\Omega]\hat{\beta} = \tau_2$, so we have $[\Omega]\Gamma \vdash \tau_1 \leq \tau_2$.

By Lemma 9 (Monotype Equality), $\tau_1 = \tau_2$.

We have $A = \hat{\beta}$ and $[\Gamma]A = A$, so $[\Gamma]\hat{\beta} = \hat{\beta}$. Thus $\hat{\beta} \in \text{unsolved}(\Gamma)$.

Let Ω' be Ω . By Lemma 20 (Reflexivity), $\Omega \longrightarrow \Omega$.

Now consider whether $\hat{\alpha}$ is declared to the left of $\hat{\beta}$, or vice versa.

- **Case** $\Gamma = (\Gamma_0, \hat{\alpha}, \Gamma_1, \hat{\beta}, \Gamma_2)$:

Let Δ be $\Gamma_0, \hat{\alpha}, \Gamma_1, \hat{\beta} = \hat{\alpha}, \Gamma_2$.

By rule `InstLReach`, $\Gamma \vdash \hat{\alpha} : \leq \hat{\beta} \dashv \Delta$.

It remains to show that $\Delta \longrightarrow \Omega$.

We have $[\Omega]\hat{\alpha} = [\Omega]\hat{\beta}$. Then by Lemma 30 (Parallel Extension Solution), $\Delta \longrightarrow \Omega$.

- **Case** $(\Gamma = \Gamma_0, \hat{\beta}, \Gamma_1, \hat{\alpha}, \Gamma_2)$:

Let Δ be $\Gamma_0, \hat{\beta}, \Gamma_1, \hat{\alpha} = \hat{\beta}, \Gamma_2$.

By rule `InstLSolve`, $\Gamma \vdash \hat{\alpha} : \leq \hat{\beta} \dashv \Delta$.

It remains to show that $\Delta \longrightarrow \Omega$.

We have $[\Omega]\hat{\beta} = [\Omega]\hat{\alpha}$. Then by Lemma 30 (Parallel Extension Solution), $\Delta \longrightarrow \Omega$.

- **Case** $A = \alpha$:

Since $A = \alpha$, we have $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq [\Omega]\alpha$.

Since $[\Omega]\alpha = \alpha$, we have $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq \alpha$.

By inversion, `≤Var` was used, so $[\Omega]\hat{\alpha} = \alpha$; therefore, since Ω is well-formed, α is declared to the left of $\hat{\alpha}$ in Ω .

We have $\Gamma \longrightarrow \Omega$.

By Lemma 17 (Reverse Declaration Order Preservation), we know that α is declared to the left of $\hat{\alpha}$ in Γ ; that is, $\Gamma = \Gamma_0[\alpha][\hat{\alpha}]$.

Let $\Delta = \Gamma_0[\alpha][\hat{\alpha} = \alpha]$ and $\Omega' = \Omega$.

By `InstLSolve`, $\Gamma_0[\alpha][\hat{\alpha}] \vdash \hat{\alpha} : \leq \alpha \dashv \Delta$.

By Lemma 30 (Parallel Extension Solution), $\Gamma_0[\alpha][\hat{\alpha} = \alpha] \longrightarrow \Omega$.

- **Case** $A = A_1 \rightarrow A_2$:

By the definition of substitution, $[\Omega]A = ([\Omega]A_1) \rightarrow ([\Omega]A_2)$.

Therefore $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq ([\Omega]A_1) \rightarrow ([\Omega]A_2)$.

Since we have an arrow as the supertype, only `≤∀L` or `≤→` could have been used, and the subtype $[\Omega]\hat{\alpha}$ must be either a quantifier or an arrow. But Ω is predicative, so $[\Omega]\hat{\alpha}$ cannot be a quantifier. Therefore, it is an arrow: $[\Omega]\hat{\alpha} = \tau_1 \rightarrow \tau_2$, and `≤→` concluded the derivation.

Inverting `≤→` gives $[\Omega]\Gamma \vdash [\Omega]A_2 \leq \tau_2$ and $[\Omega]\Gamma \vdash \tau_1 \leq [\Omega]A_1$.

Since $\hat{\alpha} \in \text{unsolved}(\Gamma)$, we know that Γ has the form $\Gamma_0[\hat{\alpha}]$.

By Lemma 28 (Unsolved Variable Addition for Extension) twice, inserting unsolved variables

$\hat{\alpha}_2$ and $\hat{\alpha}_1$ into the middle of the context extends it, that is: $\Gamma_0[\hat{\alpha}] \longrightarrow \Gamma_0[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha}]$.

Clearly, $\hat{\alpha}_1 \rightarrow \hat{\alpha}_2$ is well-formed in $(\dots, \hat{\alpha}_2, \hat{\alpha}_1)$, so by Lemma 26 (Solution Admissibility for Extension), solving $\hat{\alpha}$ extends the context: $\Gamma_0[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha}] \longrightarrow \Gamma_0[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2]$. Then by Lemma 21 (Transitivity), $\Gamma_0[\hat{\alpha}] \longrightarrow \Gamma_0[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2]$.

Since $\hat{\alpha} \in \text{unsolved}(\Gamma)$ and $\Gamma \longrightarrow \Omega$, we know that Ω has the form $\Omega_0[\hat{\alpha} = \tau_0]$. To show that we can extend this context, we apply Lemma 27 (Solved Variable Addition for Extension) twice to introduce $\hat{\alpha}_2 = \tau_2$ and $\hat{\alpha}_1 = \tau_1$, and then Lemma 26 (Solution Admissibility for Extension) to overwrite τ_0 :

$$\underbrace{\Omega_0[\hat{\alpha} = \tau_0]}_{\Omega} \longrightarrow \Omega_0[\hat{\alpha}_2 = \tau_2, \hat{\alpha}_1 = \tau_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2]$$

We have $\Gamma \longrightarrow \Omega$, that is,

$$\Gamma_0[\hat{\alpha}] \longrightarrow \Omega_0[\hat{\alpha} = \tau_0]$$

By Lemma 29 (Parallel Admissibility) (i) twice, inserting unsolved variables $\hat{\alpha}_2$ and $\hat{\alpha}_1$ on both contexts in the above extension preserves extension:

$$\underbrace{\Gamma_0[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha}]}_{\Gamma_1} \longrightarrow \underbrace{\Omega_0[\hat{\alpha}_2 = \tau_2, \hat{\alpha}_1 = \tau_1, \hat{\alpha} = \tau_0]}_{\Omega_1} \quad \begin{array}{l} \text{By Lemma 29 (Parallel Admissibility) (ii) twice} \\ \text{By Lemma 31 (Parallel Variable Update)} \end{array}$$

Since $\hat{\alpha} \notin \text{FV}(A)$, it follows that $[\Gamma_1]A = [\Gamma]A = A$.

Therefore $\hat{\alpha}_1 \notin \text{FV}(A_1)$ and $\hat{\alpha}_1, \hat{\alpha}_2 \notin \text{FV}(A_2)$.

By Lemma 51 (Finishing Completions) and Lemma 50 (Finishing Types), $[\Omega_1]\Gamma_1 = [\Omega]\Gamma$ and $[\Omega_1]\hat{\alpha}_1 = \tau_1$.

By i.h., there are Δ_2 and Ω_2 such that $\Gamma_1 \vdash A_1 \preceq \hat{\alpha}_1 \dashv \Delta_2$ and $\Delta_2 \longrightarrow \Omega_2$ and $\Omega_1 \longrightarrow \Omega_2$.

Next, note that $[\Delta_2][\Delta_2]A_2 = [\Delta_2]A_2$.

By Lemma 34 (Left Unsolvedness Preservation), we know that $\hat{\alpha}_2 \in \text{unsolved}(\Delta_2)$.

By Lemma 35 (Left Free Variable Preservation), we know that $\hat{\alpha}_2 \notin \text{FV}([\Delta_2]A_2)$.

By Lemma 21 (Transitivity), $\Omega \longrightarrow \Omega_2$.

We know $[\Omega_2]\Delta_2 = [\Omega]\Gamma$ because:

$$\begin{aligned} [\Omega_2]\Delta_2 &= [\Omega_2]\Omega_2 && \text{By Lemma 49 (Stability of Complete Contexts)} \\ &= [\Omega]\Omega && \text{By Lemma 51 (Finishing Completions)} \\ &= [\Omega]\Gamma && \text{By Lemma 49 (Stability of Complete Contexts)} \end{aligned}$$

By Lemma 50 (Finishing Types), we know that $[\Omega_2]\hat{\alpha}_2 = [\Omega_1]\hat{\alpha}_2 = \tau_2$.

By Lemma 50 (Finishing Types), we know that $[\Omega_2]A_2 = [\Omega]A_2$.

Hence we know that $[\Omega_2]\Delta_2 \vdash [\Omega_2]\hat{\alpha}_2 \leq [\Omega_2]A_2$.

By i.h., we have Δ and Ω' such that $\Delta_2 \vdash \hat{\alpha}_2 \preceq [\Delta_2]A_2 \dashv \Delta$ and $\Omega_2 \longrightarrow \Omega'$ and $\Delta \longrightarrow \Omega'$.

By rule InstLArr , $\Gamma \vdash \hat{\alpha} \preceq A \dashv \Delta$.

By Lemma 21 (Transitivity), $\Omega \longrightarrow \Omega'$.

- **Case $A = 1$:**

We have $A = 1$, so $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq [\Omega]1$.

Since $[\Omega]1 = 1$, we have $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq 1$.

The only declarative subtyping rules that can have 1 as the supertype in the conclusion are $\leq \forall L$ and $\leq \text{Unit}$. However, since Ω is predicative, $[\Omega]\hat{\alpha}$ cannot be a quantifier, so $\leq \forall L$ cannot have been used. Hence $\leq \text{Unit}$ was used and $[\Omega]\hat{\alpha} = 1$.

Let $\Delta = \Gamma[\hat{\alpha} = 1]$ and $\Omega' = \Omega$.

By InstLSolve , $\Gamma[\hat{\alpha}] \vdash \hat{\alpha} \preceq 1 \dashv \Delta$.

By Lemma 30 (Parallel Extension Solution), $\Gamma[\hat{\alpha} = 1] \longrightarrow \Omega$.

- **Case $A = \forall \beta. B$:**

We have $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq [\Omega](\forall \beta. B)$.

By definition of substitution, $[\Omega](\forall \beta. B) = \forall \beta. [\Omega]B$, so we have $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq \forall \beta. [\Omega]B$.

The only declarative subtyping rules that can have a quantifier as supertype are $\leq \forall L$ and $\leq \forall R$. However, since Ω is predicative, $[\Omega]\hat{\alpha}$ cannot be a quantifier, so $\leq \forall L$ cannot have been used.

Hence $\leq \forall R$ was used, and we have a subderivation of $[\Omega]\Gamma, \beta \vdash [\Omega]\hat{\alpha} \leq [\Omega]B$.

Let $\Omega_1 = (\Omega, \beta)$ and $\Gamma_1 = (\Gamma, \beta)$.

By $\longrightarrow_{\text{Uvar}}$, $\Gamma_1 \longrightarrow \Omega_1$.

By the definition of substitution, $[\Omega_1]B = [\Omega]B$ and $[\Omega_1]\hat{\alpha} = [\Omega]\hat{\alpha}$.

Note that $[\Omega_1]\Gamma_1 = [\Omega]\Gamma, \beta$.

Since $\hat{\alpha} \in \text{unsolved}(\Gamma)$, we have $\hat{\alpha} \in \text{unsolved}(\Gamma_1)$.

Since $\hat{\alpha} \notin \text{FV}(A)$ and $A = \forall\beta. B$, we have $\hat{\alpha} \notin \text{FV}(B)$.

By i.h., there are Ω_2 and Δ_2 such that $\Gamma, \beta \vdash \hat{\alpha} : \leq B \dashv \Delta_2$ and $\Delta_2 \longrightarrow \Omega_2$ and $\Omega_1 \longrightarrow \Omega_2$.

By Lemma 32 (Instantiation Extension), $\Gamma_1 \longrightarrow \Delta_2$, that is, $\Gamma, \beta \longrightarrow \Delta_2$.

Therefore by Lemma 24 (Extension Order), $\Delta_2 = (\Delta', \beta, \Omega'')$ where $\Gamma \longrightarrow \Delta'$.

By equality, we know $\Delta', \beta, \Omega'' \longrightarrow \Omega_2$.

By Lemma 24 (Extension Order), $\Omega_2 = (\Omega', \beta, \Omega'')$ where $\dashv \Delta' \longrightarrow \Omega'$.

We have $\Omega_1 \longrightarrow \Omega_2$, that is, $\Omega, \beta \longrightarrow \Omega', \beta, \Omega''$, so Lemma 24 (Extension Order) gives $\dashv \Omega \longrightarrow \Omega'$.

By rule InstLAllR , $\Gamma \vdash \hat{\alpha} : \leq \forall\beta. B \dashv \Delta'$.

(2) $[\Omega]\Gamma \vdash [\Omega]A \leq [\Omega]\hat{\alpha}$

These cases are mostly symmetric. The one exception is the one connective that is not treated symmetrically in the declarative subtyping rules:

- **Case $A = \forall\alpha. B$:**

Since $A = \forall\alpha. B$, we have $[\Omega]\Gamma \vdash [\Omega]\forall\beta. B \leq [\Omega]\hat{\alpha}$.

By symmetric reasoning to the previous case (the last case of part (1) above), $\leq\forall\text{L}$ must have been used, with a subderivation of $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq [\tau/\beta][\Omega]B$.

Since $[\Omega]\Gamma \vdash \tau$, the type τ has no existential variables and is therefore invariant under substitution: $\tau = [\Omega]\tau$. Therefore $[\tau/\beta][\Omega]B = [[\Omega]\tau/\beta][\Omega]B$.

By distributivity of substitution, this is $[\Omega][\tau/\beta]B$. Interposing $\hat{\beta}$, this is equal to $[\Omega][\tau/\hat{\beta}][\hat{\beta}/\beta]B$. Therefore $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq [\Omega][\tau/\hat{\beta}][\hat{\beta}/\beta]B$.

Let Ω_1 be $\Omega, \blacktriangleright_{\hat{\beta}}, \hat{\beta} = \tau$ and let Γ_1 be $\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta}$.

- By the definition of context application, $[\Omega_1]\Gamma_1 = [\Omega]\Gamma$.

- From the definition of substitution, $[\Omega_1]\hat{\alpha} = [\Omega]\hat{\alpha}$.

- It follows from the definition of substitution that $[\Omega][\tau/\hat{\beta}]C = [\Omega_1]C$ for all C . Therefore $[\Omega][\tau/\hat{\beta}][\hat{\beta}/\beta]B = [\Omega_1][\hat{\beta}/\beta]B$.

Applying these three equalities, $[\Omega_1]\Gamma_1 \vdash [\Omega_1]\hat{\alpha} \leq [\Omega_1][\hat{\beta}/\beta]B$.

By the definition of substitution, $[\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta}]B = [\Gamma]B = B$, so $\hat{\alpha} \notin \text{FV}([\Gamma_1]B)$.

Since $\hat{\alpha} \in \text{unsolved}(\Gamma)$, we have $\hat{\alpha} \in \text{unsolved}(\Gamma_1)$.

By i.h., there exist Δ_2 and Ω_2 such that $\Gamma_1 \vdash B \leq \hat{\alpha} \dashv \Delta_2$ and $\Omega_1 \longrightarrow \Omega_2$ and $\Delta_2 \longrightarrow \Omega_2$.

By Lemma 32 (Instantiation Extension), $\Gamma_1 \longrightarrow \Delta_2$, which is, $\Gamma, \blacktriangleright_{\hat{\beta}}, \hat{\beta} \longrightarrow \Delta_2$.

By Lemma 24 (Extension Order), $\Delta_2 = (\Delta', \blacktriangleright_{\hat{\beta}}, \Omega'')$ and $\Gamma \longrightarrow \Delta'$.

By equality, $\Delta', \blacktriangleright_{\hat{\beta}}, \Omega'' \longrightarrow \Omega_2$.

By Lemma 24 (Extension Order), $\Omega_2 = (\Omega', \blacktriangleright_{\hat{\beta}}, \Omega'')$ and $\dashv \Delta' \longrightarrow \Omega'$.

By equality, $\Omega, \blacktriangleright_{\hat{\beta}}, \hat{\beta} = \tau \longrightarrow \Omega', \blacktriangleright_{\hat{\beta}}, \Omega''$.

\dashv By Lemma 24 (Extension Order), $\Omega \longrightarrow \Omega'$.

By InstRAllL , $\Gamma \vdash \forall\beta. B \leq \hat{\alpha} \dashv \Delta'$. □

K'.2 Completeness of Subtyping

Theorem 14 (Generalized Completeness of Subtyping). *If $\Gamma \longrightarrow \Omega$ and $\Gamma \vdash A$ and $\Gamma \vdash B$ and $[\Omega]\Gamma \vdash [\Omega]A \leq [\Omega]B$ then there exist Δ and Ω' such that $\Delta \longrightarrow \Omega'$ and $\Omega \longrightarrow \Omega'$ and $\Gamma \vdash [\Gamma]A <: [\Gamma]B \dashv \Delta$.*

Proof. By induction on the derivation of $[\Omega]\Gamma \vdash [\Omega]A \leq [\Omega]B$.

We distinguish cases of $[\Gamma]B$ and $[\Gamma]A$ that are *impossible*, *fully written out*, and *similar to fully-written-out cases*.

		$[\Gamma]B$				
		$\forall\beta. B'$	1	α	$\hat{\beta}$	$B_1 \rightarrow B_2$
$\forall\alpha. A'$		1 (B poly)	2.Poly	2.Poly	2.Poly	2.Poly
	1	1 (B poly)	2.Units	<i>impossible</i>	2.BEx.Unit	<i>impossible</i>
$[\Gamma]A$	α	1 (B poly)	<i>impossible</i>	2.Uvars	2.BEx.Uvar	<i>impossible</i>
	$\hat{\alpha}$	1 (B poly)	2.AEx.Unit	2.AEx.Uvar	2.AEx.SameEx 2.AEx.OtherEx	2.AEx.Arrow
	$A_1 \rightarrow A_2$	1 (B poly)	<i>impossible</i>	<i>impossible</i>	2.BEx.Arrow	2.Arrows

The impossibility of the “*impossible*” entries follows from inspection of the declarative subtyping rules.

We first split on $[\Gamma]B$.

- **Case 1 (B poly):** $[\Gamma]B$ **polymorphic:** $[\Gamma]B = \forall\beta. B'$:

	$B = \forall\beta. B_0$	Γ predicative
	$B' = [\Gamma]B_0$	Γ predicative
	$[\Omega]B = [\Omega](\forall\beta. B_0)$	Applying Ω to both sides
	$= \forall\beta. [\Omega]B_0$	By definition of substitution
$\mathcal{D} ::$	$[\Omega]\Gamma \vdash [\Omega]A \leq [\Omega]B$	Given
$\mathcal{D} ::$	$[\Omega]\Gamma \vdash [\Omega]A \leq \forall\beta. [\Omega]B_0$	By above equality
$\mathcal{D}' ::$	$[\Omega]\Gamma, \beta \vdash [\Omega]A \leq [\Omega]B_0$	By Lemma 7 (Invertibility)
	$\mathcal{D}' < \mathcal{D}$	"
$\mathcal{D}' ::$	$[\Omega, \beta](\Gamma, \beta) \vdash [\Omega, \beta]A \leq [\Omega, \beta]B_0$	By definitions of substitution
	$\Gamma, \beta \vdash [\Gamma, \beta]A <: [\Gamma, \beta]B_0 \dashv \Delta'$	By i.h.
	$\Delta' \longrightarrow \Omega'_0$	"
	$\Omega, \beta \longrightarrow \Omega'_0$	"
	$\Gamma, \beta \vdash [\Gamma]A <: [\Gamma]B_0 \dashv \Delta'$	By definition of substitution
	$\Gamma, \beta \longrightarrow \Delta'$	By Lemma 32 (Instantiation Extension)
	$\Delta' = \Delta, \beta, \Theta$	By Lemma 24 (Extension Order) (i)
	$\Gamma \longrightarrow \Delta$	"
	$\Delta, \beta, \Theta \longrightarrow \Omega'_0$	By $\Delta' \longrightarrow \Omega'_0$ and above equality
	$\Omega'_0 = \Omega', \beta, \Omega_R$	By Lemma 24 (Extension Order) (i)
\dashv	$\Delta \longrightarrow \Omega'$	"
	$\Gamma, \beta \vdash [\Gamma]A <: [\Gamma]B_0 \dashv \Delta, \beta, \Theta$	By above equality
	$\Omega, \beta \longrightarrow \Omega', \beta, \Omega_R$	By above equality
\dashv	$\Omega \longrightarrow \Omega'$	By Lemma 21 (Transitivity)
	$\Gamma \vdash [\Gamma]A <: \forall\beta. [\Gamma]B_0 \dashv \Delta$	By $<: \forall R$
\dashv	$\Gamma \vdash [\Gamma]A <: \forall\beta. B' \dashv \Delta$	By above equality

• **Cases 2.*: $[\Gamma]B$ not polymorphic:**

We split on the form of $[\Gamma]A$.

– **Case 2.Poly: $[\Gamma]A$ is polymorphic: $[\Gamma]A = \forall\alpha. A'$:**

$A = \forall\alpha. A_0$	Γ predicative	
$A' = [\Gamma]A_0$	Γ predicative	
$[\Omega]A = [\Omega](\forall\alpha. A_0)$	Applying Ω to both sides	
$[\Omega]A = \forall\alpha. [\Omega]A_0$	By definition of substitution	
$[\Omega]\Gamma \vdash [\Omega]A \leq [\Omega]B$	Given	
$[\Omega]\Gamma \vdash \forall\alpha. [\Omega]A_0 \leq [\Omega]B$	By above equality	
$[\Gamma]B \neq (\forall\beta. \dots)$	We are in the “ $[\Gamma]B$ not polymorphic” subcase	
$B \neq (\forall\beta. \dots)$	Γ predicative	
$[\Omega]\Gamma \vdash [\tau/\alpha][\Omega]A_0 \leq [\Omega]B$	By inversion on $\leq \forall L$	
$[\Omega]\Gamma \vdash \tau$	"	
$\Gamma \longrightarrow \Omega$	Given	
$\Gamma, \blacktriangleright_{\hat{\alpha}} \longrightarrow \Omega, \blacktriangleright_{\hat{\alpha}}$	By \longrightarrow Marker	
$\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha} \longrightarrow \underbrace{\Omega, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha}}_{\Omega_0} = \tau$	By \longrightarrow Solve	
$[\Omega]\Gamma = [\Omega_0](\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha})$	By definition of context application (lines 16, 13)	
$[\Omega]\Gamma \vdash [\tau/\alpha][\Omega]A_0 \leq [\Omega]B$	Above	
$[\Omega_0](\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha}) \vdash [\tau/\alpha][\Omega]A_0 \leq [\Omega]B$	By above equality	
$[\Omega_0](\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha}) \vdash [[\Omega_0]\hat{\alpha}/\alpha][\Omega]A_0 \leq [\Omega]B$	By definition of substitution	
$[\Omega_0](\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha}) \vdash [[\Omega_0]\hat{\alpha}/\alpha][\Omega_0]A_0 \leq [\Omega_0]B$	By definition of substitution	
$[\Omega_0](\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha}) \vdash [\Omega_0][\hat{\alpha}/\alpha]A_0 \leq [\Omega_0]B$	By distributivity of substitution	
$\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha} \vdash [\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha}][\hat{\alpha}/\alpha]A_0 <: [\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha}]B \dashv \Delta_0$	By i.h.	
$\Delta_0 \longrightarrow \Omega''$	"	
$\Omega_0 \longrightarrow \Omega''$	"	
$\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha} \vdash [\Gamma][\hat{\alpha}/\alpha]A_0 <: [\Gamma]B \dashv \Delta_0$	By definition of substitution	
$\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha} \longrightarrow \Delta_0$	By Lemma 33 (Subtyping Extension)	
$\Delta_0 = (\Delta, \blacktriangleright_{\hat{\alpha}}, \Theta)$	By Lemma 24 (Extension Order) (ii)	
$\Gamma \longrightarrow \Delta$	"	
$\Omega'' = (\Omega', \blacktriangleright_{\hat{\alpha}}, \Omega_Z)$	By Lemma 24 (Extension Order) (ii)	
$\Delta \longrightarrow \Omega'$	"	
$\Omega_0 \longrightarrow \Omega''$	Above	
$\Omega, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha} = \tau \longrightarrow \Omega', \blacktriangleright_{\hat{\alpha}}, \Omega_Z$	By above equalities	
$\Omega \longrightarrow \Omega'$	By Lemma 24 (Extension Order) (ii)	
$\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha} \vdash [\Gamma][\hat{\alpha}/\alpha]A_0 <: [\Gamma]B \dashv \Delta, \blacktriangleright_{\hat{\alpha}}, \Theta$	By above equality $\Delta_0 = (\Delta, \blacktriangleright_{\hat{\alpha}}, \Theta)$	
$\Gamma, \blacktriangleright_{\hat{\alpha}}, \hat{\alpha} \vdash [\hat{\alpha}/\alpha][\Gamma]A_0 <: [\Gamma]B \dashv \Delta, \blacktriangleright_{\hat{\alpha}}, \Theta$	By def. of subst. ($[\Gamma]\hat{\alpha} = \hat{\alpha}$ and $[\Gamma]\alpha = \alpha$)	
$\Gamma \vdash \forall\alpha. [\Gamma]A_0 <: [\Gamma]B \dashv \Delta$	By $<: \forall L$	
$\Gamma \vdash \forall\alpha. A' <: [\Gamma]B \dashv \Delta$	By above equality	

– **Case 2.AEx: A is an existential variable $[\Gamma]A = \hat{\alpha}$:**

We split on the form of $[\Gamma]B$.

* **Case 2.AEx.SameEx: $[\Gamma]B$ is the same existential variable $[\Gamma]B = \hat{\alpha}$:**

$$\begin{array}{ll}
\Gamma \vdash \hat{\alpha} <: \hat{\alpha} \dashv \Gamma & \text{By } <: \text{Exvar} \\
\text{☞} \quad \Gamma \vdash [\Gamma]A <: [\Gamma]B \dashv \Gamma & \text{By } [\Gamma]A = [\Gamma]B = \hat{\alpha} \\
\text{☞} \quad \Delta \longrightarrow \Omega & \Delta = \Gamma \\
\text{☞} \quad \Omega \longrightarrow \Omega' & \text{By Lemma 20 (Reflexivity) and } \Omega' = \Omega
\end{array}$$

* **Case 2.AEx.OtherEx:** $[\Gamma]B$ is a different existential variable $[\Gamma]B = \hat{\beta}$ where $\hat{\beta} \neq \hat{\alpha}$:
Either $\hat{\alpha} \in \text{FV}([\Gamma]\hat{\beta})$, or $\hat{\alpha} \notin \text{FV}([\Gamma]\hat{\beta})$.

· $\hat{\alpha} \in \text{FV}([\Gamma]\hat{\beta})$:

We have $\hat{\alpha} \preceq [\Gamma]\hat{\beta}$.

Therefore $\hat{\alpha} = [\Gamma]\hat{\beta}$, or $\hat{\alpha} \prec [\Gamma]\hat{\beta}$.

But we are in Case 2.AEx.OtherEx, so the former is impossible.

Therefore, $\hat{\alpha} \prec [\Gamma]\hat{\beta}$.

Since Γ is predicative, $[\Gamma]\hat{\beta}$ cannot have the form $\forall\beta. \dots$, so the only way that $\hat{\alpha}$ can be a proper subterm of $[\Gamma]\hat{\beta}$ is if $[\Gamma]\hat{\beta}$ has the form $B_1 \rightarrow B_2$ such that $\hat{\alpha}$ is a subterm of B_1 or B_2 , that is: $\hat{\alpha} \prec [\Gamma]\hat{\beta}$.

Then by a property of substitution, $[\Omega]\hat{\alpha} \prec [\Omega][\Gamma]\hat{\beta}$.

By Lemma 18 (Substitution Extension Invariance), $[\Omega][\Gamma]\hat{\beta} = [\Omega]\hat{\beta}$, so $[\Omega]\hat{\alpha} \prec [\Omega]\hat{\beta}$.

We have $[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq [\Omega]\hat{\beta}$, and we know that $[\Omega]\hat{\alpha}$ is a monotype, so we can use Lemma 8 (Occurrence) (ii) to show that $[\Omega]\hat{\alpha} \not\prec [\Omega]\hat{\beta}$, a contradiction.

· $\hat{\alpha} \notin \text{FV}([\Gamma]\hat{\beta})$:

$$\Gamma \vdash \hat{\alpha} : \leq [\Gamma]\hat{\beta} \dashv \Delta \quad \text{By Theorem 13 (1)}$$

$$\text{☞} \quad \Gamma \vdash \hat{\alpha} <: \hat{\beta} \dashv \Delta \quad \text{By } <: \text{InstantiateL}$$

$$\text{☞} \quad \Delta \longrightarrow \Omega' \quad \text{"}$$

$$\text{☞} \quad \Omega \longrightarrow \Omega' \quad \text{"}$$

* **Case 2.AEx.Unit:** $[\Gamma]B = 1$:

$$\Gamma \longrightarrow \Omega \quad \text{Given}$$

$$1 = [\Omega]1 \quad \text{By definition of substitution}$$

$$\hat{\alpha} \notin \text{FV}(1) \quad \text{By definition of } \text{FV}(-)$$

$$[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq [\Omega]1 \quad \text{Given}$$

$$\Gamma \vdash \hat{\alpha} : \leq 1 \dashv \Delta \quad \text{By Theorem 13 (1)}$$

$$\text{☞} \quad \Omega \longrightarrow \Omega' \quad \text{"}$$

$$\text{☞} \quad \Delta \longrightarrow \Omega' \quad \text{"}$$

$$1 = [\Gamma]1 \quad \text{By definition of substitution}$$

$$\hat{\alpha} \notin \text{FV}(1) \quad \text{By definition of } \text{FV}(-)$$

$$\text{☞} \quad \Gamma \vdash \hat{\alpha} <: 1 \dashv \Delta \quad \text{By } <: \text{InstantiateL}$$

* **Case 2.AEx.Uvar:** $[\Gamma]B = \beta$:

Similar to Case 2.AEx.Unit, using $\beta = [\Omega]\beta = [\Gamma]\beta$ and $\hat{\alpha} \notin \text{FV}(\beta)$.

* **Case 2.AEx.Arrow:** $[\Gamma]B = B_1 \rightarrow B_2$:

Since $[\Gamma]B$ is an arrow, it cannot be exactly $\hat{\alpha}$.

Suppose, for a contradiction, that $\hat{\alpha} \in \text{FV}([\Gamma]B)$.

$\hat{\alpha} \prec [\Gamma]B$	$\hat{\alpha} \in \text{FV}([\Gamma]B)$
$[\Omega]\hat{\alpha} \prec [\Omega][\Gamma]B$	By a property of substitution
$\Gamma \longrightarrow \Omega$	Given
$[\Omega][\Gamma]B = [\Omega]B$	By Lemma 18 (Substitution Extension Invariance)
$[\Omega]\hat{\alpha} \prec [\Omega]B$	By above equality
$[\Gamma]B \neq \hat{\alpha}$	Given (2.AEx.Arrow)
$[\Omega][\Gamma]B \neq [\Omega]\hat{\alpha}$	By a property of substitution
$[\Omega]B \neq [\Omega]\hat{\alpha}$	By Lemma 18 (Substitution Extension Invariance)
$[\Omega]\hat{\alpha} \prec [\Omega]B$	Follows from \prec and \neq
$[\Omega]\hat{\alpha} \succ [\Omega]B$	$[\Omega]A$ has the form $\dots \rightarrow \dots$
$[\Omega]\Gamma \vdash [\Omega]\hat{\alpha} \leq [\Omega]B$	Given
$[\Omega]B$ is a monotype	Ω is predicative
$[\Omega]\hat{\alpha} \not\prec [\Omega]B$	By Lemma 8 (Occurrence) (ii)
$\Rightarrow \Leftarrow$	
$\hat{\alpha} \notin \text{FV}([\Gamma]B)$	By contradiction
$\Gamma \vdash \hat{\alpha} : \leq [\Gamma]B \dashv \Delta$	By Theorem 13 (1)
$\Delta \longrightarrow \Omega'$	"
$\Omega \longrightarrow \Omega'$	"
$\Gamma \vdash \hat{\alpha} <: \underbrace{[\Gamma]B}_{B_1 \rightarrow B_2} \dashv \Delta$	By $<:$ InstantiateL

– **Case 2.BEx:** $[\Gamma]A$ is not polymorphic and $[\Gamma]B$ is an existential variable: $[\Gamma]B = \hat{\beta}$
We split on the form of $[\Gamma]A$.

- * **Case 2.BEx.Unit** ($[\Gamma]A = 1$),
- Case 2.BEx.Uvar** ($[\Gamma]A = \alpha$),
- Case 2.BEx.Arrow** ($[\Gamma]A = A_1 \rightarrow A_2$):

Similar to Cases **2.AEx.Unit**, **2.AEx.Uvar** and **2.AEx.Arrow**, but using part (2) of Theorem 13 instead of part (1), and applying $<:$ InstantiateR instead of $<:$ InstantiateL as the final step.

– **Case 2.Units:** $[\Gamma]A = [\Gamma]B = 1$:

$\Gamma \vdash 1 <: 1 \dashv \Gamma$	By $<:$ Unit
$\Gamma \longrightarrow \Omega$	Given
$\Delta \longrightarrow \Omega$	$\Delta = \Gamma$
$\Omega \longrightarrow \Omega'$	By Lemma 20 (Reflexivity) and $\Omega' = \Omega$

– **Case 2.Uvars:** $[\Gamma]A = [\Gamma]B = \alpha$:

$\alpha \in \Omega$	By inversion on \leq Var
$\Gamma \longrightarrow \Omega$	Given
$\alpha \in \Gamma$	By Lemma 24 (Extension Order)
$\Gamma \vdash \alpha <: \alpha \dashv \Gamma$	By $<:$ Var
$\Delta \longrightarrow \Omega$	$\Delta = \Gamma$
$\Omega \longrightarrow \Omega'$	By Lemma 20 (Reflexivity) and $\Omega' = \Omega$

– **Case 2.Arrows:** $A = A_1 \rightarrow A_2$ and $B = B_1 \rightarrow B_2$:

Only rule $\leq \rightarrow$ could have been used.

$[\Omega]\Gamma \vdash [\Omega]B_1 \leq [\Omega]A_1$	Subderivation
$\Gamma \vdash [\Gamma]B_1 <: [\Gamma]A_1 \dashv \Theta$	By i.h.
$\Theta \longrightarrow \Omega_0$	"
$\Omega \longrightarrow \Omega_0$	"
$\Gamma \longrightarrow \Omega$	Given
$\Gamma \longrightarrow \Omega_0$	By Lemma 21 (Transitivity)
$\Theta \longrightarrow \Omega_0$	Above
$[\Omega]\Gamma = [\Omega]\Theta$	By Lemma 52 (Confluence of Completeness)
$[\Omega]\Gamma \vdash [\Omega]A_2 \leq [\Omega]B_2$	Subderivation
$[\Omega]\Theta \vdash [\Omega]A_2 \leq [\Omega]B_2$	By above equality
$[\Omega]A_2 = [\Omega][\Gamma]A_2$	By Lemma 18 (Substitution Extension Invariance)
$[\Omega]B_2 = [\Omega][\Gamma]B_2$	By Lemma 18 (Substitution Extension Invariance)
$[\Omega]\Theta \vdash [\Omega][\Gamma]A_2 \leq [\Omega][\Gamma]B_2$	By above equalities
$\Theta \vdash [\Theta][\Gamma]A_2 <: [\Theta][\Gamma]B_2 \dashv \Delta$	By i.h.
$\Delta \longrightarrow \Omega'$	"
$\Omega_0 \longrightarrow \Omega'$	"
$\Gamma \vdash ([\Gamma]A_1) \rightarrow ([\Gamma]A_2) <: ([\Gamma]B_1) \rightarrow ([\Gamma]B_2) \dashv \Delta$	By $<: \rightarrow$
$\Gamma \vdash [\Gamma](A_1 \rightarrow A_2) <: [\Gamma](B_1 \rightarrow B_2) \dashv \Delta$	By definition of substitution
$\Omega \longrightarrow \Omega'$	By Lemma 21 (Transitivity) □

Corollary 55 (Completeness of Subtyping). *If $\Psi \vdash A \leq B$ then there is a Δ such that $\Psi \vdash A <: B \dashv \Delta$.*

Proof. Let $\Omega = \Psi$ and $\Gamma = \Psi$.

By Lemma 20 (Reflexivity), $\Psi \longrightarrow \Psi$, so $\Gamma \longrightarrow \Omega$.

By Lemma 4 (Well-Formedness), $\Psi \vdash A$ and $\Psi \vdash B$; since $\Gamma = \Psi$, we have $\Gamma \vdash A$ and $\Gamma \vdash B$.

By Theorem 14, there exists Δ such that $\Gamma \vdash [\Gamma]A <: [\Gamma]B \dashv \Delta$.

Since $\Gamma = \Psi$ and Ψ is a declarative context with no existentials, $[\Psi]C = C$ for all C , so we actually have $\Psi \vdash A <: B \dashv \Delta$, which was to be shown. □

L' Completeness of Typing

Theorem 15 (Completeness of Algorithmic Typing). *Given $\Gamma \longrightarrow \Omega$ and $\Gamma \vdash A$:*

- (i) *If $[\Omega]\Gamma \vdash e \Leftarrow [\Omega]A$
then there exist Δ and Ω'
such that $\Delta \longrightarrow \Omega'$ and $\Omega \longrightarrow \Omega'$ and $\Gamma \vdash e \Leftarrow [\Gamma]A \dashv \Delta$.*
- (ii) *If $[\Omega]\Gamma \vdash e \Rightarrow A$
then there exist Δ , Ω' , and A'
such that $\Delta \longrightarrow \Omega'$ and $\Omega \longrightarrow \Omega'$ and $\Gamma \vdash e \Rightarrow A' \dashv \Delta$ and $A = [\Omega']A'$.*
- (iii) *If $[\Omega]\Gamma \vdash [\Omega]A \bullet e \Rightarrow C$
then there exist Δ , Ω' , and C'
such that $\Delta \longrightarrow \Omega'$ and $\Omega \longrightarrow \Omega'$ and $\Gamma \vdash [\Gamma]A \bullet e \Rightarrow C' \dashv \Delta$ and $C = [\Omega']C'$.*

Proof. By induction on the given declarative derivation.

- **Case** $\frac{(x : A) \in [\Omega]\Gamma}{[\Omega]\Gamma \vdash x \Rightarrow A}$ DeclVar

$(x : A) \in [\Omega]\Gamma$	Premise
$\Gamma \longrightarrow \Omega$	Given
$(x : A') \in \Gamma$ where $[\Omega]A' = [\Omega]A$	From definition of context application
Let $\Delta = \Gamma$.	
Let $\Omega' = \Omega$.	
☞ $\Gamma \longrightarrow \Omega$	Given
☞ $\Omega \longrightarrow \Omega$	By Lemma 20 (Reflexivity)
☞ $\Gamma \vdash x \Rightarrow A' \dashv \Gamma$	By Var
$[\Omega]A' = [\Omega]A$	Above
$= A$	FEV(A) = \emptyset
- **Case** $\frac{[\Omega]\Gamma \vdash e \Rightarrow B \quad [\Omega]\Gamma \vdash B \leq [\Omega]A}{[\Omega]\Gamma \vdash e \Leftarrow [\Omega]A}$ DeclSub

$[\Omega]\Gamma \vdash e \Rightarrow B$	Subderivation
$\Gamma \vdash e \Rightarrow B' \dashv \Theta$	By i.h.
$B = [\Omega]B'$	"
$\Theta \longrightarrow \Omega_0$	"
$\Omega \longrightarrow \Omega_0$	"
$\Gamma \longrightarrow \Omega$	Given
$\Gamma \longrightarrow \Omega_0$	By Lemma 21 (Transitivity)
$[\Omega]\Gamma \vdash B \leq [\Omega]A$	Subderivation
$[\Omega]\Gamma = [\Omega]\Theta$	By Lemma 52 (Confluence of Completeness)
$[\Omega]\Theta \vdash B \leq [\Omega]A$	By above equalities
$\Theta \longrightarrow \Omega_0$	Above
$\Theta \vdash [\Theta]B' \prec: [\Theta]A \dashv \Delta$	By Theorem 14
$\Delta \longrightarrow \Omega'$	"
$\Omega_0 \longrightarrow \Omega'$	"
☞ $\Delta \longrightarrow \Omega'$	By Lemma 21 (Transitivity)
☞ $\Omega \longrightarrow \Omega'$	By Lemma 21 (Transitivity)
☞ $\Gamma \vdash e \Leftarrow A \dashv \Delta$	By Sub

- **Case**
$$\frac{[\Omega]\Gamma \vdash A \quad [\Omega]\Gamma \vdash e_0 \Leftarrow A}{[\Omega]\Gamma \vdash (e_0 : A) \Rightarrow A} \text{DeclAnno}$$

$A = [\Omega]A$	Source type annotations cannot contain evars
$= [\Gamma]A$	"
$[\Omega]\Gamma \vdash e_0 \Leftarrow A$	Subderivation
$[\Omega]\Gamma \vdash e_0 \Leftarrow [\Omega]A$	By above equality
$\Gamma \vdash e_0 \Leftarrow [\Gamma]A \dashv \Delta$	By i.h.
$\Delta \longrightarrow \Omega$	"
$\Omega \longrightarrow \Omega'$	"
$\Gamma \vdash A$	Given
$\Gamma \vdash (e_0 : A) \Rightarrow A \dashv \Delta$	By Anno
$A = [\Omega']A$	Source type annotations cannot contain evars
$\Gamma \vdash (e_0 : [\Omega']A) \Rightarrow [\Omega']A \dashv \Delta$	By above equality

- **Case**

$$\frac{}{[\Omega]\Gamma \vdash () \Leftarrow 1} \text{Decl11}$$

We have $[\Omega]A = 1$. Either $[\Gamma]A = 1$ or $[\Gamma]A = \hat{\alpha} \in \text{unsolved}(\Gamma)$.

In the former case:

Let $\Delta = \Gamma$.	
Let $\Omega' = \Omega$.	
$\Gamma \longrightarrow \Omega$	Given
$\Omega \longrightarrow \Omega'$	By Lemma 20 (Reflexivity)
$\Gamma \vdash () \Leftarrow 1 \dashv \Gamma$	By 11
$\Gamma \vdash () \Leftarrow [\Gamma]1 \dashv \Gamma$	$1 = [\Gamma]1$

In the latter case:

$\Gamma \vdash () \Rightarrow 1 \dashv \Gamma$	By 11 \Rightarrow
$[\Omega]\Gamma \vdash 1 \leq 1$	By $\leq \text{Unit}$
$1 = [\Omega]1$	By definition of substitution
$= [\Omega][\Gamma]\hat{\alpha}$	By $[\Omega]A = 1$
$= [\Omega]\hat{\alpha}$	By Lemma 18 (Substitution Extension Invariance)
$[\Omega]\Gamma \vdash [\Omega]1 \leq [\Omega]\hat{\alpha}$	By above equalities
$\Gamma \vdash 1 <: \hat{\alpha} \dashv \Delta$	By Theorem 13 (1)
$1 = [\Gamma]1$	By definition of substitution
$\hat{\alpha} = [\Gamma]\hat{\alpha}$	$\hat{\alpha} \in \text{unsolved}(\Gamma)$
$\Gamma \vdash [\Gamma]1 <: [\Gamma]\hat{\alpha} \dashv \Delta$	By above equalities
$\Omega \longrightarrow \Omega'$	"
$\Delta \longrightarrow \Omega'$	"
$\Gamma \vdash () \Leftarrow \hat{\alpha} \dashv \Delta$	By Sub
$\Gamma \vdash () \Leftarrow [\Gamma]A \dashv \Delta$	By $[\Gamma]A = \hat{\alpha}$

- **Case**

$$\frac{[\Omega]\Gamma, \alpha \vdash e \Leftarrow A_0}{[\Omega]\Gamma \vdash e \Leftarrow \forall \alpha. A_0} \text{DeclVI}$$

$[\Omega]A = \forall\alpha. A_0$	Given
$= \forall\alpha. [\Omega]A'$	By def. of subst. and predicativity of Ω
$A_0 = [\Omega]A'$	Follows from above equality
$[\Omega]\Gamma, \alpha \vdash e \Leftarrow [\Omega]A'$	Subderivation and above equality
$\Gamma \longrightarrow \Omega$	Given
$\Gamma, \alpha \longrightarrow \Omega, \alpha$	By \longrightarrow Uvar
$[\Omega]\Gamma, \alpha = [\Omega, \alpha](\Gamma, \alpha)$	By definition of context substitution
$[\Omega, \alpha](\Gamma, \alpha) \vdash e \Leftarrow [\Omega]A'$	By above equality
$[\Omega, \alpha](\Gamma, \alpha) \vdash e \Leftarrow [\Omega, \alpha]A'$	By definition of substitution
$\Gamma, \alpha \vdash e \Leftarrow [\Gamma, \alpha]A' \dashv \Delta'$	By i.h.
$\Delta' \longrightarrow \Omega'_0$	"
$\Omega, \alpha \longrightarrow \Omega'_0$	"
$\Gamma, \alpha \longrightarrow \Delta'$	By Lemma 54 (Typing Extension)
$\Delta' = \Delta, \alpha, \Theta$	By Lemma 24 (Extension Order) (i)
$\Delta, \alpha, \Theta \longrightarrow \Omega'_0$	By above equality
$\Omega'_0 = \Omega', \alpha, \Omega_Z$	By Lemma 24 (Extension Order) (i)
☞ $\Delta \longrightarrow \Omega'$	"
☞ $\Omega \longrightarrow \Omega'$	By Lemma 24 (Extension Order) on $\Omega, \alpha \longrightarrow \Omega'_0$
$\Gamma, \alpha \vdash e \Leftarrow [\Gamma, \alpha]A' \dashv \Delta, \alpha, \Theta$	By above equality
$\Gamma, \alpha \vdash e \Leftarrow [\Gamma]A' \dashv \Delta, \alpha, \Theta$	By definition of substitution
$\Gamma \vdash e \Leftarrow \forall\alpha. [\Gamma]A' \dashv \Delta$	By \forall I
☞ $\Gamma \vdash e \Leftarrow [\Gamma](\forall\alpha. A') \dashv \Delta$	By definition of substitution

• **Case**
$$\frac{[\Omega]\Gamma \vdash \tau \quad [\Omega]\Gamma \vdash [\tau/\alpha]A_0 \bullet e \Rightarrow C}{[\Omega]\Gamma \vdash \underbrace{\forall\alpha. A_0}_{[\Omega]A} \bullet e \Rightarrow C} \text{Decl}\forall\text{App}$$

$[\Omega]\Gamma \vdash \tau$	Subderivation
$[\Omega]A = \forall\alpha. A_0$	Given
$= \forall\alpha. [\Omega]A'$	By def. of subst. and predicativity of Ω
$[\Omega]\Gamma \vdash [\tau/\alpha][\Omega]A' \bullet e \Rightarrow C$	Subderivation and above equality
$\Gamma \longrightarrow \Omega$	Given
$\Gamma, \hat{\alpha} \longrightarrow \Omega, \hat{\alpha} = \tau$	By \longrightarrow Solve
$[\Omega]\Gamma = [\Omega, \hat{\alpha} = \tau](\Gamma, \hat{\alpha})$	By definition of context application
$[\Omega, \hat{\alpha} = \tau](\Gamma, \hat{\alpha}) \vdash [\tau/\alpha][\Omega]A' \bullet e \Rightarrow C$	By above equality
$[\Omega, \hat{\alpha} = \tau](\Gamma, \hat{\alpha}) \vdash [\tau/\alpha][\Omega, \hat{\alpha} = \tau]A' \bullet e \Rightarrow C$	By def. of subst.
$([[\Omega]\tau/\alpha][\Omega, \hat{\alpha} = \tau]A') = ([\Omega, \hat{\alpha} = \tau][\hat{\alpha}/\alpha]A')$	By distributivity of substitution
$\tau = [\Omega]\tau$	$\text{FEV}(\tau) = \emptyset$
$([\tau/\alpha][\Omega, \hat{\alpha} = \tau]A') = ([\Omega, \hat{\alpha} = \tau][\hat{\alpha}/\alpha]A')$	By above equality
$[\Omega, \hat{\alpha} = \tau](\Gamma, \hat{\alpha}) \vdash [\Omega, \hat{\alpha} = \tau][\hat{\alpha}/\alpha]A' \bullet e \Rightarrow C$	By above equality
☞ $\Gamma, \hat{\alpha} \vdash [\hat{\alpha}/\alpha]A' \bullet e \Rightarrow C' \dashv \Delta$	By i.h.
☞ $C = [\Omega]C'$	"
☞ $\Delta \longrightarrow \Omega'$	"
☞ $\Omega \longrightarrow \Omega'$	"
☞ $\Gamma \vdash \forall\alpha. A' \bullet e \Rightarrow C' \dashv \Delta$	By \forall App

• **Case**
$$\frac{[\Omega]\Gamma, x : A'_1 \vdash e_0 \Leftarrow A'_2}{[\Omega]\Gamma \vdash \lambda x. e_0 \Leftarrow A'_1 \rightarrow A'_2} \text{Decl} \rightarrow \text{I}$$

We have $[\Omega]A = A'_1 \rightarrow A'_2$. Either $[\Gamma]A = A_1 \rightarrow A_2$ where $A'_1 = [\Omega]A_1$ and $A'_2 = [\Omega]A_2$ —or $[\Gamma]A = \hat{\alpha}$ and $[\Omega]\hat{\alpha} = A'_1 \rightarrow A'_2$.

In the former case:

$[\Omega]\Gamma, x : A'_1 \vdash e_0 \Leftarrow A'_2$	Subderivation
$A'_1 = [\Omega]A_1$	Known in this subcase
$= [\Omega][\Gamma]A_1$	By Lemma 18 (Substitution Extension Invariance)
$[\Omega]A'_1 = [\Omega][\Omega][\Gamma]A_1$	Applying Ω on both sides
$= [\Omega][\Gamma]A_1$	By idempotence of substitution
$[\Omega]\Gamma, x : A'_1 = [\Omega, x : A'_1](\Gamma, x : [\Gamma]A_1)$	By definition of context application
$[\Omega, x : A'_1](\Gamma, x : [\Gamma]A_1) \vdash e_0 \Leftarrow A'_2$	By above equality
$\Gamma \longrightarrow \Omega$	Given
$\Gamma, x : [\Gamma]A_1 \longrightarrow \Omega, x : A'_1$	By \longrightarrow Var
$\Gamma, x : [\Gamma]A_1 \vdash e_0 \Leftarrow A_2 \dashv \Delta'$	By i.h.
$\Delta' \longrightarrow \Omega'_0$	"
$\Omega, x : A'_1 \longrightarrow \Omega'_0$	"
$\Omega'_0 = \Omega', x : A'_1, \Theta$	By Lemma 24 (Extension Order) (v)
$\Omega \longrightarrow \Omega'$	"
$\Gamma, x : [\Gamma]A_1 \longrightarrow \Delta'$	By Lemma 54 (Typing Extension)
$\Delta' = \Delta, x : \dots, \Theta$	By Lemma 24 (Extension Order) (v)
$\Delta, x : \dots, \Theta \longrightarrow \Omega', x : A'_1, \Theta$	By above equalities
$\Delta \longrightarrow \Omega'$	By Lemma 24 (Extension Order) (v)
$\Gamma, x : [\Gamma]A_1 \vdash e_0 \Leftarrow [\Gamma]A_2 \dashv \Delta, \alpha, \Theta$	By above equality
$\Gamma \vdash \lambda x. e_0 \Leftarrow ([\Gamma]A_1) \rightarrow ([\Gamma]A_2) \dashv \Delta$	By \rightarrow I
$\Gamma \vdash \lambda x. e_0 \Leftarrow [\Gamma](A_1 \rightarrow A_2) \dashv \Delta$	By definition of substitution

In the latter case:

$[\Omega]\hat{\alpha} = A'_1 \rightarrow A'_2$	Known in this subcase
$[\Omega]\Gamma, x : A'_1 \vdash e_0 \Leftarrow A'_2$	Subderivation
$\Gamma \longrightarrow \Omega$	Given
$\Gamma, \hat{\alpha}, \hat{\beta} \longrightarrow \Omega, \hat{\alpha} = A'_1, \hat{\beta} = A'_2$	By \longrightarrow Solve twice
$[\Omega]\hat{\alpha} = [\Omega]A'_1$	By definition of substitution
$\Gamma, \hat{\alpha}, \hat{\beta}, x : \hat{\alpha} \longrightarrow \Omega, \hat{\alpha} = A'_1, \hat{\beta} = A'_2, x : A'_1$	By \longrightarrow Var
$[\Omega]\Gamma, x : A'_1 = [\Omega, \hat{\alpha} = A'_1, \hat{\beta} = A'_2, x : A'_1](\Gamma, \hat{\alpha}, \hat{\beta}, x : \hat{\alpha})$	By definition of context application
Let $\Omega_0 = (\Omega, \hat{\alpha} = A'_1, \hat{\beta} = A'_2, x : A'_1)$.	
$[\Omega_0](\Gamma, \hat{\alpha}, \hat{\beta}, x : \hat{\alpha}) \vdash e_0 \Leftarrow A'_2$	By above equality
$\Gamma, \hat{\alpha}, \hat{\beta}, x : \hat{\alpha} \vdash e_0 \Leftarrow \hat{\beta} \dashv \Delta'$	By i.h. with Ω_0
$\Delta' \longrightarrow \Omega'_0$	"
$\Omega_0 \longrightarrow \Omega'_0$	"

	$\Gamma, \hat{\alpha}, \hat{\beta}, x : \hat{\alpha} \longrightarrow \Delta'$	By Lemma 54 (Typing Extension)
	$\Delta' = \Delta, x : \hat{\alpha}, \Theta$	By Lemma 24 (Extension Order) (v)
	$\Delta, x : \hat{\alpha}, \Theta \longrightarrow \Omega'_0$	By above equality
	$\Omega'_0 = \Omega'', x : \dots, \Omega_Z$	By Lemma 54 (Typing Extension)
☞	$\Delta \longrightarrow \Omega''$	"
	$\Gamma, \hat{\alpha}, \hat{\beta} \longrightarrow \Delta$	"
	$\Omega_0 \longrightarrow \underbrace{\Omega'', x : \dots, \Omega_Z}_{\Omega'_0}$	By above equality
	$\Omega, \hat{\alpha} = A'_1, \hat{\beta} = A'_2, x : A'_1 \longrightarrow \Omega'', x : \dots, \Omega_Z$	By def. of Ω_0
	$\Omega'' = \Omega', \hat{\alpha} = \dots, \dots$	By Lemma 24 (Extension Order) (iii)
☞	$\Omega \longrightarrow \Omega'$	"
	$\Gamma, \hat{\alpha}, \hat{\beta}, x : \hat{\alpha} \vdash e_0 \leftarrow \hat{\beta} \dashv \Delta, x : \hat{\alpha}, \Theta$	By above equality
	$\Gamma \vdash \lambda x. e_0 \leftarrow \hat{\alpha} \rightarrow \hat{\beta} \dashv \Delta$	By $\rightarrow \vdash \Rightarrow$
	$[\Gamma]\hat{\alpha} = \hat{\alpha}$	By definition of substitution
	$[\Gamma]\hat{\beta} = \hat{\beta}$	By definition of substitution
	$\Gamma \vdash \lambda x. e_0 \leftarrow ([\Gamma]\hat{\alpha}) \rightarrow ([\Gamma]\hat{\beta}) \dashv \Delta$	By above equalities
☞	$\Gamma \vdash \lambda x. e_0 \leftarrow [\Gamma](\hat{\alpha} \rightarrow \hat{\beta}) \dashv \Delta$	By definition of substitution

• **Case** $\frac{[\Omega]\Gamma \vdash e_1 \Rightarrow B \quad [\Omega]\Gamma \vdash B \bullet e_2 \Rightarrow A}{[\Omega]\Gamma \vdash e_1 e_2 \Rightarrow A} \text{Decl} \rightarrow E$

	$[\Omega]\Gamma \vdash e_1 \Rightarrow B$	Subderivation
	$\Gamma \longrightarrow \Omega$	Given
	$\Gamma \vdash e_1 \Rightarrow B' \dashv \Theta$	By i.h.
	$B = [\Omega]B'$	"
	$\Theta \longrightarrow \Omega'_0$	"
	$\Omega \longrightarrow \Omega'_0$	"
	$[\Omega]\Gamma \vdash B \bullet e_2 \Rightarrow A$	Subderivation
	$[\Omega]\Gamma \vdash [\Omega]B' \bullet e_2 \Rightarrow A$	By above equality
	$\Gamma \longrightarrow \Omega'_0$	By Lemma 21 (Transitivity)
	$[\Omega]\Gamma = [\Omega]\Omega$	By Lemma 49 (Stability of Complete Contexts)
	$= [\Omega'_0]\Omega'_0$	By Lemma 51 (Finishing Completions)
	$= [\Omega'_0]\Gamma$	By Lemma 49 (Stability of Complete Contexts)
	$= [\Omega'_0]\Theta$	By Lemma 52 (Confluence of Completeness)
	$[\Omega'_0]\Theta \vdash [\Omega]B' \bullet e_2 \Rightarrow A$	By above equality
	$[\Omega]B' = [\Omega'_0]B'$	By Lemma 50 (Finishing Types)
	$[\Omega'_0]B' = [\Omega'_0][\Theta]B'$	By Lemma 18 (Substitution Extension Invariance)
	$[\Omega'_0]\Theta \vdash [\Omega][\Theta]B' \bullet e_2 \Rightarrow A$	By above equalities
	$\Theta \vdash [\Theta]B' \bullet e_2 \Rightarrow A' \dashv \Delta$	By i.h. with Ω'_0
☞	$A = [\Omega]A'$	"
☞	$\Delta \longrightarrow \Omega'$	"
	$\Omega'_0 \longrightarrow \Omega'$	"
	$\Omega \longrightarrow \Omega'$	By Lemma 21 (Transitivity)
☞	$\Gamma \vdash e_1 e_2 \Rightarrow A' \dashv \Delta$	By $\rightarrow E$

• **Case**

$$\frac{[\Omega]\Gamma \vdash e \Leftarrow B}{[\Omega]\Gamma \vdash \underbrace{B \rightarrow C}_{[\Omega]A} \bullet e \Rightarrow C} \text{Decl} \rightarrow \text{App}$$

We have $[\Omega]A = B \rightarrow C$. Either $[\Gamma]A = B_0 \rightarrow C_0$ where $B = [\Omega]B_0$ and $C = [\Omega]C_0$ —or $[\Gamma]A = \hat{\alpha}$ where $\hat{\alpha} \in \text{unsolved}(\Gamma)$ and $[\Omega]\hat{\alpha} = B \rightarrow C$.

In the former case:

$[\Omega]\Gamma \vdash e \Leftarrow B$	Subderivation
$B = [\Omega]B_0$	Known in this subcase
$\Gamma \longrightarrow \Omega$	Given
$\Gamma \vdash e \Leftarrow [\Gamma]B_0 \dashv \Delta$	By i.h.
$\Gamma \vdash ([\Gamma]B_0) \rightarrow ([\Gamma]C_0) \bullet e \Rightarrow [\Gamma]C_0 \dashv \Delta$	By \rightarrow App
☞ $\Delta \longrightarrow \Omega'$	"
☞ $\Omega \longrightarrow \Omega'$	"
Let $C' = [\Gamma]C_0$.	
$C = [\Omega]C_0$	Known in this subcase
$= [\Omega][\Gamma]C_0$	By Lemma 18 (Substitution Extension Invariance)
☞ $= [\Omega]C'$	$[\Gamma]C_0 = C'$
☞ $\Gamma \vdash [\Gamma](B_0 \rightarrow C_0) \bullet e \Rightarrow [\Gamma]C_0 \dashv \Delta$	By definition of substitution

In the latter case, $\hat{\alpha} \in \text{unsolved}(\Gamma)$, so the context Γ must have the form $\Gamma_0[\hat{\alpha}]$.

$\Gamma \longrightarrow \Omega$	Given
$\Gamma_0[\hat{\alpha}] \longrightarrow \Omega$	$\Gamma = \Gamma_0[\hat{\alpha}]$
$[\Omega]A = B \rightarrow C$	Above
$[\Omega]\hat{\alpha} = B \rightarrow C$	$A = \hat{\alpha}$
$\Omega = \Omega_0[\hat{\alpha} = A_0]$ and $[\Omega]A_0 = B \rightarrow C$	Follows from $[\Omega]\hat{\alpha} = B \rightarrow C$
Let $\Gamma' = \Gamma_0[\hat{\alpha}_2, \hat{\alpha}_1, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2]$.	
Let $\Omega'_0 = \Omega_0[\hat{\alpha}_2 = [\Omega]C, \hat{\alpha}_1 = [\Omega]B, \hat{\alpha} = \hat{\alpha}_1 \rightarrow \hat{\alpha}_2]$.	
$\Gamma' \longrightarrow \Omega'_0$	By Lemma 29 (Parallel Admissibility) (ii) twice
$[\Omega]\Gamma \vdash e \Leftarrow B$	Subderivation
$\Omega \longrightarrow \Omega'_0$	By Lemma 27 (Solved Variable Addition for Extension) then Lemma 29 (Parallel Admissibility) (iii)
$[\Omega]\Gamma = [\Omega]\Omega$	By Lemma 49 (Stability of Complete Contexts)
$= [\Omega'_0]\Omega'_0$	By Lemma 51 (Finishing Completions)
$= [\Omega'_0]\Gamma'$	By Lemma 52 (Confluence of Completeness)
$B = [\Omega'_0]\hat{\alpha}_1$	By definition of Ω'_0
$[\Omega'_0]\Gamma' \vdash e \Leftarrow [\Omega'_0]\hat{\alpha}_1$	By above equalities
$\Gamma' \vdash e \Leftarrow [\Gamma']\hat{\alpha}_1 \dashv \Delta$	By i.h.
☞ $\Delta \longrightarrow \Omega'$	"
☞ $\Omega'_0 \longrightarrow \Omega'$	"
☞ $\Omega \longrightarrow \Omega'$	By Lemma 21 (Transitivity)
$[\Gamma']\hat{\alpha}_1 = \hat{\alpha}_1$	$\hat{\alpha}_1 \in \text{unsolved}(\Gamma')$
$\Gamma' \vdash e \Leftarrow \hat{\alpha}_1 \dashv \Delta$	By above equality

$$\begin{array}{l}
\Gamma \vdash \hat{\alpha} \bullet e \Rightarrow \hat{\alpha}_2 \dashv \Delta \quad \text{By } \hat{\alpha}\text{App} \\
\text{Let } C' = \hat{\alpha}_2. \\
C = [\Omega_0'] \hat{\alpha}_2 \quad \text{By definition of } \Omega_0' \\
= [\Omega'] \hat{\alpha}_2 \quad \text{By Lemma 50 (Finishing Types)} \\
\Rightarrow = [\Omega'] C' \quad \text{By above equality} \\
\Rightarrow \Gamma \vdash [\Gamma] A \bullet e \Rightarrow C' \dashv \Delta \quad \hat{\alpha} = [\Gamma] A \text{ and } \hat{\alpha}_2 = C'
\end{array}$$

• Case

$$\begin{array}{l}
\frac{}{[\Omega] \Gamma \vdash () \Rightarrow 1} \text{Decl}1\Rightarrow \\
1 = A \quad \text{Given} \\
\Gamma \vdash () \Rightarrow 1 \dashv \Gamma \quad \text{By } 1\Rightarrow \\
\text{Let } \Delta = \Gamma. \\
\text{Let } \Omega' = \Omega. \\
\Gamma \longrightarrow \Omega \quad \text{Given} \\
\Rightarrow \Delta \longrightarrow \Omega \quad \text{By above equality} \\
\Rightarrow \Omega \longrightarrow \Omega' \quad \text{By Lemma 20 (Reflexivity)} \\
\text{Let } A' = 1. \\
\Rightarrow \Gamma \vdash () \Rightarrow A' \dashv \Delta \quad \text{By above equalities} \\
\Rightarrow 1 = [\Omega] A' \quad \text{By definition of substitution}
\end{array}$$

• Case

$$\begin{array}{l}
\frac{[\Omega] \Gamma \vdash \sigma \rightarrow \tau \quad [\Omega] \Gamma, x : \sigma \vdash e_0 \Leftarrow \tau}{[\Omega] \Gamma \vdash \lambda x. e_0 \Rightarrow \sigma \rightarrow \tau} \text{Decl} \rightarrow \Leftarrow \\
(\sigma \rightarrow \tau) = A \quad \text{Given} \\
[\Omega] \Gamma, x : \sigma \vdash e_0 \Leftarrow \tau \quad \text{Subderivation} \\
\text{Let } \Gamma' = (\Gamma, \hat{\alpha}, \hat{\beta}, x : \hat{\alpha}). \\
\text{Let } \Omega_0 = (\Omega, \hat{\alpha} = \sigma, \hat{\beta} = \tau, x : \sigma). \\
\Gamma \longrightarrow \Omega \quad \text{Given} \\
\Gamma' \longrightarrow \Omega_0 \quad \text{By } \longrightarrow\text{Solve twice, then } \longrightarrow\text{Var} \\
[\Omega_0] \Gamma' = ([\Omega] \Gamma, x : \sigma) \quad \text{By definition of context application} \\
\tau = [\Omega_0] \hat{\beta} \quad \text{By definition of } \Omega_0 \\
[\Omega_0] \Gamma' \vdash e_0 \Leftarrow [\Omega_0] \hat{\beta} \quad \text{By above equalities} \\
\Gamma' \vdash e_0 \Leftarrow \hat{\beta} \dashv \Delta' \quad \text{By i.h.} \\
\Delta' \longrightarrow \Omega_0' \quad \text{"} \\
\Omega_0 \longrightarrow \Omega_0' \quad \text{"} \\
\Delta' = (\Delta, x : \hat{\alpha}, \Theta) \quad \text{By Lemma 24 (Extension Order) (v)} \\
\Gamma, \hat{\alpha}, \hat{\beta}, x : \hat{\alpha} \vdash e_0 \Leftarrow \hat{\beta} \dashv \Delta, x : \hat{\alpha}, \Theta \quad \text{By above equalities} \\
(\Delta, x : \hat{\alpha}, \Theta) \longrightarrow \Omega_0' \quad \text{By above equality} \\
\Omega_0' = \Omega', x : \sigma, \Omega_Z \quad \text{By Lemma 24 (Extension Order) (v)} \\
\Rightarrow \Delta \longrightarrow \Omega' \quad \text{"} \\
\Rightarrow \Gamma \vdash \lambda x. e_0 \Rightarrow \hat{\alpha} \rightarrow \hat{\beta} \dashv \Delta \quad \text{By } \rightarrow \Leftarrow \Rightarrow
\end{array}$$

Let $A' = (\hat{\alpha} \rightarrow \hat{\beta})$.

$\Gamma \vdash \lambda x. e_0 \Rightarrow A' \dashv \Delta$	By above equality
$\sigma \rightarrow \tau = ([\Omega_0]\hat{\alpha}) \rightarrow ([\Omega_0]\hat{\beta})$	By definition of Ω_0
$\sigma \rightarrow \tau = [\Omega_0](\hat{\alpha} \rightarrow \hat{\beta})$	By definition of substitution
$A = [\Omega_0]A'$	By above equalities
$A = [\Omega']A'$	By Lemma 50 (Finishing Types)
$\Gamma' \longrightarrow \Delta'$	By Lemma 54 (Typing Extension)
$\Omega \longrightarrow \Omega'$	By Lemma 21 (Transitivity) \square

References

Frank Pfenning. Structural cut elimination. In *LICS*, 1995.